

PESQUISAS

BOTÂNICA, Nº 36

Ano 1984

BALDUÍNO RAMBO, S.J.

GRAMINEAE RIOGRANDENSES

Instituto Anchietano de Pesquisas
São Leopoldo - Praça Tiradentes, 35 - Rio Grande do Sul - Brasil

PESQUISAS

BOTÂNICA, Nº 36

Ano 1984

BALDUÍNO RAMBO, S.J.

GRAMINEAE RIOGRANDENSES

Instituto Anchietano de Pesquisas
São Leopoldo - Praça Tiradentes, 35 - Rio Grande do Sul - Brasil

Edição financiada pela
Universidade do Vale do Rio dos Sinos – UNISINOS

APRESENTAÇÃO

O alentado trabalho “Gramineae Riograndenses” que B. Rambo deixou em forma de manuscrito é resultado de exaustiva pesquisa de campo e laboratório sobre umas das maiores e, sem dúvida, a mais importante das famílias de plantas do Estado do Rio Grande do Sul. O farto material estudado e as preciosas anotações de habitat, deram oportunidade ao Padre Balduino Rambo de apresentar um precioso trabalho de cunho taxonômico e ecológico.

Somos de opinião que este rico depositário de informações agrostológicas deva ser publicado, o que honrará a entidade que o acolher. Desta forma um público bem maior terá ao seu alcance a enorme experiência de Rambo refletida neste trabalho.

Como todo e qualquer trabalho científico, este também apresenta lacunas que pretendemos analisar.

1 – O trabalho não apresentou chaves para os gêneros e espécies, constante que se repete em todos os trabalhos publicados deste renomado botânico gaúcho.

2 – Rambo não é cioso em usar a nomenclatura atualizada dos nomes científicos de acordo com o Código Internacional de Nomenclatura Botânica. Acresce que muita coisa mudou após sua morte prematura.

Verificamos que diversos binômios citados, hoje jazem no vasto cemitério da sinonímia. Rambo cultivava um receio por criação de “espécies novas”, e por isso, “forçava” o enquadramento de plantas em taxa já publicados, que hoje reconhecidamente são boas espécies aceitas.

Para exemplificar aqui vai uma lista parcial de binômios atualizados:

- *Agrostis ramboi* Parodi = *Agrostis hygrometrica* Nees
- *Andropogon virgatus* (Spreng.) Nees = *Hypogynum virgatum* (Desvaux) Dandy
- *Arthrostylidium trinii* (Rupr.) Munro = *Chusquea mimosa* McClure & Smith
- *Briza calotheca* (Trin.) Hack. = *Poidium calotheca* (Trinii) Matthei
- *Briza juergensii* Hack. = *Poidium juergensii* (Hack.) Matthei

- *Briza lamarckiana* Nees = *Chascolytrum lamarckianum* (Nees) Matthei
- *Briza poaemorpha* (Presl) Heunr. = *Poidium poaemorphum* (Presl.) Matthei
- *Briza rufa* (Presl) Steud. = *Poidium rufum* (Presl) Matthei
- *Briza stricta* (Hoock. et Arn.) Steud = *Poidium monandrum* (Hackel) Matthei
- *Briza subarista* Lam. = *Chascolytrum subaristatum* (Lamarck) Desvaux
- *Briza triloba* Nees = *Chascolytrum subaristatum* (Lamarck) Desvaux
- *Briza uniolae* Nees = *Poidium uniolae* (Nees) Matthei
- *Cortaderia rudiuscula* Stapf = *Cortadeira vaginata* Swallen
- *Guadua trinii* Nees = *Bambusa trinii* Nees
- *Ischaemum urvilleanum* Kunth = *Ischaemum minus* Presl
- *Merostachys burchellii* Munro = *Merostachys multiramea* Hackel
- *Paspalum vaginatum* Swartz = *Paspalum distichum* L.
- *Sporobolus poiretii* (B. et S.) Hitchc. = *Sporobolus indicus* (L.) R. Brown

Tais "senões" não diminuem o valor deste último manuscrito de Rambo, sobre o qual ele prostrou a cabeça, quando acometido de um mal fulminante, *dormiens in Domino*. Ao agrostólogo é fácil detectar e solucionar os problemas de sinonímia.

Somos de opinião que "Gramineae Riograndenses" deva ser publicado na forma como Rambo deixou o manuscrito. É valioso, portanto útil aos pesquisadores da flora rio-grandense e brasileira.

Itajaí, 10 de maio de 1983

Dr. Raulino Reitz

Prof. Ronaldo Wasum conferiu o manuscrito com as anotações do fichário do Herbarium Anchieta e anotou ao pé da página as modificações introduzidas desde a morte do autor.

O editor

GRAMINEAE RIOGRANDENSES

B. Rambo SJ.

Resumo: Em trinta anos de herborização no extremo sul do Brasil, o autor reuniu 3124 números de Graminae em ... gêneros e 301 espécies, determinadas na maioria pelos especialistas americanos da família; o resultado é aqui apresentado como primeira aproximação à agrostologia do Rio Grande do Sul.

Zusammenfassung: Verfasser hat in dreissig Jahren Sammelarbeit im südlichsten Brasilien 3124 Nummern aus der Familie der Gräser zusammengetragen, die zum grössten Teil von den amerikanischen Fachkennern der Familie bearbeitet wurden und ... Arten in 301 Gattungen enthalten; die folgende Aufzählung will weiter nichts sein als ein Beitrag zur zukünftigen Gesamtdarstellung der Gramineen in Rio Grande do Sul.

Altough the Grasses represent at least 10% of the Riograndean phanerogamous flora, their knowledge is very far from being satisfactory. Flora Brasiliensis II 2,3 (1871-1883, Döll and Hackel) mentions no more than 15 species as collected within this State. Lindman records passing (1900) 69 components of the Campos. Bornmüller (1934) collected only 13 species on the northeastern Highlands, and Malme (1936) 46 from the southernmost sea coast.

Somewhat better was the situation as to the collected and partially determined material. The Herbarium Dutra (about 1890-1939), among 1629 collections of ferns, legums and compositis, has 253 ones of mostly determined Grasses; and the Herbarium Araujo (about 1920-1961) contains, in round number, 250 species determined by Chase and Parodi. Both of them remain unpublished.

As in almost all other Families, the Grasses of the Republic of Uruguay is by far better known than that of Rio Grande do Sul. Arechavaleta (1894) has about 180 species, Herter (1930) about 280 natives ones.

Santa Catarina, northern neighbor State to Rio Grande do Sul, had, according to Reitz (1956), 185 known species.

The total number of Gramineae in Rio Grande do Sul and its immediate neighborhood, cited in literature or herbaria, published or unpublished, is 570 (taken at face value.).

The Herbarium Anchieta (1931-1961, PACA) contains 3124 collections of Gramineae with 301 species in ... genera. For the following enumerations some geographic symbols may be useful for an easier orientation:

NEL – northeastern Lowlands: Campos, swampy plains, dry granite or sandstone hills, forests, woodlets, gallery.

NEC – northeastern sea coast: sand dunes, woodlets, swamps, dry Campos.

NES – northeastern slope of the Serra Geral: subtropical rain forest.

NEH – northeastern Highlands: Campos, swamps, Araucaria forests.

SEC – south eastern sea coast: Campos, swamps, dry plains, woodlets, gallery.

SEH – southeastern hills lands: dry Campos, woodlets, gallery.

SW – southwestern plains: dry Campos, swamps, gallery, sandstone hills.

CL – central Lowlands: similar to the precedent.

CS – central slope of the Serra Geral: subtropical rain forest.

CH – central highlands: Campos, swamps, woodlets, gallery.

NWH – northwestern Highlands: partly Campos, partly rain forest.

NWU – rain forest belt in the valley of the Uruguay River.

RGS – Rio Grande do Sul.

SC – Santa Catarina.

Ur – Uruguay.

In almost all cases, one or more specimens have been determined by Chase, Parodi, or Swallen, sometimes by two or all of them; divergences are duly annotated. Where there is no indication as to collector and determinator, collection was made by B. Rambo SJ, and determination by B. Rambo SJ and K. Emrich in common work; the latter also obtains when determinator is omitted.

AGROSTIS L.

Agrostis hygronometrica Nees, Agrost. Bras. (1829) 404; Malme, Sv. Bot. Tidskr. 30 (1936) 11. — *Bromidium hygrometricum* Nees et Mey., Nov. Act. Cur. 17, Suppl. 2 ()154; Döll, Fb 3,2 (1876) 103.

29318 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus humidis, 26.10.1945, fl., det. Swallen. (2)

37986 RGS, Barra do Ribeiro p. P. Alegre, NEL, in arenosis humidis, 14.11.1948, fr., det. Swallen. (2)

54418 RGS, Vila Manresa p. P. Alegre, NEL, in agro subhumido, 12.11.1953, fl., fr., det. Swallen. (2)

Habit: 20 cm high, in dense stands.

Habitat: Wet, rocky, more or less bare places.

Distribution in RGS: Central and northeastern Lowlands.

General distribution: RGS.

***Agrostis juergensii* Hack..**

2714 S. Leopoldo, NEL, in humidis graminosis, 25.11.1935, fr., leg. C. Orth SJ, det. Chase, Parodi. (1)

37867 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus alte graminosis, 6.11.1948, fr. (1)

38005 RGS, Barrá do Ribeiro, NEL, in arenosis humidis, 14.11.1948, fr., det. Swallen. (1)

Habit: 60 cm high.

Habitat: Somewhat wet, grassy to shrubby Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: Central and northeastern Lowland Campos of RGS.

***Agrostis koelerioides* E. Desv. E. Desv. in Gay, Flor. Chil. 6 () 317; Malme, Sv. Bot. Tidskr. 30 (1936) 11.**

749 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus, 15.10.1932, fr., leg. C. Orth SJ, det. Chase; Parodi: var. pampeana Par. (2)

57012 RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus parce humosis, 26.10.1955, fl. (2)

57214 RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 15.10.1955, fl. (2)

62575 RGS, Garibaldi, NEH, in incultis, 13.10.1957, fl., leg. O. Camargo 2066. (2)

(1) = *Agrostis hygrométrica* Nees.

(2) = *Agrostis tandilensis* (O.K.) Parodi, det. A. Normann, 1974.

Habit: 20 cm, in dense stands.

Habitat: Somewhat wet, rocky places with thin soil layer.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: Chile; RGS.

Agrostis longiberbis Hack.,

36481 RGS, Cambará p. S. Francisco de Paula, NEH, in humidis graminosis, 2.1948, fl., det. Swallen. (3)

62970 RGS, Lageadinho p. Bom Jesus, NEH, in campestribus, 10.1.1958, fl., leg. O. Camargo 3113. (3)

Habit: Dense bunches, 30 cm high.

Habitat: Somewhat wet, grassy places.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: RGS.

Agrostis montevidensis Spreng. ap. Nees, Agrost. Bras. (1829) 403; Döll, FB 2,3 (1878) 28, tab. 7, fig. 2; Ekman, Ark. f. Bot. 13, n. 10 (1913); Reitz, Sellowia 7 (1956) 96; Arechavaleta, Gram. Ur. (1894) 286.

860 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus subhumidis, 15.10.1933, fr., leg. C. Orth SJ, det. Chase.

6994 SC. Lages, in graminosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

33539 Ur, Juan Jackson p. Soriano, in campestribus subhumidis, 1.1944, fr., leg. et det. Rosengurtt PE-5415.

36468 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus subhumidis, 2.1948, fr., det. Swallen.

37097 Chile, Cautín, in prato, 3.12.1945, fl., leg. H. Gunkel, det. Parodi.

38074 RGS, Vila Manresa p. P. Alegre, NEL, in subhumidis, 17.11.1948, fr., det. Swallen.

48370 SC, Campo dos Padres p. Bom Retiro, in campestribus humidis, 17.12.1948, fr., leg. R. Reitz 3409, det. Swallen.

54319 RGS, Fachinal Preto p. Bom Jesus, NEH, in dense graminosis, 29.1.1950. leg. R. Reitz 4997, det. Swallen.

54996 RGS, Vila Oliva p. Caxias, NEH, in campestribus alte graminosis, 24.2.1954, fr.

55060 RGS, ibidem, 24.2.1954, fr., det. Swallen.

56378 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in campestribus subhumidis, 2.1.1955, fl.

59678 RGS, Farroupilha, NEH, in subhumidis, 16.5.1956, fr., leg. O. Camargo 672.

60086 SC, Campo dos Padres p. Bom Retiro, in subhumidis, 22.1.1957, fr.

60616 RGS, Farroupilha, NEH, in humidis, 16.5.1956, fr., leg. O. Camargo 672.

60821 RGS, ibidem, 5.6.1957, fl., leg. O. Camargo 1580.

64211 SC, S. Joaquim, in campestribus, 10.1.1958, fl., leg. Mattos.

Habit: Thick bunches, very smooth, 30 cm high.

Habitat: More or less wet Campos.

Distribution in RGS: Over the whole of the Campos area.

General distribution: From Minas Gerais to Ur.

Agrostis ramboi Par., Bol. Soc. Arg. Bot. 1,n.2 (1946) 119, fig. 1; Reitz, Sellowia 7 (1956) 96.

9068 RGS, Bom Jesus, NEH, ad rupes in rivulo, 16.1.1942, fr., det. Parodi.

35146 RGS, Fazenda da Ronda p. Vacaria, NEH, ad rupes in rivo, 15.1.1947, fl., det. Parodi.

36444 RGS, Cambará p. S. Francisco de Paula, NEH, ad rupes in rivo, 2.1948, fr., det. Swallen.

44633 RGS, Vila Oliva p. Caxias, NEH, ad rupes in rivo, 3.12.1949, fr., det. Parodi.

45458 RGS, Serra da Rocinha p. Bom Jesus, NEH, ad rivum in rupestribus, 18.1.1950, fr., det. Parodi.

48371 SC, Campo dos Padres p. Bom Retiro, in ripariis alte graminosis, 17.12.1948, fr., leg. R. Reitz 3416, det. Swallen.

54717 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, ad rupes in rivulo, 1.1.1954, fr., det. Swallen.

56403 RGS, ibidem, 2.1.1955, fl.

60074 SC, Campo dos Padres p. Bom Retiro, ad rupes in flumine Uruguay supremo, 22.1.1957, fl.

Habit: Dense stands, 40 cm high.

Habitat: Flooded rocks.

Distribution in RGS: Northeastern Highlands.

General distribution: Highlands of SC and RGS.

AMPHIBROMUS Nees

Amphibromus rubrivalvis (Trin.) Swallen, Herter, Fl. Ur. (1930) 35. *Avena scabririvalvis* Trin., Bull. Sci. Ac. Pétersb. 1 (1836) 67.

57141 RGS, Vila Manresa p. P. Alegre, NEL, in palude, 4.11.1955, fl.

Habit: Erect, slender, 1 m high.

Habitat: Sparsely vegetated swamp.

Distribution in RGS: Northeastern Lowlands.**General distribution:** Chile, Bolívia, Argentina, Ur, RGS.**ANDROPOGON L.*****Andropogon arenarius* Hack.,**

Arechavaleta, Gram. Ur. (1894) 224; Herter, Fl. Ur. (1930) 26; Malme, Sv. Bot. Tidskr. 30 (1936) 9; Reitz, Sellowia 7 (1956) 96.

54344 SC, Laguna, ad litus in arenosis, 29.2.1952, fl., leg. R. Reitz 5040, det. Swallen.**63088** RGS, Pelotas, SEC, in siccis graminosis, 11.3.1956, fl., leg. Schlichting.**Habit:** Dense bunches, 30 cm high.**Habitat:** Dry sandy places.**Distribution in RGS: Sea coast.****General distribution:** SC, RGS, Ur.***Andropogon bicornis* L., Spec. Plant. (1753) 1046; Hackel, FB 2,3 (1883) 283; Ekman, Ark. f. Bot. 11, n. 4 (1912) 6; idem, Ark. f. Bot. 13, n. 10 (1913) 8; Reitz, Sellowia 7 (1956) 96.****1002** RGS, S. Leopoldo, NEL, in campestribus alte dumosis, 10.4.1934, fr., leg. C. Orth SJ, det. Chase.**31689** SC, Sombrio p. Araranguá, in alte dumosis, 6.2.1946, fr.**58920** RGS, Estação de Silvicultura p. Sta. Maria, CL, in campestribus alte graminosis, 20.2.1956, fl., leg. O. Camargo 22.**63658** RGS, P. Alegre, NEL, in dumosis, 8.3.1957, fl., leg. Camargo.**Habit:** Tall bunches, up to 1 m high.**Habitat:** More or less wet, shrubby Campos.**Distribution in RGS: Central and northeastern Lowlands, sea coast.****General distribution:** Through the whole of tropical and subtropical America, down to Uruguay.***Andropogon condensatus* H. B. K., Nov. Gen. 1 ()188; Hackel, FB 2,3 (1883) 296; Arechavaleta, Gram. Ur. (1894) 220; Herter, Flor. Ur. (1930) 26; Reitz, Sellowia 7 (1956) 96.****6953** SC, Lages, in campestribus siccis alte graminosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.**6955** SC, ibidem, leg. A. Bruxel SJ, det. Swallen.**7005** SC, ibidem, 1935, fr., leg. A. Bruxel, det. Swallen.**9841** RGS, Tupanciretan, NWH, in campestribus siccis, 29.1.1942, fr.**26275** RGS, Fazenda do Jarau p. Quaraí, SW, in campestribus siccis dumosis, 1.1945, fr.

- 31207** RGS, sine loco, sine die, fl.
- 33467** RGS, S. Leopoldo, NEL, in campestribus siccis dumosis, 25.7.1946, fr., leg. E. Henz SJ.
- 52759** RGS, Belém Novo p. P. Alegre, NEL, in campestribus alte graminosis, 3.7.1952, fr., leg. et det. A. Beetle 1586.
- 54994** RGS, Vila Oliva p. Caxias, NEH, in subhumidis alte graminosis, 24.2.1954, fr., det. Swallen.
- 55343** SC, Mafra, in campestribus subhumidis, 26.1.1933, fr., leg. R. Reitz 7188, det. Swallen.
- 55673** RGS, Fachinal Preto p. Bom Jesus, NEH, 29.01.1950, in campestribus subhumidis leg. R. Reitz 5172, det. Swallen. (4)
- 56728** RGS, Vila Oliva p. Caxias, NEH, in campestribus humidis, 8.2.1955, fr. (5)

Habit: Dense bunches, up to 1 m high.

Habitat: Dry to wet, open to shrubby Campos.

Distribution in RGS: Fairly common all over de Campos area.

General distribution: From Ecuador to Paraguay, South Brazil and Uruguay.

Andropogon consanguineus Kunth, Enum. 1 () 494; Hackel, FB 2,3 (1883) 237; Herter, Fl. Ur. (1930) 27.

848 RGS, Sapucaia p. S. Leopoldo, NEL, in campestribus arenosis dumosis, 10.6.1933, fl., leg. C. Orth SJ, det. Chase.

38033 RGS, Vila Manresa p. P. Alegre, NEL, in rupestribus parce humosis, 17.11.1948, fr.

44467 RGS, Granja Neugebauer p. Itapoan, NEL, in humidis parce humosis, 19.11.1949, fr., det. Swallen.

45317 RGS, Serra da Rocinha p. Bom Jesus, NEH, in turfosis, 18.1.1950, fr.

52742 RGS, Uruguaiana, SW, in alte graminosis, 1.8.1952, fr., leg. et det. Beetle.

Habit: Dense bunches, 40 cm high.

Habitat: More or less dry, grassy, Campos.

Distribution in RGS: Southwestern plains, northeastern Lowlands; probably over the whole of the Campos area.

General distribution: From Amazonia to Uruguay.

Andropogon glaucescens H. B. K., Nov. Gen. et Spec. 1 () 186; Hackel, FB 2,3 (1883) (var. *genuinus* Hack.); Arechavaleta, Gram. Ur. (1894) 225.

(4) = **Axonopus affinis** Chase, det Swallen, 04.04.1935.

(5) = **Hypogynium virgatum** (Desv.) Dandy, det. A. M. Barcellos Hervé, 27.09.1982.

870 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 15.10.1933, fr., leg. C. Orth SJ, det. Chase.

Habit: Dense bunches, 50 cm high.

Habitat: Dry, open, Campos.

Distribution in RGS: NEL, apparently rare.

General distribution: From Goiás and Minas Gerais to Uruguay.

Andropogon hassleri Hack.,

52756 RGS, Sto Antônio da Patrulha, NEL, in campestribus alte graminosis, 29.6.1952, fr., leg. et det. A. Beetle.

58944 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 1.3.1956, fr., leg. O. Camargo 137.

Habit: Up to 1 m high.

Habitat: Dry, secondary growth.

Distribution in RGS: Central and northeastern Lowlands, probably of wider dispersion.

Andropogon hypogynus Hack.,

2516 RGS, Portão p. S. Leopoldo, NE, in subpaludosis alte dumosis, 17.6.1936, fr., Leg. C. Orth SJ, det. Chase.

Habit: 40 cm high.

Habitat: Swampy, more or less shrubby places.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Piauí to RGS.

Andropogon imberbis Hack.,

Arechavaleta, Gram. Ur. (1894) 216; Herter, Fl. Ur. (1930) 27.

37842 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 4.11.1948, fr., det. Swallen.

57542 RGS, ibidem, 19.11.1955, fl.

61992 RGS, Farroupilha, NEH, in siccis dense graminosis, 18.11.1957, fl., leg. O. Camargo 2572.

64342 SC, S. Joaquim, in campestribus, 1.2.1958, fl., leg. J. Mattos.

Habit: Dense, stiff bunches, up to 60 cm high.

Habitat: Dry, open Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: RGS and Uruguay.

Andropogon lateralis Nees, Agrost. Bras. (1829) 329; Ekman, Ark. f. Bot. 11, n. 4 (1912) 6; idem, Ark. f. Bot. 13, n. 10 (1913) 8; Herter, Fl. Ur. (1930) 27. — An. glaucescens H. B. K. var. lateralis (Nees) Hack., FB 2,3 (1883) 289; Arechavaleta, Gram. Ur. (1894) 227. Reitz, Sellowia 7 (1956) 96.

- 7007 SC, Lages, in campestribus subhumidis, 1935, fr., leg. A. Bruxel SJ, det. Swallen, (var. *incanus* (Hack.) Henr.).
- 10833 RGS, Lagoa Vermelha, NEH, in campestribus subhumidis, 1.1943, fr., leg. E. Friderichs SJ.
- 25576 RGS, Fazenda Sta Cecília, SW, p. S. Gabriel, in campestribus subhumidis, 1.1944, fr.
- 25581 RGS, ibidem, 1.1944, fr.
- 25606 RGS, ibidem, 1.1944, fr., det. Swallen (var. *incanus* (Hack.) Henr.)
- 27759 S. Paulo, Campos do Jordão, in subhumidis, 1.1944, fr., leg. E. Friderichs SJ, det. Swallen (var. *incanus* (Hack.) Henr.).
- 27804 S. Paulo, ibidem, 1.1944, fr., leg. E. Friderichs SJ, det. Swallen (var. *incanus* (Hack.) Henr.)
- 30220 Ur. Tacuarembó, in campestribus arenosis, 3. 1945, fr. leg. et det. B. Rosengurtt 5-4739, (var. *incanus* (Hack.) Henr.)
- 30727 RGS, Vila Oliva p. Caxias, NEH, in campestribus subhumidis, 28. 1.1946, fr., det. Swallen (var. *incanus* (Hack.) Henr.).
- 36458 RGS. Cambará p. S. Francisco de Paula, NEH, in campestribus graminosis, 2.1948, fr., det. Swallen (var. *incanus* (Hack.) Henr.).
- 39337 RGS, Belém Novo p. P. Alegre, NEL, in campestribus subsiccis, 31.12.1948, fr.
- 40797 RGS, Sapucaia p. S. Leopoldo, NEL, in campestribus dumosis, 1.4.1949, fr., det. Swallen.
- 40898 RGS, ad montem das Cabras p. S. Leopoldo. NEL, in campestribus subhumidis, 8.4.1949, fr., det. Swallen.
- 44950 RGS, Canela p. Gramado in campestribus siccis, 26.12.1949, fl.
- 45372 RGS. Serra da Rocinha p. Bom Jesus, NEH, in campestribus turfosis, 18.1.1950, fr.
- 45579 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in campestribus subhumidis, 30.1.1950, fr.
- 50423 RGS. Serra da Rocinha p. Bom Jesus, NEH, in campestribus subhumidis, 19.1.1950, fr., leg. A. Sehnem SJ 4268.
- 50456 RGS, ibidem, 19.1.1950, fr., leg. A. Sehmeh SJ 4251.
- 50849 Paraná, S. Luiz de Purunha p. C. Largo, in campestribus subturfosis, 4.12.1949, fr., leg. G. Hatschbach 1635, det. Parodi.
- 50900 Paraná, Piraquara, in subpaludosis dumosis, 17.12.1950, fl., leg. G. Hatschbach 2051, det. Parodi.
- 52758 RGS, Belém Novo p. P. Alegre, NEL, in campestribus alte graminosis, 13.7.1952, fl.
- 54313 RGS, Fachinal Preto p. Bom Jesus, NEH, in campestribus dense graminosis, 29.1.1950, leg. R. Reitz 4986, det. Swallen.
- 54321 RGS. ibidem, 29.1.1950, leg. R. Reitz 4999, det. Swallen.
- 54343 SC, Laguna, ad litus in arenosis, 29.2.1952, leg. R. Reitz 5039, det. Swallen.

- 54362** SC, Cambajuva p. S. Joaquim, in campestribus dense graminosis, 29.1.1950, leg. R. Reitz 5075, det. Swallen.
- 54364** SC, Bom Jardim p. S. Joaquim, in campestribus dense graminosis, 29.1.1950, fl., leg. R. Reitz 5077, det. Swallen.
- 54716** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in campestribus dumosis, 1.1.1954, fr., det. Swallen.
- 54995** RGS, Vila Oliva p. Caxias, NEH, in campestribus alte graminosis, 24.2.1954, fr., det. Swallen.
- 55345** SC, Massiambú p. Palhoça, in arenosis dense graminosis, 19.12.1952, leg. R. Reitz 7186, det. Swallen.
- 55430** SC. Massiambú p. Palhoça, in graminosis arenosis, 19.12.1952, fl., leg. R. Reitz 7056, det. Swallen.
- 55434** SC, Serra da Boa Vista p. S. José, in dense graminosis, 2.2.1953, fl., leg. R. Reitz 7048, det. Swallen.
- 55439** SC, Mafra, in dense graminosis, 26.1.1953, fl., leg. R. Reitz 7043, det. Swallen.
- 55672** RGS, Fachinal Preta p. Bom Jesus, NEH, in campestribus dense graminosis, 29.1.1950, fl., leg. R. Reitz, 5179, det. Swallen.
- 56302** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in dense graminosis, 2.1.1955, fl.
- 58048** SC, Massiambú p. Palhoça, in arenosis, 24.9.1953, fr., leg. R. Reitz 8321, det. Swallen.
- 59168** RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus parce humosis, 3.2.1956, fr.
- 60445** RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus subdumosis, 8.4.1956, fr., leg. O. Camargo 563.
- 61671** RGS, Taquarí, NEL, in campestribus, 10.12.1957, fl., leg. O. Camargo 2849.
- 62870** RGS, P. Alegre, NEL, in campestribus, 8.2.1958, fl., leg. O. Camargo 3140.
- 62957** RGS, Lageadinho p. Bom Jesus, NEH, in campestribus, 20.1.1958, fl., leg. O. Camargo 3130.
- 63090** RGS, Pelotas, in campestribus, 11.3.1956, fl., leg. Schlichting.
- 63091** RGS, ibidem, 10.3.1951, fl., leg. Schlichting.
- 64269** SC, S. Joaquim, in campestribus, 20.1.1958, fl., leg. J. Mattos.

Habit: Dense bunches, stalks up to 1 m high.

Habitat: Dry to moderately wet Campos.

Distribution in RGS: One of the most common Campos grasses throughout the whole of the Campos area, especially on the northeastern Highlands where it is dominant.

General distribution: From Minas Gerais to Uruguay.

Andropogon leucostachyus H.B.K., Nov. Gen. et Spec. 1 () 187; Arechavaleta, Gram. Ur. (1894) 221; Herter, Flor. Ur. (1930) 27; Reitz, Sellowia 7 (1956) 96. — *A. virginicus* L., ssp. *leucostachyus* (H.B.K.) Hack., FB 2,3 (1883) 286.

38204 RGS, Sapucaia p. S. Leopoldo, NEL, in campestribus siccis, 22.11.1948, fr., det. Swallen.

39330 RGS, Belém Novo p. P. Alegre, NEL, in campestribus siccis dumosis, 31.12.1948, fr., det. Swallen.

39931 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in campestribus siccis, 12.1.1949, fr.

39937 RGS, ibidem, in secundariis, 12.1.1949, fr.

40757 RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis, 1.4.1949, fr., det. Swallen.

40808 RGS, ibidem, 1.4.1949, fr., det. Swallen.

47583 SC, Morro do Baú p. Itajaí, in summo monte in dumosis, 29.1.1948, fr., leg. R. Reitz 3070, det. Asplund.

48356 SC, Figueiredo p. Bom Retiro, in alte dumosis, 28.12.1948, fr., leg. R. Reitz 3459, det. Swallen.

55373 SC, Massiambú p. Palhoça, in arenosis, 19.12.1952, fl., leg. R. Reitz 7153, det. Swallen.

55401 SC, Praia Braba p. Itajaí, in arenosis, 5.3.1953, fl., leg. R. Reitz 7109, det. Swallen.

55410 SC, Massiambú p. Palhoça, in arenosis, 5.2.1953, fl., leg. R. Reitz 7088, det. Swallen.

55423 SC, ibidem, 12.3.1953, leg. R. Reitz 7068, det. Swallen.

57027 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in graminosis siccis, 26.10.1955, fl.

59383 RGS, Farroupilha, NEH, in alte graminosis, 15.11.1956, fl., leg. O. Camargo 956.

59384 RGS, ibidem, 25.11.1956, fl., leg. O. Camargo 1027.

61994 RGS, Taquarí, NEL, in alte graminosis, 8.12.1957, fl., leg. O. Camargo 2688.

61995 RGS, ibidem, 10.12.1957, fl., leg. O. Camargo 2854.

61996 RGS, P. Alegre, NEL, in alte graminosis, 19.11.1957, fl., leg. O. Camargo 2588.

63660 RGS, ibidem, 8.3.1957., leg. O. Camargo.

Habit: Thin bunches, stalks slender, up to 1 m high.

Habitat: Common on more or less dry Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: From Guyana and Venezuela to Uruguay.

Andropogon lindmanii Hack.,

Malme, Sv. Bot. Tidskr. 30 (1936) 9.

1960 RGS, Vacaria, NEH, in campestribus subhumidis, 6.1.1935, fr., leg. et det. Dutra.

2514 RGS, in summo monte Steinkopf p. S. Leopoldo, NEL, in campestribus subhumidis, 10.6.1936, fr., leg. C. Orth SJ, det. Parodi.

7439 RGS, S. Leopoldo, NEL, in alte graminosis, 1907, fr., leg. F. Theissen SJ.

Habit: Dense bunches, stalks up to 75 cm high.

Habitat: Dry to more or less wet places.

Distribution in RGS: Southeastern coast, northeastern Lowlands and Highlands.

General distribution: Apparently RGS only.

Andropogon macrothrix Trin., Mém. Ac. St. Pétersb. sér. 6,2 () 270; A. ternatus (Spreng.) Nees, ssp. macrothrix (Trin.) Hack., FB. 2,3 (1883) 287; Ekman, Ark. f. Bot. 13,n.10 (1913) 8.

36448 RGS, Cambará p. S. Francisco de Paula, NEH, in alte graminosis, 2.1948, fr., det. Swallen.

48304 SC, Campo dos Padres p. Bom Retiro, in campestribus, 22.2.1948, fr., leg. R. Reitz 3376, det. Swallen.

48355 SC, Figueiredo p. Bom Retiro, in campestribus humidis dumosis, 28.12.1948, fr., Leg. R. Reitz 3471, det. Swallen.

50852 Paraná, Piraquara, in campestribus, 13.11.1949, fr., leg. G. Hatschbach 1615, det. Parodi.

54335 RGS, Fachinal Preto p. Bom Jesus, NEH, in campestribus subhumidis, 28.1.1950, fr., leg. Reitz 5023, det. Swallen.

54703 RGS, Fazenda Englert p. S. Francisco, NEH, in campestribus dense graminosis, 1.1.1954, fr., det. Swallen.

59382 RGS, Farroupilha, NEH, in alte graminosis, 15.11.1956, fl., leg. O. Camargo 933.

60079 SC, Campo dos Padres p. Bom Retiro, in campestribus subsiccis, 20.1.1957, fr.

Habit: Thin bunches, stalks about 50 cm high.

Habitat: Dry to moderately wet Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: From Goiás and Minas Gerais to RGS.

Andropogon paniculatus Kunth, Enum. 1 () 494. — A. condensatus H.B.K. var. paniculatus (Kunth) Hack., FB 2,3 (1883) 297; Arechavaleta, Gram. Ur. (1894) 221.

747 RGS, S. Leopoldo, NEL, in siccis dumosis, 10.1.1933, fr., leg. C. Orth SJ. det. Chase.

7049 SC, Lages, in siccis dumosis, 1935, fr., leg. A. Bruxel SJ. det. Swallen.

- 25565** RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus siccis alte graminosis, 1.1944, fr., det. Swallen.
- 28850** RGS, Caracol p. Canela, NEH, in campestribus dumosis, 5.2.1945, fr., leg. K. Emrich. (6).
- 30741** RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis dumosis, 28.1.1946, fr.
- 32973** RGS, Parecí p. Montenegro, NEL, in campestribus siccis dumosis, 15.1.1946, fr., leg. C. Ritter SJ, det. Swallen. (6)
- 35167** RGS, Fazenda da Ronda p. Vacaria, NEH, in siccis dumosis, 15.1.1947, fr.
- 39928** RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in secundariis arenosis, 12.1.1949, fr.
- 41334** RGS, Morretes p. P. Alegre, NEL, in arenosis dumosis, 2.5.1949, fr.
- 46223** RGS, Cristo Rei p. S. Leopoldo, NEL, in campestribus siccis dumosis, 10.3.1950, fr., det. Parodi.
- 46806** RGS, Fazenda do Arroio p. Osório, NEC, in campestribus arenosis siccis, 14.4.1950, fr.
- 47715** SC, Brusque, in dumosis secundariis, 6.3.1948, fr., leg. R. Reitz 2876, det. Parodi. (6)
- 48923** RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus alte dumosis, 28.2.1956, fr., leg. O. Camargo 11.
- 59821** RGS, Farroupilha, NEH, in dumosis, 25.3.1947, fl., leg. O. Camargo 1319.
- 58922** RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 28.2.1956, fl., leg. O. Camargo 16.
- 60246** RGS, Farroupilha, NEH, in alte graminosis, 14.1.1957, fr., leg. O. Camargo 1111.
- 60439** RGS, Estação de Silvicultura p. Sta Maria, CL, in alte dumosis, 8.4.1956, fr., leg. O. Camargo 592.
- 60732** RGS, Passo Fundo, CH, in campestribus, 4.1957, fr., leg. P. Freddiani.
- 61993** RGS, Taquari, NEL, in alte graminosis, 8.12.1957, ster., leg. O. Camargo 2861.
- 63089** RGS, Pelotas, SEC, in alte graminosis, 10.3.1956, fl., leg. Schlichting.

Habit: Large brownisch bunches, up to 1 m high.

Habitat: Shrubby Campos and secondary growth.

Distribution in RGS: Apparently over the whole of the Campos and cultivated area.

General distribution: Southern Brazil to Uruguay.

(6) = *Schizachyrium microstachium* (Desv.) R. A. I. ssp. *microstachyum*, det. A. M. Barcellos Hervé, 27.09.1982.

Andropogon saccharoides Swartz, Prodr. Veg. Ind. Occid. () 26; Hackel, FB 2,3 (1883) 292; Arechavaleta, Gram. Ur. (1894) 228; Ekman, Ark. f. Bot. 11,n.4 (1912) 8; Herter, Fl. Ur. (1930) 27.

2721 RGS, Kappesberg p. Montenegro, NES, in dumosis secundariis, 4.1.1936, fr., leg. C. Orth SJ, det. Chase (Parodi: A. hassleri Hack.).

26311 RGS, Fazenda do Jarau p. Quaraí, SW, in siccis alte graminosis, 1.1945, fr.

40704 RGS, Cristal p. P. Alegre, NEL, in arenosis alte graminosis, 31.3.1949, fr., det. Swallen.

41216 RGS, ad fl. Caí inferior p. P. Alegre, NEL, in alte dumosis, 27.4.1949, fr., det. Swallen.

41572 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in secundariis arenosis, 16.5.1949, fr., det. Swallen.

42436 RGS, Pareci p. Montenegro, NEL, in secundariis alte graminosis, 7.7.1949, fr., det. Swallen.

56499 RGS, Torres, NEC, ad litus in arenosis, 19.1.1955, fr., det. Swallen (Parodi: A. laguroides DC.).

59677 RGS, Farroupilha, NEH, in alte graminosis, 16.5.1956, fr., leg. O. Camargo 669.

59822 RGS, ibidem, 25.3.1957, fl., leg. O. Camargo 1207.

60608 RGS, 8.5.1957, fr., leg. O. Camargo 1400.

60617 RGS, ibidem, 16.5.1957, fl., leg. O. Camargo 669.

Habit: Medium-sized bunches, stalks up to 75 cm high.

Habitat: Dry to moderately wet Campos and secondary growth.

Distribution in RGS: Apparently over the whole of the Campos and cultivated area.

General distribution: Through the whole of tropical and subtropical America.

Andropogon sellianus Hack., Bull. Herb. Boiss. 2. sér. 4,n.3 (1904) 266; Ekman, Ark. f. Bot. 11, n.4 (1912) 6; idem, Ark. f. Bot. 13, n.10 (1913) 8; Reitz, Sellowia 7 (1956) 97.

846 RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 5.1.1933, fr., leg. C. Orth SJ, det. Swallen (Khase: A. macrothrix Trin.).

6862 SC, Florianópolis, in arenosis alte graminosis, 1935, fr., leg. A. Bruxel SJ, det. R. Swallen.

6956 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

7068 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

9580 RGS, Tupanciretan, NWH, in campestribus siccis, 28.1.1942, fr., det. Swallen.

28003 RGS, Tramandaí, NEC, in arenosis submobilibus, 1.1945, fr., leg. P. Gonçalves.

- 35449** RGS, S. Leopoldo, NEL, in arenosis alte graminosis, 6.11.1946, fr., leg. E. Henz SJ.
- 37393** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 5.7.1948, fr.
- 38115** RGS, Morro da Polícia p. P. Alegre, NEL, in siccis alte graminosis, 20.11.1948, fr., det. Swallen.
- 40947** RGS, ad montem das Cabras p. S. Leopoldo, NEL, in arenosis dumosis, 8.4.1949, fr., det. Swallen.
- 42790** RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis siccis, 5.7.1948, fr., det. Swallen.
- 46781** RGS, Fazenda do Arroio p. Osório, NEC, in arenosis dumosis, 14.4.1950, fr.
- 46860** RGS, ad fl. Capivarí p. Palmares, NEL, in arenosis siccis, 17.4.1950, fr.
- 51187** Paraná, Matinhos p. Paranaguá, in arenosis, 1.1.1950, fr., leg. G. Hatschbach 1687, det. Parodi.
- 52757** RGS, Belém Novo p. P. Alegre, NEL, in campestribus arenosis, 13.7.1952, fr., leg. et det. A. Beetle.
- 54859** RGS, Torres, prope litus in arenosis, 11.2.1954, fr., det. Swallen.
- 58051** SC, Massiambú p. Palhoça, in arenosis, 4.11.1953, leg. R. Reitz 8327, det. Swallen.
- 58058** SC, ibidem, 5.11.1953, fr., leg. R. Reitz 6350, det. Swallen.
- 59019** RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus alte dumosis, 12.2.1956, fr., leg. O. Camargo 172.
- 58921** RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus arenosis, 28.2.1956, fr., O. Camargo 15.

Habit: Dense bunches, 30 cm high.

Habitat: Dry, sandy places.

Distribution in RGS: Apparently over the whole of the Campos area, notably near the sandy sea coast.

General distribution: Paraguay, SC, RGS, Ur.

- Andropogon tener** (Nees) Kunth, Revisio Gram. 2 () 565; Hackel, FB 2,3 (1883) 301; Ekman, Ark. f. Bot. 11 (1912) n. 4, 5; idem, Ark. f. Bot. 13, n.10 (1913) 7; Herter, Flor. Urug. (1930) 27; Reitz, Sellowia 7 (1956) 97. — *Schizachyrium tenerum* Nees, Agrost. Bras. (1829) 336.
- 7012** SC, Lages, in campestribus, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

7033 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

55057 RGS, Vila Oliva p. Caxias, NEH, in campestribus dense graminosis, 24.2.1954, fr., det. Swallen.

60390 RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus

alte graminosis, 8.4.1956, fl., leg. O. Camargo 593.

Habit: Dense bunches, stalks up to 50 cm high.

Habitat: Dry Campos.

Distribution in RGS: Widely scattered, apparently rare.

General distribution: From southern USA to Uruguay.

Andropogon ternatus (Spreng.) Nees, Agrost. Bras. (1829) 326; Hackel, FB 2,3 (1883) 287 (var. *genuinus* Hack.); Ekman, Ark. f. Bot. 13, n.10 (1913) 8; Herter, Flor. Urug. (1930) 27; Reitz, Sellowia 7 (1956) 97.

8600 RGS, Serra da Rocinha p. Bom Jesus, NEH, in campestribus subhumidis, 14.1.1942, fr., det. Swallen.

9711 RGS, Tupanciretan, NWH, in campestribus, 29.1.1942, fr., det. Beetle.

29343 RGS, Vila Manresa p. P. Alegre, in campestribus siccis, 26.10.1945, fr.

33197 RGS, Vila Oliva p. Caxias, NEH, in campestribus, 16.1.1946, fr., det. Beetle.

55411 SC, Serra da Boa Vista p. S. José, in dense graminosis, 2.2.1953, leg. R. Reitz 7087, det. Swallen.

Habit: Middle-sized bunches, 30 cm high.

Habitat: Moderately wet to dry places.

Distribution in RGS: Widely scattered, apparently over the whole of the Campos area.

General distribution: From Minas Gerais and Paraguay to Uruguay.

Andropogon virgatus Desv., Ham. Prodr. Ind. Occ. () 9; Ekman, Ark.f. Bot. 11, n.4 (1912) 6; idem, Ark. f. Bot. 13, n.10 (1913) 7; Herter, Flor. Urug. (1930) 27; Reitz, Sellowia 7 (1956) 97. — *A. spathiflorus* (Nees) Kunth, Hackel, FB 2,3 (1883) 295, tab. 68, fig. 1.

745 RGS, ad montem das Cabras p. S. Leopoldo, NEL, in campestribus siccis arenosis, 15.11.1933, fl., leg. C. Orth SJ, det. Chase, Parodi.

6992 SC, Lages, in campestribus siccis arenosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

7016 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

26035 RGS, Fazenda do Jarau p. Quaraí, SW, in campestribus siccis, 29.12.1944, fr.

30741 RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis, 28.1.1946, fr., det. Beetle.

31976 SC, Sombrio p. Araranguá, in campestribus siccis, 9.5.1945, fr., leg. R. Reitz, det. Beetle.

- 32771 RGS, Parecí p. Montenegro, NEL, in arenosis siccis, 15.11.1945, fr., leg. E. Henz SJ, det. Beetle.
- 35174 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis dumosis, 15.1.1947, fr., det. Beetle.
- 38944 RGS, Chácara Weber p. Itapoan, NEL, in campestribus siccis alte dumosis, 18.12.1948, fr., det. Beetle.
- 39527 RGS, Cachoeirinha p. Gravataí, NEL, in campestribus dumosis, 7.1.1949, fl.
- 52755 RGS, Sto Antônio da Patrulha, in campestribus alte dumosis, 29.6.1952, fr., det. Beetle.
- 40993 RGS, ad fl. Gravataí inferius, NEL, in campestribus siccis, 10.4.1949, fr.
- 53403 RGS, Caaró p. S. Luiz, NWH, in campestribus graminosis, 24.12.1952, fr.
- 53412 RGS, ibidem, 24.11.1952, fr.
- 55000 RGS, Vila Oliva p. Caxias, NEH, in campestribus alte dumosis, 24.2.1954, Fr., det. Swallen.
- 56926 RGS, Caracol p. Canela, NEH, in campestribus alte graminosis, 15.1.1955, fl., leg. K. Emrich.
- 58924 RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus alte graminosis, 28.2.1956, fl., leg. O. Camargo 19.
- 59731 RGS, Farroupilha, in alte graminosis, 28.5.1956, fr., leg. O. Camargo 732.
- 60440 RGS, Estação de Silvicultura p. Sta Maria, in campestribus alte graminosis, 8.4.1956, fr., leg. O. Camargo 591.
- 61991 RGS, Taquarí, NEL, in alte graminosis, 12.12.1957, fl., leg. O. Camargo 2892.

Habit: Strong bunches, up to 1,5 m high.

Habitat: Dry, shrubby Campos and secondary growth.

Distribution in RGS: Over the whole of the Campos and cultivated area.

General distribution: From the Antilles to Uruguay.

ANTHOXANTHUM L.

Anthoxanthum odoratum L., Spec. Plant. (1753) 28; Ekman, Ark.f.Bot. 13, n.10 (1913) 37; Herter, Fl. Ur. (1930) 32.

43456 RGS, Kappesberg p. Montenegro, NEL, in agro, 11.9.1949, fr., det. Swallen.

59521 RGS, Farrroupilha, NEH, in graminosis, 30.8.1956, fl., leg. O. Camargo 729.

62159 RGS, Bento Gonçalves, NEH, in incultis, 6.10.1957, fl., leg. O. Camargo 1957.

62239 RGS, Garibaldi, NEH, in incultis, 13.10.1957, fl., leg. O Camargo 2044.

62375 RGS, Farroupilha, NEH, in incultis, 1.9.1957, fl., leg. O. Camargo 1674.

62403 RGS, ibidem, 12.9.1957, fl., leg. O. Camargo 1701.

Habit: 75 cm. high.

Habitat: Tilled or otherwise disturbed soil.

Distribution in RGS: Northeastern Highlands.

General distribution: Native to the Mediterranean region; in Brasil, from Paraná to RGS and Ur.

ARISTIDA L.

Aristida altissima Arech., An. Mus. Nac. Mont. 4 (1903) 80, tab. 5; Herter, Fl. Ur. (1930) 33.

744 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 5.10.1932, fr., det. Beetle (Chaze: *A. teretifolia* Arech.).

1094 RGS, Novo Hamburgo, NEL, in campestribus siccis alte graminosis, 13.9.1934, fl., leg. C. Orth SJ, det. Chase.

9641 RGS, Tupanciretan, NWH, in campestribus siccis alte graminosis, 28.1.1942, fr., det. Swallen.

38534 RGS, S. Leopoldo, NEL, in campestribus siccis dense graminosis, 4.12.1948, fr., det. Swallen.

44398 RGS, Fião p. S. Leopoldo, NEL, in campestribus arenosis siccis, 14.11.1949, fl., det. Beetle.

56006 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 19.11.1954, fl., det. Swallen.

57023 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fl.

57492 RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus siccis, 18.11.1955, fl., leg. O. Camargo 9.

61967 RGS, P. Alegre, NEL, in siccis alte graminosis, 19.11.1957, fl., leg. O. Camargo 2593.

61969 RGS, Taquarí, NEL, in alte graminosis, 10.12.1957, fl., leg. O. Camargo 2827.

61969 RGS, Farroupilha, NEH, in secundariis alte dumosis, 25.11.1957, fl., leg. O. Camargo 2638.

62749 RGS, P. Alegre, NEL, in alte graminosis, 1.11.1957, fl., leg. O. Camargo 2395.

Habit: Strong bunches, stalks up to 1,5 m high.

Habitat: Dry, poor, strongly mineralized soil.

Distribution in RGS: Frequent in the whole southern Campos area;

also on the northwestern Highlands.

General distribution: RGS and Uruguay.

Aristida circinnalis Lindman, Kongl. Sv. Vet. Ak. Handl. 34, n.6 (1900) 13; Malme, Sv. Bot. Tidskr. 30 (1936) 10.

29316 RGS, Vila Manresa p. P. Alegre, in campestribus siccis alte graminosis, 26.10.1945, fr.

30191 Ur, Atlantida p. Canelones, in campestribus siccis, 3.1.1945, fr., leg. et det. B. Rosengurtt B-4374.

38666 RGS, Rio dos Sinos p. S. Leopoldo, NEL, in campestribus siccis alte graminosis, 10.12.1948, fr., det. Swallen.

41457 RGS, Estação Azevedo p. Montenegro, NEL, in campestribus siccis alte graminosis, 6.5.1949, fr.

43591 RGS, Portão p. S. Leopoldo, NEL, in siccis alte graminosis, 23.9.1949, fr., det. Beetle.

44396 RGS p. S. Leopoldo, NEL, in arenosis siccis, 14.11.1949, fl., det. Beetle.

44468 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis, 19.11.1949, fr.

52751 RGS, Guaíba p. P. Alegre, NEL, in campestribus alte graminosis, 7.7.1952, fr., leg. et det. Beetle.

Habit: Dense, slender bunches, up to 50 cm high.

Habitat: Very dry and poor Campos..

Distribution in RGS: Apparently southern half of the State only.

General distribution: Paraguay, RGS, Uruguay.

Aristida implexa Trin., Act. Petrop. 1836, 48; Döll, FB 2,3 (1878) 24.

1926 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in arenosis siccis, 1.5.1936, fr., leg. C. Orth SJ, det. Chase.

7019 SC, Lages, in campestribus siccis, 1935, fr., leg. A. Bruxel SJ, det. Swallen. (7)

30728 RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis alte graminosis, 28.1.1946, fr., det. Swallen.

35178 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 15.1.1947, fr. (7)

36471 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus subsiccis, 2.1948, fr., det. Swallen.

36479 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus siccis, 2.1948, fr., det. Swallen.

(7) = **Aristida megapotamica** Spr., det. Branca Severo, 01.1979.

38927 RGS, Granja Weber p. P. Alegre, NEL, in campestribus siccis alte graminosis, 18.12.1948, fr.

39872 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in arenosis siccis alte dumosis, 12.1.1949, fr.

40800 Sapucaia p. S. Leopoldo, NEL, in arenosis alte dumosis, 1.4.1949, fr., det. Swallen.

55397 SC, Mafra, in campestribus siccis, 27.1.1953, fr., leg. R. Reitz 7113, det. Swallen.

58977 RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus siccis alte graminosis, 25.1.1956, fr., leg. O. Camargo 88.

62860 RGS, Lageadinho p. Bom Jesus, NEH, in siccis, 20.1.1958, fl., leg. O. Camargo 3158.

Habit: Stiff bunches, stalks up to 50 cm high.

Habitat: Poor, more or less mineralized Campos.

Distribution in RGS: Over the whole Campos area.

General distribution: From Bahia to Uruguay.

Aristida laevis Kunth, Enum. 1 () 192. — *Chaetaria laevis* Nees, Agrost. Bras. (1829) 384. — *A. Complanata* Trin., Act. Petrop. 1829, 85; Döll, FB 2,3 (1878) 15; Arechavaleta, Gram. Ur. (1894) 246; Herter, Fl. Ur. (1930) 33; Reitz, Sellowia 7 (1956) 97.

1001 RGS, S. Leopoldo, NEL, in arenosis siccis, 5.5.1934, fr., leg. C. Orth SJ, det. Chase.

29609 RGS, Parecí p. Montenegro, NEL, in campestribus siccis, 3.10.1945, fl., det. Swallen.

33363 RGS, S. Leopoldo, NEL, in campestribus siccis, 15.9.1946, fl., leg. H. Simas SJ.

34127 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 16.10.1946, fl.

39924 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in arenosis siccis, 12.1.1949, fr.

55384 SC, Massiambú p. Palhoça, in arenosis, 19.12.1952, fr., leg. R. Reitz 7134, det. Swallen.

57022 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fl.

57125 RGS, Esteio p. S. Leopoldo, NEL, in siccis alte graminosis, 14.11.1955, fl.

Habit: Dense bunches, up to 75 cm high.

Habitat: Dry, poor soils.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Paraguay to Uruguay.

General distribution: From Paraguay to Uruguay.

Aristida pallens Cav., Ic. et Descr. 5 () 43, tab. 468; Döll FB 2,3 (1878) 13; Ekman, Ark. f. Bot. 11, n.4 (1912) 30; idem, Ark. f. Bot. 13, n.10 (1913) 37; Arechavaleta, Gram. Ur. (1894) 244; Herter, Flor. Ur. (1930) 33.

- 1071** RGS, S. Leopoldo, NEL, in siccis arenosis, 15.12.1933, fr., leg. C. Orth SJ, det. Chase.
- 29317** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 26.10.1945, fl.
- 32759** RGS, Parecí p. Montenegro, NEL, in arenosis siccis, 17.10.1945, fl, leg. E. Henz SJ.
- 38132** RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis, 20.11.1948, fr., det. Swallen.
- 38539** RGS, S. Leopoldo, NEL, in arenosis siccis, 4.12.1948, fr., det. Beetle.
- 38928** RGS, Granja Weber p. Itapoan, NEL, in campestribus siccis, 18.12.1948, fr., det. Beetle.
- 43727** RGS, Morro Sta Teresa p. P. Alegre, NEL, in campestribus siccis, 3.10.1949, fr., det. Swallen
- 44484** RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis, 19.11.1949, fl., det. Swallen.
- 53213** RGS, Cerro Largo p. S. Luiz, NWH, in agro derelicto, 20.11.1952, fr.
- 55996** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 19.11.1954, fl., det. Swallen.
- 55998** RGS, ibidem, 19.11.1954, fr., det. Swallen.
- 57021** RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fl.
- 57442** RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in ruprestribus subsiccis, 9.10.1955, fl.
- 57494** RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus, 15.11.1955, fl., leg. O. Camargo 32.
- 58517** RGS, Vila Manresa p. P. Alegre, NEL, in dense graminosis, 19.11.1955, fl.
- 61861** RGS, Passo Fundo, CH, in campestribus siccis, 20.10.1957, fl., leg. O. Camargo 2159.
- 61880** RGS, Garibaldi, NEH, ad viam, 20.10.1957, fl., leg. O. Camargo 2280.
- 61970** RGS, Taquarí, NEL, in campestribus siccis, 8.12.1957, fl., leg. O. Camargo 2703.
- 62601** RGS, P. Alegre, NEL, in campestribus, 1.11.1957, fl., leg. O. Camargo 2410.
- 62924** RGS, Taquarí, NEL, in campestribus siccis, 10.12.1957, fl., leg. O. Camargo 3011.

Habit: Dense bunches, up to 50 cm high.

Habitat: Poor, mineralized, Campos soils.

Distribution in RGS: Over the whole of the Campos area, especially frequent in the West.

General distribution: From Minas Gerais to Uruguay and Paraguay.

Aristida teretifolia Arech., An. Mus. Nac. Mont. 4 (1903) 79, tab. 4; Herter, Flor. Urug. (1930) 33.

29315 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis graminosis, 26.10.1945, fr., det. Swallen.

29342 RGS, ibidem, 26.10.1945, fr., det. Swallen.

38127 RGS, Morro da Polícia p. P. Alegre, NEL, in siccis graminosis, 20.11.1948, fr., det. Swallen.

43662 RGS, Morro das Abertas p. P. Alegre, NEL, in campestribus siccis graminosis, 30.9.1949, fl., det. Swallen.

56006 RGS, Vila Manresa p. P. Alegre, NEL, in rupestribus alte graminosis, 19.11.1954, fl., det. Swallen.

58518 RGS, ibidem, 19.11.1955, fl.

Habit: Small, rather stiff, bunches, 30 cm high.

Habitat: Dry, stony, Campos.

Distribution in RGS: P. Alegre only.

General distribution: RGS and Uruguay.

Aristida uruguayensis Henr.,

3900 RGS, Morro Vigia p. Livramento, SW, in siccis graminosis, 12.1.1941, fr., det. Parodi.

Habit: Small bunches, leaves less than 10, stalks 25 cm high.

Habitat: Rocky, dry places.

Distribution in RGS: Apparently southwestern frontier only.

General distribution: RGS, Uruguay.

Aristida venustula Arech., An. Mus. Nac. Mont. 4 (1903) 77; Herter, Flor. Urug. (1930) 33.

1713 RGS, Novo Hamburgo p. S. Leopoldo, NEL, in campestribus siccis, 10.9.1934, fr., leg. C. Orth SJ, det. Beetle.

2711 RGS, Vila Mafresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 19.11.1935, fr., leg. C. Orth SJ, det. Parodi (Chase: A. spegazzinii Arech.).

37840 RGS, ibidem, 4.11.1948, fl., det. Swallen.

44178 RGS, Morro de Santana p. P. Alegre, NEL, in siccis alte graminosis, 2.11.1949, fl., det. Beetle.

61971 RGS, P. Alegre, NEL, in campestribus, 19.11.1957, fl., leg. O. Camargo 2599.

61972 RGS, *ibidem*, 19.11.1957, fr., leg. O. Camargo 2597.

Habit: Large, smooth bunches, inflorescences brownish, 30 cm high.

Habitat: Very dry, rocky Campos.

Distribution in RGS: Over the whole of the southern half of the State.

General distribution: RGS and Uruguay.

ARTHROSTYLIUM Rupr.

Arthrostylidium trinii (Rupr.) Munro, Monogr. () 45; Döll, FB 2,3 (1880) 173; Reitz, *Sellowia* 7 (1956) 97.

4613 RGS, Serra da Rocinha p. Bom Jesus, NEH, in sphagneto, 12.2.1941, ster.

7014 SC, Lages, in silvula subpaludosa ster., leg. A. Bruxel SJ.

8688 RGS, Serra da Rocinha p. Bom Jesus, NEH, in silvula nebulari, 14.1.1942, ster.

9011 RGS, Bom Jesus, NEH, ad rivum in silvula, 16.1.1942, ster.

30717 RGS, Vila Oliva p. Caxias, NEH, ad rivum in dumosis, 28.1.1946, ster.

35180 RGS, Fazenda da Ronda p. Vacaria, NEH, in silvula subpaludosa 15.1.1947, ster.

36480 RGS, Cambará p. S. Francisco de Paula, NEH, in silvula ad rivum, 2.1948, ster.

53788 RGS, Serra da Rocinha p. Bom Jesus, NEH, in silvula nebulari, 3.2.1953, ster.

54074 RGS, *ibidem*, 20.2.1953, ster.

56761 RGS, Vila Oliva p. Caxias, NEH, ad rivum in araucarieto, 8.2.1945, ster.

59828 RGS, Farroupilha, NEH, ad rivulum in dumosis, 2.4.1957, ster., leg. O. Camargo 1224.

60085 SC, Campo dos Padres p. Bom Retiro, in silvula paludosa, 22.1.1957, ster.

Habit: Loose bunches, up to 5 m high.

Distribution in RGS: Northeastern Highlands only.

Habitat: Wet to swampy places.

General distribution: From Minas Gerais to RGS.

ARUNDINARIA Michx.

Arundinaria ulei Hack.,

32475 RGS, Serra da Rocinha p. Bom Jesus, NEH, in sphagneto, 28.2.1946, ster.

35197 RGS, ibidem, 14.2.1947, fr.

36470 RGS, Cambará p. S. Francisco de Paula, NEH, in sphagneto, ster., 2.1948.

45330 RGS, Serra da Rocinha p. Bom Jesus, NEH, in sphagneto, 18.1.1950, fr.

50067 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in spagneto, 21.2.1951, fl.

60084 SC, Campo dos Padres p. Bom Retiro, in sphagneto, 22.1.1957, fl.

Habit: Dense clumps and stands, up to 1,5 m high.

Habitat: Sphagnum swamps.

Distribution in RGS: Northeasternmost Highlands.

General distribution: SC and RGS.

ARUNDINELLA Raddi

Arundinella hispida (Willd.) O. K.,

Ekman, Ark. f. Bot. 11, n.4 (1912) 10; idem, Ark. f. Bot. 13, n.10 (1913) 10. — A. brasiliensis Raddi, Agrostol. brasiliensis () 37, tab. 1, fig. 3; Döll, FB 2,2 (1877) 298, tab. 38.

2873 RGS, Schwabenschneis p. Novo Hamburgo, NES, in alte dumosis, 14.9.1937, fr., det. Parodi.

35144 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis dumosis, 15.1.1947, fl., det. Parodi.

60340 RGS, Estação de Silvicultura p. Sta Maria, CL, in humidis alte graminosis, 8.4.1956, fr., leg. O. Camargo.

60341 RGS, ibidem, 8.4.1956, fr., leg. Camargo 570.

60342 RGS, ibidem, in humidis, 8.4.1956, fl., leg. O. Camargo 575.

Habit: Up to 2 m high.

Habitat: More or less wet, shrubby places.

Distribution in RGS: Widely scattered, probably over the whole of the northern and central Campos area of the State.

General distribution: From northern South America to RGS.

AXONOPUS Hook.

Axonopus affinis Chase,

743 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus humidis, 5.12.1932, fr., det. Beetle (Parodi, Chase: A. fissifolius (Raddi) Kuhn).

29045 RGS, ibidem, 25.8.1945, fl., det. Beetle.

29045 RGS, ibidem, 25.8.1945, fl., det. Beetle.

- 36450** RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus dense graminosis, 2.1948, fl., det. Swallen.
- 38524** RGS, S. Leopoldo, NEL, ad viam ferream in humidis, 4.12.1948, fr., det. Swallen.
- 38641** RGS, Morro da Polícia p. P. Alegre, NEL, in humidis graminosis, 22.7.1948, fr.
- 38786** RGS, Conceição p. Caí, NEH, in humidis arenosis, 14.12.1948, fr., det. Swallen.
- 39121** RGS, Esteio p. P. Alegre, NEL, in humidis graminosis, 22.12.1948, fl., det. Swallen.
- 39923** RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in humidis alte graminosis, 12.1.1949, fr., det. Swallen.
- 41071** RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in humidis graminosis, 18.4.1949, fr., Swallen.
- 41185** RGS, Ilha das Flores p. P. Alegre, NEL, in humidis graminosis, 22.4.1949, fr., det. Swallen.
- 41333** RGS, Morretes p. P. Alegre, NEL, in humidis graminosis, 2.5.1949, fr., det. Swallen.
- 41794** RGS, Esteio p. S. Leopoldo, NEL, in humidis graminosis, 1.6.1949, fl.
- 44644** RGS, Vila Oliva p. Caxias, NEH, in humidis graminosis, 3.12.1949, fr.
- 52745** RGS, Belém Novo p. P. Alegre, NEL, in campestribus arenosis, 13.7.1952, fr. deciso, leg. et det. Beetle 1570.
- 55409** SC, Massiambú p. Palhoça, in arenosis, 5.2.1953, fl., leg. R. Reitz 7089, det. Swallen.
- 55413** SC, Serra da Boa Vista p. S. José, in campestribus dense graminosis 2.2.1953, fl. R. Reitz, 7085 det. Swallen.
- 56772** RGS, Vila Oliva p. Caxias, NEH, in pascuo, 8.2.1955, fl.
- 58044** SC, Caixa d'Água p. Itajaí, in humidis, 18.3.1954, fr., leg. R. Reitz 8309, det. Swallen.
- 59387** RGS, Farroupilha, NEH, in alte graminosis, 25.11.1956, fl., leg. O. Camargo 1007.

Habit: Loose bunches, up to 40 cm high.

Habitat: Wet places.

Distribution in RGS: Widely distributed, apparently over the whole of the open area.

General distribution: From Paraná to RGS.

Axonopus brasiliensis (Spreng.) Kuhlm.,
Reitz, Sellowia 7 (1956) 97.

31791 SC, Sombrio p. Araranguá, in arenosis graminosis, 7.2.1946, fr., det. Beetle.

31892 SC, ibidem, 10.10.1945, fr., leg. R. Reitz 1293, det. Swallen.

Habit: Loose bunches, leaves 20, stalks 40 cm high.

Habitat: Dry, sandy places.

Distribution in RGS: Northeasternmost sea coast.

General distribution: Paraná to RGS.

Axonopus compressus (Swartz) Beauv.,

1015 RGS, S. Leopoldo, ad silvam campestrem in dumosis, 15.10.1934, fr., leg. C. Orth SJ, det. Chase.

6998 SC, Lages, in umbrosis graminosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

9859 RGS, Tupanciretan, NWH, in umbrosis, 29.1.1942, fr., det. Beetle.

28633 RGS, Nonoai ad fl. Uruguay, NWU, ad silvam campestrem in umbrosis, 3.1945, fl., det. Swallen.

29048 RGS, Vila Manresa p. P. Alegre, NEL, in silva campestri aperta, 25.8.1945, fr., det. Swallen.

30731 RGS, Vila Oliva p. Caxias, NEH, in araucarieto aperto, 28.1.1946, fr., det. Swallen.

30757 RGS, ibidem, 29.1.1946, fr., det. Swallen.

30848 RGS, ibidem, 31.1.1946, fl.

35177 RGS, Fazenda da Ronda p. Vacaria, NEH, in araucarieto aperto, 15.1.1947, fr.

35277 RGS, Serra da Rocinha p. Bom Jesus, NEH, in umbrosis dense graminosis, 14.2.1947, fr.

36447 RGS, Cambará p. S. Francisco de Paula, NEH, in umbrosis graminosis, 2.1948, fr., det. Swallen.

38529 RGS, S. Leopoldo, NEL, ad silvam campestrem in umbrosis, 4.12.1948, fr.

39585 RGS, Cachoeirinha p. Gravataí, NEL, in silva campestri aperta, 7.1.1949, fr., det. Swallen.

40469 RGS, Vila Manresa p. P. Alegre, NEL, in umbrosis, 14.3.1949, fr.

41792 RGS, Esteio, p. S. Leopoldo, NEL, in silva campestri aperta, 1.6.1949, det. Swallen.

46344 RGS, Cristo Rei p. S. Leopoldo, NEL, in silva campestri aperta, 17.3.1950, fr.

48353 SC, Figueiredo p. Bom Retiro, in umbrosis, 28.12.1948, fr. deciso, leg. R. Reitz 3465, det. Swallen.

55061 RGS, Vila Oliva p. Caxias, NEH, in araucarieto umbroso subhumido, det. Swallen.

56379 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in umbrosis dense graminosis, 2.1.1955, fl.

58537 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in campestribus subumbrosis, 12.2.1956, fl.

60609 RGS, Farroupilha, NEH, in graminosis, 8.5.1957, fr., leg. O. Camargo 1402.

Habit: Slender, leaves 20, stalks up to 40 cm high.

Habitat: Grassy, more or less humid and shady places at the edge of the forest.

Distribution in RGS: Apparently over the whole of the Campos area.

General distribution: South Brazil and Uruguay.

Axonopus suffultus (Mikan) Parodi,

Panicum suffultum Mikan, ex Sprengel, Neue Entdeck. 2 () 46. —

Panicum scoparium Flügge, Monogr. () 124; Döll, FB 2,2 (106).

865 RGS, Vila Manresa p. P. Alegre, (NEL) in campestribus siccis alte graminosis, 5.11.1933, fr., det. Beetle (Chase, Parodi: *A. arcuatus*) Mez.

1057 RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis siccis alte dumosis, 10.11.1934, fr., leg. C. Orth SJ, det. Parodi (Chase: *A. scoparius* (Flügge) Hitchc.).

29313 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 25.10.1945, fr.

29374 RGS, ibidem, 3.11.1945, fr.

30735 RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis alte graminosis, 28.1.1946, fr.

32989 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis dense alteque graminosis, 12.12.1945, fr.

34138 RGS, ibidem, 9.10.1946, fl.

34201 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis dense graminosis, 16.10.1946, fr.

35153 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis alte graminosis, 15.1.1947, fr.

35457 RGS, S. Leopoldo, NEL, in campestribus siccis alte graminosis, 6.11.1946, fr., leg. H. Simas.

35657 RGS, Kappesberg p. Montenegro, NES, in secundariis, 27.12.1946, fr., leg. E. Henz SJ.

38101 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 17.11.1948, fl., det. Swallen.

38140 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis alte graminosis, 20.11.1948, fr., det. Swallen.

38352 RGS, S. Leopoldo, NEL, in campestribus siccis, 4.12.1948, fr.

38704 RGS, Rio dos Sinos p. S. Leopoldo, NEL, in campestribus siccis graminosis, 10.12.1948, fr., det. Swallen.

38940 RGS, Granja Weber p. Itapoan, NEL, in siccis alte graminosis, 18.12.1948, fl., det. Swallen.

- 43592** RGS, Portão p. S. Leopoldo, NEL, in campestribus siccis alte graminosis, 23.9.1949, fr., det. Swallen.
- 44003** RGS, Morro do Osso p. P. Alegre, NEL, in campestribus siccis alte graminosis, 21.10.1949, fr., det. Swallen.
- 44092** RGS, Barreto Vianna p. S. Leopoldo, NEL, in campestribus siccis, 24.10.1949, fl., det. Swallen.
- 45409** RGS, Serra da Rocinha p. Bom Jesus, NEH, in subhumidis alte graminosis, 18.1.1950, fr.
- 50691** RGS, ibidem, 18.1.1950, fr., leg. A. Sehnem SJ 4249.
- 52739** RGS, Uruguaiana, SW, in campestribus alte graminosis, 1.8.1952, fr., leg. et det. Beetle.
- 54723** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in subhumidis alte dumosis, 1.1.1954, fr., det. Swallen.
- 57544** RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 19.11.1955, fl.
- 60080** SC, Campo dos Padres p. Bom Retiro, in subhumidis graminosis, 22.1.1957, fr.
- 61978** RGS, P. Alegre, NEL, in alte graminosis, 19.11.1957, fl., leg. O. Camargo 2590.
- 61979** RGS, ibidem, 19.12.1957, fl., leg. O. Camargo 2999.
- 61980** RGS, Taquarí, NEL, in alte graminosis, 14.12.1957, fl., leg. O. Camargo 2962.
- 62959** RGS, Lageadinho p. Bom Jesus, NEH, in alte graminosis, 20.1.1958, fl., leg. O. Camargo 3128.

Habit: Dense and large bunches, stalks up to 75 cm high.

Habitat: Dry to more or less wet Campos.

Distribution in RGS: Apparently over the whole of the Campos area, especially frequent in the northeastern Lowlands and Highlands.

General distribution: From Amazonia to Uruguay.

Axonopus ulei Hack..

Reitz, Sellowia 7 (1956) 98.

1017 RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus parce humosis, 4.4.1934, fr., leg. C. Orth SJ, det. Chase, Parodi.

6962 SC, Lages, in campestribus, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

40802 RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus alte graminosis, 1.3.1949, fr.

41016 RGS, in summo monte das Cabras p. S. Leopoldo, NEL, in rupestribus alte graminosis, 8.4.1949, fr., det. Swallen.

45451 RGS, Serra da Rocinha p. Bom Jesus, NEH, in subhumidis alte graminosis, 18.1.1950, fr.

55164 SC, Faxinal Cambajuva p. S. Joaquim, in humidis alte graminosis, 29.1.1950, fl., leg. R. Reitz 4885, det. Swallen.

Habit: Loose bunches, wiry, 40 cm high.

BRACHIARIA Gris.

Brachiaria plantaginea (Linck) Hitchc.,

Reitz, Sellowia 7 (1956) 98.

853 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 5.2.1933, fr., leg. C. Orth SJ, det. Chase.

7030 SC, Lages, in incultis efferata, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

28260 RGS, Nonoai ad fl. Uruguay, NWU, in incultis, 3.1945, fr.

40697 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 26.3.1943, fr., det. Swallen.

41370 RGS, Morretes p. P. Alegre, NEL, in incultis ad viam ferream, 2.5.1949, fr.

41482 RGS, Estação Azevedo p. Montenegro, NEL, in incultis efferata, 6.5.1949, fr., det. Swallen.

47074 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 15.5.1950, fl.

60375 RGS, Estação de Silvicultura p. Sta Maria, CL, in incultis, 8.4.1957, fr., leg. O. Camargo 610.

60630 RGS, Farroupilha, NEH, efferata, 7.5.1958, fr., leg. O. Camargo 1392.

Habit: Prostrate to ascending, stalk several m long.

Habitat: Cultivated and subs spontaneous in secondary growth.

Distribution in RGS: Over the whole of the cultivated area not too frequent.

General distribution: From southern USA to Paraguay, not native to RGS.

BRIZA L.

Briza calotheca (Trin.) Hack.,

Ekman, Ark. f. Bot. 11, n.4 (1912) 47; idem, Ark. f. Bot. 13, n.10 (1913) 52; Reitz, Sellowia 7 (1956) 98.

2511 SC, Porto União, in incultis, 29.11.1936, fl., leg. C. Orth SJ, det. Chase.

- 9870** RGS, Tupanciretan, NWH, in subpaludosis alte graminosis, 29.1.1942, fr., det. Swallen.
- 35137** RGS, Fazenda da Ronda p. Vacaria, NEH, in subpaludosis alte graminosis, 15.1.1947, fr.
- 44383** RGS, Fião p. S. Leopoldo, NEL, in subpaludosis alte dumosis, 14.11.1949, fr.
- 44636** RGS, Vila Oliva p. Caxias, NEH, in subpaludosis alte graminosis, fr. deciso.
- 49154** RGS, Esteio p. S. Leopoldo, NEL, in paludosis altissime graminosis, 20.11.1950, fr. deciso.
- 49399** RGS, Taimbesinho p. S. Francisco de Paula, NEH, in humidis alte graminosis, 18.12.1950, fr.
- 50866** Paraná, Piraquara, in paludosis alte graminosis, 30.10.1949, fr., leg. G. Hatschbach 1558.
- 57030** RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in humidis alte graminosis, 26.10.1955, fl.
- 57128** RGS, Esteio p. S. Leopoldo, NEL, in humidis alte graminosis, 14.11.1955, fr.
- 57233** RGS, ibidem, 11.11.1955, fl.
- 62769** RGS, Farroupilha, NEH, in alte graminosis, 10.11.1957, fl., leg. O. Camargo 2504.

Habit: Up to 1 m high, slender, inflorescence hanging.

Habitat: Swampy, densely vegetated places.

Distribution in RGS: Widely scattered, apparently over the whole of the swampy Campos area.

General distribution: From Paraná to Uruguay.

Briza hackelli (Lindm.) Ekm., Ark.f.Bot. 13, n.10 (1913) 61. — *Isachne* *hackelli* Lindm., Kong. Sv. Ak. Handl. 34, n.6 (1900) 83.

63381 RGS, Pelotas, SEC, in alte graminosis, 19.10.1954, fr., leg. Sacco, det. Parodi.

Habit: Isolated or in loose bunches.

Habitat: More or less dry, grassy, Campos.

Distribution in RGS: Southeastern coast, northeastern Lowlands.

General distribution: Apparently only RGS.

Briza juergensii Hack.

Reitz, Sellowia 7 (1956) 98.

44259 RGS, Sapucaia p. S. Leopoldo, NEL, in paludosis alte dumosis, 5.11.1949, fr.

48311 SC, Campo dos Padres p. Bom Retiro, in alte graminosis, 18.12.1948, fr., leg. R. Reitz 3367, det. Swallen.

48358 SC, ibidem, in paludosis, 16.12.1948, fr., leg. R. Reitz 3395, det. Swallen.

55920 RGS, Fazenda do Arroio p. Osório, NEC, in arenosis alte graminosis, 3.10.1954, fr., det. Parodi (Swallen: *B. calotheca* (Trin.) Hack.).

56317 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in alte graminosis, 2.1.1955, fr., det. Parodi.

Habit: Isolated or in thin bunches, up to 50 cm high.

Habitat: Wet, densely vegetated places.

Distribution in RGS: Northeastern quadrant of the State.

General distribution: SC and RGS.

Briza lamarckiana NEES, Agrost. Bras. (1829) 481.

43612 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus alte graminosis, 28.9.1949, fr.

43669 RGS, Morro das Abertas p. P. Alegre, NEL, in campestribus alte graminosis, 30.9.1949, fr.

44032 RGS, Morro do Osso p. P. Alegre, NEL, in campestribus alte graminosis, 21.10.1949, fr.

48827 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 22.9.1950, fr., det. Beetle.

57008 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 11.10.1955, fr.

57031 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fl.

Habit: Small bunches, wiry, up to 40 cm high.

Habitat: Dry, grassy, places.

Distribution in RGS: Apparently over the whole of the Campos area.

General distribution: RGS, Uruguay.

Briza minor L., Spec. Plant. (1753) 70; Herter, Flor. Ur. (1930) 35; Reitz, Sellowia 7 (1956) 98.

740 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 5.9.1931, fr., det. Parodi (Chase: *B. virens* Lam.).

6647 RGS, Sta Clara p. Lageado, NEL, in incultis, 18.11.1940, fr.

7445 RGS, S. Leopoldo, NEL, in incultis, 1907, fr., leg. F. Theissen S.J.

29627 RGS, sine loco, sine die, fr.

33923 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 18.9.1946, fr.

43396 RGS, Kappesberg p. Montenegro, NES, in agro, 11.9.1949, fr., det. Swallen.

43735 RGS, Morro do Sabiá p. P. Alegre, NEL, ad viam in incultis, 5.10.1949, fr.

43880 RGS, ad fl. Caí inferius p. P. Alegre, NEL, in agro, 12.10.1949, fr.

44204 RGS, Morro Santana p. P. Alegre, NEL, in incultis, 2.11.1949, fr.

47350 RGS, S. Leopoldo, NEL, ad viam in graminosis, 1947, fr., leg. A. Spies.

48989 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 18.10.1950, fr.

54407 RGS, ibidem, 12.12.1953, fr., det. Swallen.

57009 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in incultis, 11.10.1955, fr.

59633 RGS, Farroupilha, NEH, in incultis, 16.9.1956, fl., leg. O. Camargo 752.

61263 RGS, Pelotas, SEC, in incultis, 26.10.1957, fl., leg. Sacco 672.

62080 RGS, Bento Gonçalves, NEH, in incultis, 6.10.1957, fl., leg. O. Camargo 1957.

62339 RGS, Montenegro, NEL, in incultis, 19.9.1957, fr., leg. O. Camargo 1779.

Habit: Isolated or in thin bunches, up to 50 cm high.

Habitat: On disturbed soil only, often in extensive stands.

Distribution in RGS: Over the whole of the cultivated area.

General distribution: From Europe to the greater part of America.

Briza neesii Döll, FB 2,3 (1878) 132; Arechavaleta, Gram. Ur. (1894) 407; Reitz, Sellowia 7 (1956) 98.

48333 SC, Campo dos Padres p. Bom Retiro, in subpaludosis alte graminosis, 17.12.1948, fr., leg. R. Reitz 3484, det. Swallen.

50709 SC, ibidem, 16.12.1948, fr., leg. R. Reitz 3770, det. Swallen.

60078 SC, ibidem, 22.1.1957, fr.

Habit: Thin bunches, wiry, up to 1 m high.

Habitat: Wet, densely vegetated places.

Distribution in RGS: To be found on the northeastern Highlands.

General distribution: From Minas Gerais to RGS.

Briza poaeomorpha (Presl.) Henr.,

Herter, Flor. Urug. (1930) 35; Malme, Sv. Bot. Tidskr. 30 (1936) 11.

2731 RGS, S. Leopoldo, NEL, in alte dumosis humidis, 23.10.1933, fr., Leg. C. Orth SJ, det. Chase.

33206 RGS, Parecí p. Montenegro, NEL, in subpaludosis alte graminosis, 25.11.1945, fr., leg. E. Henz SJ, det. Swallen.

38340 RGS, Esteio p. S. Leopoldo, NEL, in subpaludosis altissime graminosis, 24.11.1948, fr., det. Swallen.

44114 RGS, Barreto Vianna p. S. Leopoldo, NEL, in humidis alte dumosis, 24.10.1949, fr., det. Swallen.

44345 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in humidis alte graminosis, 11.11.1949, fr.

- 44368** RGS, Fião p. S. Leopoldo, NEL, in subhumidis alte graminosis, 14.11.1949, fr.
- 44483** RGS, Granja Neugebauer p. P. Alegre, NEL, in humidis alte graminosis, 19.11.1949, fr.
- 54708** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in humidis alte graminosis, 1.1.1954, fr., det. Swallen.
- 56437** RGS, ibidem, 2.1.1955.
- 57135** RGS, Vila Manresa p. P. Alegre, NEL, in humidis, 4.11.1955, fl.
- 59458** RGS, Farroupilha, NEH, in humidis alte graminosis, 15.11.1956, fl. leg. O. Camargo 950.
- 59541** RGS, ibidem, 30.10.1956, fl., leg. O. Camargo 909.
- 62662** RGS, P. Alegre, NEL, in humidis alte graminosis, 2.11.1957, fl., leg. O. Camargo 2375.
- 62731** RGS, Farroupilha, NEH, in alte graminosis, 31.10.1957, fl., leg. O. Camargo 2337.

Habit: Very thin, threadlike, recumbent, up to 1 m long and more.

Habitat: High, grassy swamps and wet Campos.

Distribution in RGS: Apparently over the whole swampy area.

General distribution: From Minas Gerais to Uruguay.

Briza rotundata Hack.,

- 35149** RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis alte graminosis, 15.1.1947, fr., det. Parodi.

Habit: Dense, rather stiff bunches, 50 cm high.

Habitat: Dry Campos.

Distribution in RGS: Apparently northeastern Highlands only.

General distribution:

Briza rufa (Presl.) Steud.,

- B. scabra (Nees) Ekm., Ark. f. Bot. 13, n.10 (1913) 53; Herter, Flor. Urug. (1930) 35. — B. neesii Döll, FB 2,3 (1878) 132.

- 856** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 5.10.1931, fl., leg. C. Orth SJ (Chase, Parodi: B. scabra (Nees) Ekm.).

- 29322** RGS, ibidem, 26.10.1945, det. Swallen.

- 38077** RGS, ibidem, 17.11.1948, fr., det. Swallen.

- 30713** RGS, Vila Oliva p. Caxias, NEH, in subhumidis umbrosis, 28.1.1946, fr., det. Swallen.

- 34152** RGS, Faz. da Ronda p. Vacaria, NEH, in umbrosis, 15.1.1947, fr.

- 43900** RGS, ad fl. Caí inferius p. P. Alegre, NEL, ad silvam campes-trem in umbrosis, 12.10.1949, fr., det. Swallen.

57010 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 11.10.1955, fl.

57252 RGS, Vila Manresa p. P. Alegre, NEL, in subumbrosis, 15.10.1955, fl.

59527 RGS, Farroupilha, NEH, in humidis, 24.9.1956, fl., leg. O. Camargo 762.

59528 RGS, ibidem, 16.9.1956, fl., leg. O. Camargo 755.

62190 RGS, ibidem, in alte graminosis, 4.10.1957, fl., leg. O. Camargo 1885.

61828 RGS, Garibaldi, NEH, in alte graminosis, 29.10.1957, fl., leg. O. Camargo 2252.

62356 Farroupilha, NEH, 12.9.1957, in alte graminosis, fl., Leg. O. Camargo 1708.

Habit: Moderately thick bunches, up to 75 cm high.

Habitat: Dry to moderately wet places.

Distribution in RGS: Apparently over the whole of the Campos area.

General distribution: From Minas Gerais to Uruguay.

Briza stricta (Hook. et Arn.) Steud.,

11735 RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, 1943, fr.

25586 RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus dense graminosis, 1.1944, fr. deciso.

29518 RGS, Sapucaia p. S. Leopoldo, NEL, in campestribus arenosis, 5.9.1945, det. Swallen.

33198 RGS, Vila Oliva p. Caxias, NEH, in campestribus subhumidis, 16.1.1946, fr.

33382 RGS, S. Leopoldo, NEL, in campestribus alte graminosis, 10.9.1946, fr., leg. E. Henz SJ.

33914 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 18.9.1946, fr.

37743 RGS, ibidem, 22.9.1948, fr., det. Beetle.

37852 RGS, ibidem, 4.11.1948, fr., det. Swallen.

38119 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis, 20.11.1948, fr., det. Beetle.

43263 RGS, Est. Azevedo p. Montenegro, NEL, in alte graminosis, 5.9.1949, fr., det. Swallen.

43590 RGS, Portão p. S. Leopoldo, NEL, in alte graminosis, 23.9.1949, fr., det. Swallen.

43606 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus alte graminosis, 28.9.1949, det. Swallen.

43750 RGS, Morro do Sabiá p. P. Alegre, NEL, in arenosis alte graminosis, 5.10.1949, fr.

43872 RGS, ad fl. Caí inferius p. P. Alegre, NEL, in arenosis alte

- graminosis, 12.10.1949, fr., det. Swallen.
- 43980** RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis alte graminosis, 19.10.1949, fr., det. Swallen.
- 48990** RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 18.10.1950, fr., det. Beetle.
- 56229** RGS, Torres, prope litus in arenosis, 12.11.1954, fr., det. Parodi.
- 57443** RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus graminosis, 9.10.1955, fr.
- 62066** RGS, Bento Gonçalves, NEH, in alte graminosis, 6.10.1957, fl., leg. O. Camargo 1934.
- 62067** RGS, Farroupilha, NEH, in alte graminosis, 10.10.1957, fl., leg. O. Camargo 1912.
- 63361** RGS, Pelotas, SEC, in campestribus, 5.10.1954, fl., leg. J. Sacco, det. Parodi, Swallen.

Habit: Dense bunches, up to 50 cm high.

Habitat: Dry Campos.

Distribution in RGS: Widely scattered, probably over the whole of the area.

General distribution: Southern Brazil to Uruguay.

Briza subaristata Lam.,

Illustrat. des Genres 1 (1791) 187.

- 26775** RGS, Cerro Largo p. S. Luiz, NWH, in campestribus dumosis, 9.1944, fr., leg. E. Friderichs SJ, det. Swallen.
- 29323** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 26.10.1945, fr., det. Swallen.
- 34224** Chile, Mal Paso p. Valdivia, in arenosis, 15.11.1931, fr., leg. et det. H. Gunkel.
- 54411** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 12.11.1953, fr., det. Swallen.
- 54569** RGS, Taimbesinho p. S. Francisco de Paula, NEH, in paludosis alte graminosis, 13.11.1953, fl., det. Swallen.
- 55905** RGS, Fazenda do Arroio p. Osório, NEC, in arenosis graminosis, 3.10.1954, fr., det. Parodi (Swallen: B. triloba Nees).
- 57016** RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fl.
- 57122** RGS, Esteio p. S. Leopoldo, NEL, in siccis alte graminosis, 14.11.1955, fl.
- 57212** RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 22.10.1955, fl.
- 57222** RGS, ibidem, 15.10.1955, fl.
- 57228** RGS, ibidem, 15.10.1955, fl.

- 57232** RGS, ibidem, in alte graminosis, 15.10.1955, fl.
- 57234** RGS, Esteio p. S. Leopoldo, NEL, in graminosis, 11.11.1955, fl.
- 57358** RGS, Estação de Silvicultura p. Sta Maria, CL, 15.11.1955, fr., leg. O. Camargo 30.
- 59542** RGS, Farroupilha, NEH, in alte graminosis, 22.10.1956, fl., leg. O. Camargo 845.
- 62044** RGS, P. Alegre, NEL, in alte graminosis, 1.10.1957, fl., leg. O. Camargo 1838.
- 62352** RGS, Montenegro, NEL, in alte graminosis, 19.9.1957, fr., leg. O. Camargo 1764.
- 62580** RGS, Passo Fundo, CH, in campestribus siccis, 20.10.1957, fl., leg. O. Camargo 2216.
- 63365** RGS, Pelotas, SEC, in campestribus, 29.9.1954, fl., leg. J. Sacco 177, det. Parodi.
- 63366** RGS, ibidem, 5.10.1954, fl., leg. J. Sacco 186, det. Parodi.
- Habit:** Thick, rather stiff bunches, up to 50 cm high.
- Habitat:** Dry Campos.
- Distribution in RGS:** Over the whole of the Campos area.
- General distribution:** Southern Brazil and Uruguay.
- Briza triloba** Nees, Agrost. Bras. (1929) 482; Döll, FB 2,3 (1878) 133; Arechavaleta, Gram. Ur. (1894) 405; Herter, Flor. Ur. (1930) 35; Reitz, Sellowia 7 (1956) 98.
- 741** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 10.12.1932, fr., leg. C. Orth SJ, det. Parodi, Swallen.
- 7453** RGS, S. Leopoldo, NEL, in campestribus alte graminosis, 1907, fr., leg. F. Theissen SJ.
- 11770** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 1943, fr.
- 26457** RGS, Parecí p. Montenegro, NEL, in campestribus alte graminosis, 1944, fr., leg. E. Henz SJ.
- 30725** RGS, Vila Oliva p. Caxias, NEL, in campestribus graminosis, 28.1.1946, fr., det. Swallen.
- 30726** RGS, ibidem, 28.1.1946, fr., det. Swallen.
- 31798** SC, Sombrio p. Araranguá, in arenosis siccis, 8.2.1946, fr.
- 31903** SC, ibidem, 9.10.1945, fr., leg. R. Reitz 1284, det. Swallen.
- 32580** RGS, Parecí p. Montenegro, NEL, in campestribus graminosis, 26.9.1945, fr., leg. E. Henz SJ.
- 34151** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 16.10.1946, fr.
- 34203** RGS, ibidem, 16.10.1946, fr.
- 34349** RGS, Cerro Largo p. S. Luiz, NWH, in alte graminosis, 31.10.1946, fr., leg. Irm. Augusto.

35136 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus alte graminosis, 15.1.1947, fr.

35148 RGS, ibidem, 15.1.1947, fr.

43776 RGS, Parecí p. Montenegro, NEL, in campestribus alte graminosis, 7.10.1949, fr., det. Swallen.

48344 SC, Sombrio p. Araranguá, in agro, 9.10.1945, fr., leg. R. Reitz 3445, det. Swallen.

Habit: More or less thick bunches, up to 50 cm high.

Habitat: Dry Campos.

Distribution in RGS: Over the whole of the Campos area.

General distribution: From SC to Uruguay, Argentina and Chile.

Briza uniolae Nees, Agrost. Bras. (1829) 482; Döll, FB 2,3 (1878) 130; Arechavaleta, Gram. Ur. (1894) 408; Herter, Flor. Ur. (1930) 36; Ekman, Ark. f. Bot. 13, n.10 (1913) 52; Malme, Sv. Bot. Tidskr. 30 (1936) 11; Reitz, Sellowia 7 (1956) 98.

1096 RGS, S. Leopoldo, NEL, in campestribus alte graminosis, 20.10.1934, fr., leg. C. Orth SJ, det. Parodi (Chase: *B. subaristata* Lam.).

1109 RGS, ibidem, 13.10.1934, fr., leg. C. Orth SJ, det. Chase, Parodi.

9874 RGS, Tupanciretan, NWH, in siccis graminosis, 29.1.1942, fr., det. Swallen.

29324 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus graminosis, 26.10.1945, fr.

35415 RGS, S. Leopoldo, NEL, in arenosis, 18.10.1946, fr., leg. B. Wagner SJ.

37866 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus alte graminosis, 6.11.1948, fr., det. Swallen.

37946 RGS, Sapucaia p. S. Leopoldo, NEL, in campestribus arenosis, 10.11.1949, fr., det. Swallen.

38236 RGS, ibidem, 22.11.1948, fr., det. Swallen.

38342 RGS, Esteio p. S. Leopoldo, NEL, in subpaludosis alte graminosis, 24.11.1948, fr.

43897 RGS, ad f. Caí inferius p. P. Alegre, NEL, in arenosis alte graminosis, 12.10.1949, fr., det. Swallen.

44050 RGS, Barreto Vianna p. S. Leopoldo, NEL, in campestribus siccis alte graminosis, 24.10.1949, fr., det. Swallen.

44179 RGS, Morro Santana p. P. Alegre, NEL, in campestribus dumosis, 2.11.1949, fr.

44239 RGS, Sapucaia p. S. Leopoldo, NEL, in campestribus arenosis, 5.11.1949, fr., det. Beetle.

49038 RGS, Esteio p. P. Alegre, NEL, in campestribus alte graminosis, 25.10.1950, fr.

- 54567** RGS, Taimbesinho p. S. Francisco de Paula, NEH, in paludosis alte graminosis, 13.11.1953, fr., det. Swallen.
- 54713** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in humidis alte graminosis, 1.1.1954, fr., det. Swallen.
- 56001** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 19.11.1954, fr.
- 57204** RGS, ibidem, 22.10.1955, fl.
- 57242** RGS, Esteio p. S. Leopoldo, NEL, in alte graminosis, 11.11.1955, fr.
- 57441** RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus alte graminosis, 9.10.1955, fl.
- 57509** RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 15.11.1955, fr., leg. O. Camargo 22.
- 59385** RGS, Farroupilha, NEH, in dumosis alte graminosis, 15.11.1956, fl., leg. O. Camargo 992.
- 61989** RGS, Farroupilha, NEH, in alte dumosis, 18.11.1957, fl., leg. O. Camargo 2570.
- 61990** RGS, ibidem, 25.11.1957, fl., leg. O. Camargo 2637.
- 62830** RGS, P. Alegre, NEL, in campestribus siccis, 1.11.1957, fl., leg. O. Camargo 2412.
- 63368** RGS, Pelotas, SEC, in alte graminosis, 19.10.1954, fr., leg. J. Sacco 209, det. Parodi, Swallen.

Habit: Thin bunches, rather wiry, up to 1 m high.

Habitat: Dry shrubby Campos.

Distribution in RGS: Over the whole of the Campos area.

General distribution: From Paraná to Uruguay and Argentina.

BROMUS L.

Bromus auleticus Trin., ex Nees, Agrost. Bras. (1828) 468;— **B. erectus** Huds. var. **auleticus** (Trin.) Döl, FB 2,3 (1878) 109, tab. 32; Herter, Flor. Urug. (1930) 37.

- 841** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 5.10.1931, fr., leg. C. Orth SJ, det. Chase.
- 2725** RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus alte dumosis, 20.10.1935, fr., leg. C. Orth SJ, det. Chase.
- 25603** RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus dense graminosis, 1.1944, fr.
- 37757** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte dumosis, 1.10.1948, fl.
- 38099** RGS, ibidem, 17.11.1948, det. Swallen.
- 43571** RGS, Portão p. S. Leopoldo, NEL, in campestribus alte dumosis, 23.9.1949, fl., det. Swallen.

- 43611** RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus alte graminosis, 28.9.1949, fr., det. Swallen.
- 43978** RGS, Vila Manresa p. P. Alegre, 19.10.1949, fr., det. Swallen.
- 43995** RGS, Morro do Osso p. P. Alegre, NEL, in alte graminosis, 21.10.1949, fr.
- 44430** RGS, Granja Neugebauer p. Itapoan, in alte graminosis, 19.11.1949, fr.
- 48993** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus aperte dumosis, 28.11.1950, fl.
- 56000** RGS, ibidem, 19.11.1954, fl.
- 57139** RGS, ibidem, 4.11.1955, fl.
- 57501** RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 18.11.1955, fl., leg. O. Camargo 20.

Habit: Large, dense bunches, up to 1 m high.

Habitat: More or less dry Campos slopes.

Distribution in RGS: Apparently southern half of the State only.

General distribution: SC, RGS, Ur.

- Bromus brachyanthera** Döll, FB 2,3 (1878) 110: Ekman, Ark. f. Bot. 13, n. 10 (1913) 62; Reitz, Sellowia 7 (1956) 98.
- 8739** RGS, Serra da Rocinha p. Bom Jesus, NEH, in silvula nebulari, 14.1.1942, fr.
- 8748** RGS, ibidem, 14.1.1942, fr. deciso.
- 40231** RGS, Sta Rita p. Farroupilha, NEH, in araucarieto, 28.1.1949, fr.
- 44631** RGS, Vila Oliva p. Caxias, NEH, in araucarieto, 3.12.1949, fr.
- 44862** RGS, S. Francisco de Paula, NEH, in araucarieto, 18.12.1949, fl.
- 48352** SC, Figueiredo p. Bom Retiro, in dumosis humidis, 28.12.1948, fr. deciso, leg. R. Reitz, 3466, det. Swallen.
- 49315** RGS, Taimbesinho p. S. Francisco de Paula, NEH, in araucarieto, 18.12.1950, fl.
- 54479** RGS, ibidem, in umbrosis subhumidis, 13.11.1953, fl., det. Swallen.
- 56315** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in paludosis alte dumosis, 2.1.1955, fl.
- 56357** RGS, ibidem, 2.1.1955, fl.
- 59717** RGS, Farroupilha, NEH, in dumosis, 30.10.1956, fr., leg. O. Camargo 912.
- 59718** RGS, ibidem, 28.5.1956, fl., leg. O. Camargo 703.
- 59681** RGS, ibidem, 28.5.1956, fl., leg. O. Camargo 722.

Habit: Isolated or in thin, slender bunches, up to 75 cm high.

Habitat: Wet to swampy cloud and Araucaria forests.

Distribution in RGS: Northeastern Highlands.

General distribution: From Paraná to RGS.

Bromus catharticus Vahl, Symb. Bot. 2 () 22; Herter, Flor. Ur. (1930) 36; Ekman, Ark. f. Bot. 11, n.4 (1912) 47; idem, Ark. f. Bot. 13, n.10 (1913) 63. — *B. unioloides* (Willd.) Nees, Agrost. Bras. (1829) 470; Döll, FB 2,3 (1878) 110.

738 RGS, Vila Manresa p. P. Alegre, NEL, in agro inculto, 6.10.1932, fl., leg. C. Orth SJ, det. Parodi.

2907 RGS, S. Borja, SW, in incultis, 1943, fl., leg. Fr. Baglione.

2664 RGS, Parecí p. Montenegro, NEL, in incultis, 1944, fl., leg. E. Henz SJ.

26758 RGS, Cerro Largo p. S. Luiz, NWH, in agro, 9.1944, fr., leg. E. Friderichs SJ.

26824 RGS, P. Alegre, NEL, in incultis, 13.9.1944, fl., leg. P. Caparelli, det. Parodi, Swallen.

26980 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 10.1944, fr.

28338 RGS, Nonoai, NWU, in agro, 3.1945, fr.

29184 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 12.9.1945, fr., leg. Swallen.

30715 RGS, Vila Oliva p. Caxias, NEH, in campestribus, 28.1.1946, fr.

32881 RGS, Kappesberg p. Montenegro, NES, in agro, 24.10.1944, fl., leg. E. Friderichs SJ.

33357 RGS, S. Leopoldo, NEL, in incultis, 17.7.1946, fl., leg. E. Henz SJ.

33909 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 18.9.1946, fl., det. Swallen.

35497 RGS, S. Leopoldo, NEL, in agro, 2.10.1946, fl., leg. C. Ritter SJ.

43736 RGS, Morro do Sabiá p. P. Alegre, NEL, ad viam in incultis, 5.10.1949, fr., det. Swallen.

54449 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 12.11.1953, fr., det. Swallen.

55989 RGS, ibidem, 19.11.1954, fr.,

57218 RGS, ibidem, 15.10.1955, fl.

59522 RGS, Farroupilha, NEH, in dumosis alte graminosis, 17.9.1956, leg. O. Camargo 751.

61580 RGS, Taquarí, NEL, in incultis, 8.12.1957, fl., leg. O. Camargo 2690.

64341 SC, S. Joaquim, in incultis, 10.1958, fl., leg. J. Mattos.

Habit: Isolated, slender, up to 50 cm high.

Habitat: Disturbed soil.

Distribution in RGS: Over the whole of the cultivated area; not native.

General distribution: Subtropical to temperate America.

CALAMAGROSTIS Rott.

Calamagrostis armata (Döll) Parodi, (8)

- 37444 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 11.8.1948, fr.
 41559 RGS, ibidem, 14.5.1949, fr.
 43977 RGS, ibidem, 19.10.1949, fr.
 44463 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis alte graminosis, 19.11.1949, fr.
 45940 RGS, Lagoa dos Quadros p. Torres, NEC, in alte dumosis, 21.2.1950, fr.
 48986 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis graminosis, 18.10.1950, fr.
 57020 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fl.

Habit: Thin bunches, up to 75 cm high.

Habitat: More or less dry Campos.

Distribution in RGS: Northeastern Lowlands and Sea Coast.

General distribution: RGS, Ur, Argentina.

Calamagrostis beyrichiana Nees, ap. Döll, FB 2,3 (1878) 54, tab. 16; Parodi, Rev. Arg. Agr. 16, n.2 (1949) 61, fig. 4.

2729 RGS, S. Leopoldo, NEL, ad rivum in alte dumosis, 16.10.1935, fr., leg. C. Orth SJ, det. Chase.

54679 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in alte graminosis, 1.1.1954, fr., (Swallen: C. longearistata (Wedd.) Hack.).

54715 RGS, ibidem, 1.1.1954, fr.

59680 RGS, Farroupilha, NEH, in alte graminosis, 16.5.1956, fr., leg. O. Camargo 655.

59830 RGS, ibidem, 28.12.1956, fr., leg. O. Camargo 1064.

60083 SC, Campo dos Padres p. Bom Retiro, in subhumidis, 22.2.1957, fr.

64339 SC, S. Joaquim, in campestribus, 10.2.1958, fr., leg. J. Mattos.

Habit: Isolated or in small bunches, up to 75 cm high.

Habitat: More or less dry, grassy and shrubby places.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: From Rio de Janeiro to Uruguay.

Calamagrostis longearistata (Wedd.) Hack., ap. Sodiro, An. Univ. Quito 1899, 8.

(8) = **Calamagrostis alba** (Presl.) Stendel, det. A. Normann, 1974.

- 8720** RGS, Serra da Rocinha p. Bom Jesus, NEH, in campestribus subsiccis, 14.1.1942, fr., det. Swallen.
- 9746** RGS, Tupanciretan, NWH, in campestribus siccis alte graminosis, 29.1.1942, fr., det. Swallen.
- 27808** S. Paulo, Campos do Jordão, in campestribus alte graminosis, 1.1944, fr., leg. E. Friderichs SJ, det. Swallen.
- 35150** RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis alte graminosis, 15.1.1947, fr.
- 36445** RGS, Cambará p. S. Francisco de Paula, NEH, in humidis alte graminosis, 2.1948, fr.
- 36472** RGS, ibidem, 2.1948, det. Swallen.
- 44630** RGS, Vila Oliva p. Caxias, NEH, in campestribus alte graminosis, 3.12.1948, fr.
- 44643** RGS, ibidem, 3.12.1949, fr.
- 44846** RGS, S. Francisco de Paula, NEH, in campestribus alte graminosis, 18.12.1949, fr.
- 44960** RGS, Gramado p. Canela, NEH, in dumosis, 26.12.1949, fr.
- 45000** RGS, ibidem, 26.12.1949, fr. (9)
- 48322** SC, Figueiredo p. Bom Retiro, in campestribus alte graminosis, 28.12.1948, fr., leg. R. Reitz 3500, det. Swallen.

Habit: Thin bunches, up to 75 cm high.

Habitat: Dry to moderately wet Campos.

Distribution in RGS: All findplaces from the Highlands, especially the northeastern sector.

General distribution: From Ecuador to Uruguay.

Calamagrostis montevidensis Nees, Agrost. Bras. (1829) 401; Parodi, Rev. Arg. Agr. 16, n.2 (1949) 61, fig. 1.

54572 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in humidis alte graminosis, 13.11.1953, fl. (Swallen: C. longearistata (Wedd.) Hack.).

54691 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in humidis alte graminosis, 1.1.1954, fr. (Swallen: C. longearistata (Wedd.) Hack.).

56321 RGS, ibidem, 2.1.1955.

56673 RGS, Vila Oliva p. Caxias, NEH, in campestribus alte graminosis, 8.2.1955, fr.

56046 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis 19.11.1954, fl., det. Parodi (Deyeuxia spendens Brongn.; Swallen: C. viridiflavescens (Poir.) Steudel).

57216 RGS, ibidem, 15.10.1955, fl.

57238 RGS, Esteio p. S. Leopoldo, NEL, in humidis alte graminosis, 11.11.1955, fl.

57342 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 15.11.1955, fl., leg. O. Camargo 36.

58688 RGS, ibidem, in alte graminosis, 28.2.1956, fl., leg. O. Camargo 5.

61211 SC, Itapiranga ad fl. Uruguay superius, in incultis, 7.10.1957, fl.

62963 RGS, Taquarí, NEL, in campestribus, 10.12.1957, fl., leg. O. Camargo 3050.

Habit: Thin, slender bunches, up to 1 m high.

Habitat: Dry to moderately wet, more or less shrubby Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: From Peru and Bolivia to South Brazil, Uruguay, Central Argentina.

Calamagrostis viridi-flavescens (Poir.) Steud. ed. 2, 1 (1840) 251; Ekman, Ark. f. Bot. 11, n.4 (1912) 34; Ekman, Ark. f. Bot. 13., n.10 (1913) 42; Malme, Sv. Bot. Tidskr. 30 (1936) 11; Parodi, Rev. Arg. Agr. 16, n.2 (1949) 66, fig. 2; Reitz, Sellowia 7 (1956) 98.

839 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 8.2.1933, fr., leg. C. Orth SJ, det. Swallen.

2912 RGS, S. Borja, SW, in campestribus alte graminosis, 1943, fr., det. Fr. Baglione.

4913 RGS, Sta Clara p. Lageado, NEL, in graminosis, 18.11.1940, fr., det. Swallen.

6989 SC, Lages, ad viam in dumosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

30744 RGS, Vila Oliva p. Caxias, NEH, in graminosis siccis, 28.1.1946, fr.

31955 SC, Sombrio p. Araranguá, in arenosis alte graminosis, 3.12.1945, fr., leg. R. Reitz 1320, det. Beetle.

32601 RGS, Parecí p. Montenegro, NEL, in campestribus alte graminosis, 24.10.1945, fr., leg. E. Henz SJ, det. Beetle.

33492 RGS, S. Leopoldo, NEL, in campestribus alte graminosis, 17.7.1946, fr., leg. E. Henz SJ.

37945 RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis alte graminosis, 10.11.1948, fr., det. Swallen.

38100 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 18.11.1948, fr., det. Swallen.

39920 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in arenosis alte graminosis, 12.11.1949, fr.

40002 RGS, Linha Bonita p. Montenegro, in agro, 19.1.1949, fr., det. Swallen.

- 40213** RGS, Sta Rita p. Farroupilha, NEH, in incultis, 27.1.1949, fr., det. Swallen.
- 41052** RGS, Morro Santana p. P. Alegre, NEL, in dumosis siccis, 13.4.1949, fr., det. Swallen.
- 41102** RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in siccis alte graminosis, 8.4.1949, fr., det. Swallen.
- 41330** RGS, Morretes p. P. Alegre, NEL, in arenosis graminosis, 2.5.1949, fr.
- 42435** RGS, Pareci p. Montenegro, NEL, ad viam ferream in dumosis, 7.7.1949, fr., det. Swallen.
- 43899** RGS, ad fl. Caí inferius p. P. Alegre, NEL, ad viam ferream in ruderatis, 12.10.1949, fr., det. Swallen.
- 50902** Paraná, Campo Magro p. Timoneira, in siccis alte graminosis, 26.9.1950, fr., leg. G. Hatschbach 2043, det. Parodi.
- 52754** RGS, Sto Antônio da Patrulha, NEL, in dumosis ad viam, 29.6.1952, fr., leg. et det. A. Beetle.
- 53296** RGS, Caaró p. S. Luiz, NWH, in campestribus alte dumosis, 24.11.1952, fr.
- 54431** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 12.11.1953, fl., det. Swallen.
- 54958** RGS, Vila Oliva p. Caxias, NEH, in dumosis, 24.2.1954, fr.
- 55087** RGS, ibidem, 24.2.1954, fr.
- 59388** RGS, Farroupilha, NEH, in alte graminosis, 25.11.1956, fl., leg. O. Camargo 1026.
- 59523** RGS, ibidem, 26.10.1956, fl., leg. O. Camargo 863.
- 59831** RGS, ibidem, 25.3.1957, fl., leg. O. Camargo 1321.
- 60814** RGS, ibidem, 5.6.1957, fl., leg. O. Camargo 1567.
- 61966** RGS, ibidem, 25.11.1957, fl., leg. O. Camargo 2633.
- 62748** RGS, Garibaldi, NEH, in alte graminosis, 29.10.1957, fl., leg. O. Camargo 2316.

Habit: Isotated or in thin bunches, up to 1,5 m high.

Habitat: Dry to moderately wet, open or shrubby Campos.

Distribution in RGS: Over the whole of the Campos area.

General distribution: From Rio de Janeiro to Uruguay and B. Aires.

CENCHRUS L.

Cenchrus echinatus L., Spec. Plant. ed. 2 (1762) 1150; Döll, FB 2,3 (1877) tab. 63; Arechavaleta, Gram. Urug. (1894) 155; Ekman, Ark. f. Bot. 11, n.4 (1912) 28; Herter, Flor. Ur. (1930) 28; Reitz, Sellowia 7 (1956) 98.

861 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 5.3.1933, fr., leg. C. Orth SJ, det. Chase.

- 2904** RGS, S. Borja, SW, in incultis, 1943, fr., leg. F. Baglione.
- 7438** RGS, S. Leopoldo, NEL, in arenosis siccis, 1907, fr., leg. F. Theissen SJ.
- 3708** RGS, S. José do Hortênsio, NES, in incultis ad viam, 3.1.1941, fr.
- 28875** SC, Trindade p. Florianópolis, in arenosis incultis, 18.3.1945, fr., leg. A. Rohr SJ.
- 29583** RGS, Parecí p. Montenegro, NEL, in incultis arenosis siccis, 26.9.1945, fr., leg. E. Henz SJ.
- 31401** RGS, Vila Oliva p. Caxias, NEH, in incultis ad viam, 31.3.1946, fr.
- 38434** RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 1.12.1948, fr.
- 41987** RGS, Morro Santana p. P. Alegre, NEL, in agro, 15.6.1949, fl.
- 58608** RGS, Estação de Silvicultura p. Sta Maria, CL, in incultis, 10.1.1956, fr., leg. O. Camargo 25.
- 61965** RGS, Taquarí, NEL, in incultis, fr., 8.12.1957, fr., leg. O. Camargo 2685.

Habit: Radiating from the fixing point, prostrate to ascending.

Habitat: Dry, sandy or otherwise more or less bare places.

Distribution in RGS: Over the whole of the cultivated area, sometimes dominant weed.

General distribution: From northern South America to Uruguay.

Cenchrus myosuroides H.B.K., Nov. Gen. et Spec. 1 () 115, tab. 35; Ekman, Ark. f. Bot. 11, n.4 (1912) 28; Herter, Flor. Urug. (1930) 28; Arechavaleta, Gram. Ur. (1894) 157.

1066 RGS, S. Leopoldo, NEL, in arenosis dumosis, 20.2.1934, fr., leg. C. Orth SJ, det. A. Chase.

32587 RGS, Parecí p. Montenegro, NEL, in arenosis dumosis, 31.10.1945, fr., leg. E. Henz SJ.

33379 RGS, ibidem, 10.9.1946, fr., leg. E. Henz SJ.

38210 RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis alte graminosis, 22.11.1948, fr., det. Swallen.

40923 RGS, ad montem das Cabras p. S. Leopoldo, NEL, in arenosis alte dumosis, 8.4.1949, fr., det. Swallen.

Habit: Climbing and low shrubs, up to 2 m high.

Habitat: Dry, sandy, shrubby, places.

Distribution in RGS: Northeastern Lowlands.

General distribution: From southern USA to Uruguay and Argentina.

Cenchrus pauciflorus Benth., Bot. Voy. Sulph. () 56; Malme, Sv. Bot. Tidskr. 30 (1936) 10; Reitz, Sellowia 7 (1956) 99.

26404 RGS, Fazenda das Almas p. Palmares, NEC, in arenosis siccis, 1.1945, fr., leg. P. Buck SJ, det. R. Swallen.

55369 SC, Barra do Sul p. Araquari in arenosis, 8.4.1953, leg. R. Reitz 7157, det. Swallen.

55404 SC, Praia Braba p. Itajaí, prope litus in arenosis, 5.3.1953, fr., leg. R. Reitz 7106, det. Swallen.

63094 RGS, Pelotas, SEC, in incultis, 11.3.1956, fl., fr., leg. Schlichting.

Habit: The same as *C. echinatus*, differences in the fruit.

Habitat: Dry, sandy, Campos.

Distribution in RGS: Sea coast.

General distribution: From southern USA to Uruguay and Argentina.

CHAETOTROPIS Kunth

Chaetotropis chilensis Kunth, Rev. Gram. () 72, 271, tab. 47. —

Polypogon chilensis (Kunth) Pilger,

Herter, Flor. Urug. (1930) 22.

1717 RGS, S. Leopoldo, NEL, in incultis, 8.11.1934, fr., det. Chase.

10834 RGS, Marcelino Ramos, NWU, in dumosis, 1.1943, fr., leg. E. Friderichs SJ, det. Swallen.

38815 RGS, Conceição p. Caí, NEL, in campestribus dumosis, 14.12.1948, fl., det. Swallen.

38221 RGS, Sapucaia p. S. Leopoldo, NEL, in dumosis siccis, 22.11.1948, fl., det. Swallen.

44728 RGS, Lagoa dos Barros p. Osório, NEC, in humidis alte graminosis, 14.12.1949, fr.

59389 RGS, Farroupilha, NEH, in dumosis graminosis, 15.11.1956, fl., leg. O. Camargo 991.

Habit: Isolated, up to 75 cm high.

Habitat: Dry, more or less densely vegetated places.

Distribution in RGS: Apparently over the whole Campos area, not too frequent.

General distribution: From Chile to Argentina, Uruguay, and RGS.

CHLORIS Sw.

Chloris argentina (Hack.) Lillo et Parodi,

41809 RGS, Esteio p. S. Leopoldo, NEL, in campestribus graminosis, 1.6.1949, fr., det. Swallen.

Habit: Medium bunches, stalks up to 75 cm high.

Habitat: Dry, grassy, Campos.

Distribution in RGS: No other findplace.

General distribution:

Chloris berroi Arech., Gram. Urug. (1894) 328, lam 44; Herter, Flor. Urug. (1930) 38.

60446 RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus siccis, 8.4.1956, fl., leg. O. Camargo 556.

Habit: Moderately dense bunches, stalks up to 90 cm high.

Habitat: Dry, sandy, grassy places.

Distribution in RGS: No other findplaces.

General distribution: RGS, Uruguay.

Chloris canterai Arech., Gram. Urug. (1894) 325; Herter, Flor. Urug. (1930) 38.

25567 RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus alte graminosis, 1.1944, fr., det. Swallen.

25585 RGS, ibidem, 1.1944, fr.

Habit: Moderately dense bunches, up to 60 cm high.

Habitat: Dry Campos.

Distribution in RGS: Southeastern plains.

General distribution: RGS, Uruguay.

Chloris caribaea Spreng., Syst. Veg. 1 (1825) 295. *C. barbata* Swartz, Flor. Ind. occid. 1() 200; Döll, FB 2,3 (1878) 66.

31399 SC, Sombrio p. Araranguá, in arenosis alte graminosis, 31.1.1946, fr., det. Swallen.

39938 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in dumosis arenosis, 12.1.1949, fr.

42046 RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis dumosis, 18.6.1949, fr.

44380 RGS, Fião p. S. Leopoldo, NEL, in arenosis dumosis, 14.11.1949, fr.

46780 RGS, Fazenda do Arroio p. Osório, NEC, in arenosis dumosis, 14.4.1950, fr.

47057 RGS, Lagoa da Pinguela p. Osório, NEC, in arenosis dumosis, 8.5.1950, fr.

Habit: Moderately large bunches, stalks up to 50 cm high.

Habitat: Dry, grassy Campos.

Distribution in RGS: Apparently all over the Campos area.

General distribution: From the West Indies to Uruguay; also extra-American.

Chloris ciliata Swartz, Prodr. () 24; Döll, FB 2,3 (1878) 66; Arechavaleta, Gram. Urug. (1894) 324; Herter, Flor. Urug. (1930) 38.

3906 RGS, Morro Vigia p. Livramento, SW, in rupestribus siccis parce graminosis, 12.1.1941, fr., det. Parodi.

Habit: Loose, small bunches, stalk 25 cm high.

Habitat: Rocky, dry, grassy, places.

Distribution in RGS: Southwestern frontier.

General distribution: From the West Indies to Uruguay.

Chloris distichophylla Lag., Gen. et Spec. nov. diagn. () 4; Döll, FB 2,3 (1878) 70; Arechavaleta, Gram. Urug. (1894) 323; Ekman, Ark. f. Bot. 11, n.4 (1912) 34; idem, Ark f. Bot. 13, n.10 (1913) 46; Herter, Flor. Urug. (1930) 38; Reitz, Sellowia 7 (1956) 99.

1060 RGS, S. Leopoldo, NEL, in agro in culto, 3.2.1937, fr., leg. C. Orth SJ, det. Beetle.

4912 RGS, Sta Clara p. Lageado, CS, in agro, 18.11.1940, fr.

7450 RGS, S. Leopoldo, NEL, in campestribus alte graminosis, 1907, fr., leg. F. Theissen SJ.

26564 RGS, Parecí p. Montenegro, NEL, in campestribus alte dumosis, 1944, fr., leg. E. Henz SJ.

26772 RGS, Cerro Largo p. S. Luiz, NWH, ad viam in in cultis, 9.1944, fr., leg. E. Friderichs SJ, det. Swallen.

29474 RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis dumosis, 5.9.1945, fr., det. Swallen.

29640 RGS, Parecí p. Montenegro, NEL, in campestribus arenosis, 10.9.1945, fr., leg. E. Henz SJ, det. Swallen.

33441 RGS, S. Leopoldo, NEL, in arenosis, 17.7.1946, fr., leg. E. Henz SJ.

33457 RGS, ibidem, 18.7.1946, fl., leg. C. Ritter SJ.

38241 RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis alte graminosis, 22.11.1948, fr., det. Swallen.

38615 RGS, ad montem das Cabras p. S. Leopoldo, NEL, in dumosis arenosis, 7.12.1948, fl.

39748 RGS, Parecí p. Montenegro, NEL, in arenosis dumosis, 14.1.1949, fr.

40000 RGS, Linha Bonita p. Montenegro, NES, in in cultis, 19.1.1949, fr., det. Swallen.

40539 RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis dumosis, 17.3.1949, fr.

41446 RGS, Estação Azevedo p. Montenegro, NEL, in arenosis graminosis, 6.5.1949, fr., det. Swallen.

41574 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in dumosis secundariis, 16.5.1949, fr.

- 42429** RGS, Parecí p. Montenegro, NEL, in campestribus dumosis, 7.7.1949, fr., det. Swallen.
- 42631** Portão p. S. Leopoldo, NEL, in campestribus alte dumosis, 20.7.1949, fr., det. Swallen.
- 43274** RGS, Estação Azevedo p. Parecí, NEL, in campestribus arenosis alte graminosis, 5.7.1949, fr.
- 43827** RGS, Parecí p. Montenegro, NEL, in arenosis alte graminosis, 7.10.1949, fl.
- 44125** RGS, Barreto Viana p. S. Leopoldo, NEL, in campestribus alte graminosis, 24.10.1949, fr.
- 44240** RGS, Sapucaia p. S. Leopoldo, NEL, in dense alte graminosis, 5.11.1949, fr.
- 45942** RGS, Lagoa dos Quadros p. Torres, NEC, in arenosis graminosis, 21.2.1950, fr.
- 46451** Lagoa da Pinguela p. Osório, NEC, in arenosis subnudis, 27.3.1950, fr.
- 55403** SC, Praia Braba p. Itajaí, prope litus in arenosis, 5.3.1953, leg. R. Reitz 7109, det. Swallen.
- 57134** RGS, Esteio p. S. Leopoldo, NEL, in alte graminosis, 14.11.1955, fl.
- 58668** RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 10.1.1956, fl., leg. O. Camargo 52.
- 59170** RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus parce humosis, 3.2.1956, fl., fr.
- 60266** RGS, Farroupilha, NEH, in incultis, 14.1.1957, fr., leg. O. Camargo 1091.
- 60397** RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fr., leg. O. Camargo 573.
- 60398** RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 558.
- 60399** RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 574.
- 61986** RGS, Farroupilha, NWH, in alte graminosis, 18.11.1957, fl., leg. O. Camargo 2560.
- 61987** RGS, Taquarí, NEL, in alte graminosis, 8.12.1957, fl., leg. O. Camargo 2766.

Habit: Thin, loose bunches, stalks 50-75 cm high.

Habitat: Dry to moderately wet places, often as a weed on disturbed soil.

Distribution in RGS: Frequent over the whole of the Campos and cultivated area.

General distribution: From Bahia to Uruguay and northeastern Argentina.

Chloris dusenii Ekman, Ark. f. Bot. 10, n.17 (1911) 26, tab. 4, fig. 2; tab. 6, fig. 13.

9747 RGS, Tupanciretan, NWH, in humidis graminosis, 29.1.1942, fl., det. Swallen.

36478 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus graminosis, 2.1948, fr., det. Swallen.

Habit: Thin bunches, stalks up to 1 m high.

Habitat: Dry to moderately wet Campos among high grasses.

Distribution in RGS: Highlands.

General distribution: Paraná, RGS.

Chloris gayana Kunth, Rev. Gram. 1 () 89, 293, tab. 58.

41982 RGS, Morro Santana p. P. Alegre, NEL, in dumosis, 15.6.1949, fr., det. Swallen.

42119 RGS, Schwabenschneis p. Novo Hamburgo, NES, in dumosis secundariis, 23.6.1949, fr.

Habit: Loose, thin bunches, 30 cm high.

Habitat: Dry, more or less shrubby places.

Distribution in RGS: Northeastern Lowlands.

General distribution: Paraná to Uruguay, apparently an escape.

Chloris polydactyla Swartz, Prodr. () 26; Döll, FB 2,3 (1878) 67; Ekman, Ark. f. Bot. 11, n.4 (1912) 34; Herter, Flor. Urug. (1930) 38.

7446 RGS, S. Leopoldo, NEL, in campestribus alte dumosis, 1907, fr., leg. F. Theissen SJ, det. Swallen.

57127 RGS, Esteio p. S. Leopoldo, NEL, in alte graminosis, 14.11.1955, fl.

58928 RGS, Estação de Silvicultura p. Sta Maria, CL, in subhumidis alte graminosis, 28.2.1956, fr., leg. O. Camargo 24.

60400 RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 585.

61988 RGS, Taquarí, NEL, in alte graminosis, 10.2.1957, fl., leg. O. Camargo 2837.

Habit: Thin bunches, up to 1 m high.

Habitat: Dry to somewhat wet Campos among high grasses.

Distribution in RGS: Apparently over the whole of the southern Campos.

General distribution: From the Antilles to Uruguay.

Chloris pycnothrix Trin., De gramin. unifl. et plurifl. () 234, C. radiata Swartz, Flor. Ind. occ. 1 () 201; Döll, FB 2,3 (1878) 63; Ekman, Ark. f. Bot. 11, n.4 (1912) 34; idem, Ark. f. Bot. 13, n.10 (1913) 46.

2519 RGS, sine loco, sine die, fr., leg. C. Orth, SJ, det. Parodi.

4884 RGS, Sta Clara p. Lageado, NEL, in incultis, 18.11.1940, fl.

6691 RGS, ibidem, 18.11.1940, fr.

32769 RGS, Parecí p. Montenegro, NEL, in campestribus siccis, 14.11.1945, fr.

33022 RGS, ibidem, 1.10.1945, fl., leg. A. Strieder SJ.

Habit: Low bunches, stalks up to 30 cm high.

Habitat: Dry, grassy places.

Distribution in RGS: Apparently over the whole area, not frequent.

General distribution: From the West Indies to RGS.

***Chloris uliginosa* Hach..**

Herter, Flor. Urug. (1930) 38.

2995 RGS, Quebradente p. S. Leopoldo, NEL, in agro, 12.8.1936, fr., leg. C. Orth SJ, det. Parodi, Swallen (Chase: *C. pycnothrix* Trin.).

28259 RGS, Nonoai, NWU, in campestribus siccis, 3.1945, fr.

30719 RGS, Vila Oliva p. Caxias, in campestribus alte graminosis, 28.1.1946, fr.

35151 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus alte graminosis, 15.1.1947, fr., det. Parodi.

37874 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis graminosis, 11.11.1948, fr., det. Swallen.

43296 RGS, Estação Azevedo p. Montenegro, NEL, 5.9.1949, fr.

44469 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus alte graminosis, 19.11.1949, fr.

56433 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in campestribus graminosis, 2.1.1955, fr., det. Parodi.

57123 RGS, Esteio p. S. Leopoldo, NEL, in siccis alte graminosis, 14.11.1955, fr.

57024 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fr.

57357 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 15.11.1955, fr., leg. O. Camargo 21.

60401 RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 572.

60402 RGS, ibidem, 8.4.1956, leg. O. Camargo 571.

63387 RGS, Pelotas, SEC, in campestribus, 19.10.1954, fl., leg. H. Sacco, det. Parodi, Swallen.

Habit: Loose, thin bunches, up to 50 cm high, ears brown.

Habitat: Dry to moderately wet places.

Distribution in RGS: Over the whole of the Campos and cultivated area.

General distribution: From Paraná to RGS.

***CHUSQUEA* Kunth**

***Chusquea meyeriana* Rupr.**, ex Döll, FB 2,3 (1880) 203; Ekman, Ark. f. Bot. 11, n.4 (1912) 68.

53915 RGS, Serra do Fachinal p. S. Francisco de Paula, NEH, in silva nebulari, recinata, fl., det. Swallen.

Habit: Recumbent, up to 4 m high.

Habitat: At the edge of dense cloud woodlet.

Distribution in RGS: Northeasternmost Highlands.

General distribution: From Paraná to RGS.

Chusquea ramosissima Lindman, Kongl. Sv. Ak. Handl., 34, n.6 (1900) 24, tab. 14; Herter, Flor. Urug. (1930) 39; Parodi, Rev. Arg. Agron. 8 (1941) 338.

6682 RGS, Sta Clara p. Lageado, CS, in silva pluviali, 18.11.1940, ster.

7031 SC, Lages, in silva campestri, 1935, fr., leg. A. Bruxel.

9587 RGS, Tupanciretan, NWH, in silva campestri, 28.1.1942, ster.

10052 RGS, Ijuisinho p. Tupanciretan, NWH, in silvula campestri, 30.1.1942, ster.

42566 RGS, Parecí p. Montenegro, NEL, in silva pluviali, 1.7.1949, fl.

42670 RGS, Portão p. S. Leopoldo, NEL, in silva paludosa, 20.7.1949, fr.

42882 RGS, Schwabenschneis p. S. Leopoldo, NEL, in silva pluviali, 12.8.1949, fl., det. Swallen.

42951 RGS, Parecí p. Montenegro, NEL, in silva pluviali, 17.8.1949, fr., det. Swallen.

58615 RGS, Estação de Silvicultura p. Sta Maria, CL, in silva, 14.1.1956, ster., leg. O. Camargo 18.

59829 RGS, Farroupilha, NEH, in silva, 12.4.1957, ster., leg. O. Camargo 1270.

61544 RGS, ibidem, 22.11.1957, fl., leg. O. Camargo 2613.

61821 RGS, ibidem, 1.11.1957, fl., leg. O. Camargo 2300.

61976 RGS, ibidem, 18.11.1957, fl., leg. O. Camargo 2565.

62638 RGS, ibidem, in silva pluviali, 10.11.1957, fl., leg. O. Camargo 2483.

62640 RGS, ibidem, 10.11.1957, fl., leg. O. Camargo 2481.

62676 RGS, ibidem, 5.11.1957, fl., leg. O. Camargo 2441.

Habit: Recumbent, up to 5 m high.

Habit: Rain forest.

Distribution in RGS: Through the whole of the forested area.

General distribution: From Paraguay to RGS and Uruguay.

Chusquea tenella Nees, Linnaea 9 (1834) 492; Döll, Fb 2,3 (1880) 200; Reitz, Sellowia 7 (1956) 99.

871 RGS, Alto Feliz p. Caí, in silva pluviali, 5.3.1933, fl., det. Chase.

- 35528** RGS, Dois Irmãos p. S. Leopoldo, NES, in silva pluviali, 10.10.1946, fl.
- 37351** RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in silva, 8.7.1948, fl.
- 41560** RGS, Vila Manresa p. P. Alegre, NEL, in silva ad rivum, 15.5.1949, fl., det. Swallen.
- 41726** RGS, Schwabenschneis p. Novo Hamburgo, NES, in silva pluviali, 25.5.1949, fl.
- 42109** RGS, ibidem, 23.6.1949, fl.
- 42386** RGS, ad montem Ferrabraz p. Novo Hamburgo, NES, in silva pluviali, 5.7.1949, fl.
- 42473** RGS, Parecí p. Montenegro, NES, in silva pluviali, 7.7.1949, fl., det. Swallen.
- 47076** RGS, Vila Manresa p. P. Alegre, NEL, in silva, 15.8.1950, fl.
- 50863** Paraná, Pinhal p. Piraquara, in silva pluviali, 3.7.1949, fl., leg. G. Hatschbach, det. Parodi.

Habit: Recumbent, up to 5 m high.

Habitat: Rain forest.

Distribution in RGS: Northern forested area.

General distribution: From Paraná to RGS.

CORTADERIA Stapf

Cortaderia rudiuscula Stapf,
Reitz, Sellowia 7 (1956) 99.

50724 SC, Campo dos Padres p. Bom Retiro, in rupestribus dumosis, 20.12.1948, fr., leg. R. Reitz 3756, det. Swallen.

Habit: In all parts very smaller than the following (if really different).

Habitat: Rocky, somewhat wet places.

Distribution in RGS: To be found on the northeasternmost Highlands.

General distribution: SC, RGS.

Cortaderia selloana (Schult.) Asch. et Graebn., Syn. Mittel. Flora 2 (1900) 325; Ekman, Ark. f. Bot. 13, n. 10 (1913) 47; Herter, Flor. Urug. (1930) 36; Reitz, Sellowia 7 (1956) 99. — *Arundo selloana* Schult., Mant. 3, Add. 1 (1827) 605. — *Gynerium argenteum* Nees, Agrost. Bras. (1829) 462; Döll, FB 2,3 (1878) 49. 14,15.

844 RGS, Osório, in dumosis secundariis, 4.2.1933, fr., leg. C. Orth SJ, det. Chase. (*C. argentea* Stapf).

7146 SC, Lages, in dumosis secundariis, 1935, leg. A. Bruxel SJ, det. Swallen (*C. argentea* Stapf).

- 29945** RGS, Kappesberg p. Montenegro, NES, in secundariis, 3.1945, fl., leg. E. Friderichs SJ.
- 31310** RGS, Vila Oliva p. Caxias, NEH, in secundariis, 24.2.1946, fr.
- 45479** RGS, Serra da Rocinha p. Bom Jesus, NEH, in silvula montana, 18.1.1950, fr.
- 45666** RGS, Sta Rita p. Farroupilha, NEH, in dumosis secundariis, 7.2.1950, fr.
- 53853** RGS, Serra da Rocinha p. B. Jesus, NEH, in rupestribus subhumidis, 3.2.1953, fr.
- 54923** RGS, Vila Oliva p. Caxias, NEH, in secundariis, 24.2.1954, fr., det. Swallen.
- 58845** RGS, Estação de Silvicultura p. Sta Maria, CL, in secundariis alte dumosis, 1.3.1956, fl., leg. O. Camargo 122.
- 59736** RGS, Farroupilha, NEH, in dumosis, 28.5.1957, fl., leg. O. Camargo 699.
- 59998** RGS, ibidem, 25.1.1956, fl., leg. O. Camargo 1134.
- 60919** RGS, ibidem, 5.6.1957, fl., leg. O. Camargo 1578.

Habit: Very large bunches, up to 1 m in diameter at the base, stalks 2 m and more high.

Habitat: Native in rocky, somewhat wet places, dominant in secondary growth.

Distribution: Common over the whole area, especially in the northern secondary forest zone.

General distribution: From Minas Gerais to Uruguay and Argentina.

CYNODON Pers.

Cynodon dactylon (L.) Pers. Syn. Pl. 1 (1805) 85; Döll, FB 2,3 (1878) 77, tab. 21, fig. 3; Herter, Flor. Urug. (1930) 38; Cabrera, Fl. Alred. B. Aires (1953) 85; Reitz, Sellowia 7 (1956) 100. - *Panicum dactylon* L., Spec. Plant. (1753) 58.

851 RGS, Vila Manresa p. P. Alegre, NEL, in tritis siccis, 5.10.1933, fr., leg. C. Orth SJ, det. Chase, Parodi.

25583 RGS, Fazenda S. Gabriel, SW, in siccis tritis, 1.1944, fl., det. Swallen.

29195 RGS, Vila Manresa p. P. Alegre, NEL, in siccis tritis, 12.9.1945, fr., det. Swallen.

29522 RGS, Sapucaia p. S. Leopoldo, NEL, in siccis, 5.9.1945, fr., det. Beetle.

38122 RGS, Morro da Polícia p. P. Alegre, NEL, in siccis tritis, 20.11.1948, fr.

39462 RGS, Teresópolis p. P. Alegre, NEL, in tritis, 4.1.1949, fr.

39483 RGS, Cristal p. P. Alegre, NEL, in arenosis, 5.1.1949, fl., det. Swallen.

- 41228** RGS, ad fl. Caí inferius p. P. Alegre, NEL, in siccis graminosis, 27.4.1949, fl., det. Swallen.
- 44340** RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in arenosis siccis, 11.11.1949, fl.
- 53133** RGS, Cerro Largo p. S. Luiz, NWH, ad viam, 20.11.1952, fr.
- 55365** SC, Barra do Sul p. Araquari, in arenosis, 8.4.1953, fr., leg. R. Reitz, 7161, det. Swallen.
- 55418** SC, Palhoça p. Florianópolis, ad litus in arenosis, 22.12.1952, fr., leg. R. Reitz 7077, det. Swallen.
- 55993** RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 19.11.1954, fr.
- 57229** RGS, ibidem, in incultis, 10.10.1955, fr.
- 61581** RGS, Taquarí, NEL, in graminosis, 8.12.1957, fl., leg. O. Camargo 2689.
- 62576** RGS, Passo Fundo, CH, in tritis, 20.10.1957, fl., leg. O. Camargo 2220.

Habit: Widely spreading, prostate to ascending.

Habitat: Dry, trodden places.

Distribution in RGS: Apparently over the whole of the inhabited area.

General distribution: Cosmopolitan from tropical to temperate climates.

DANTHONIA Lam. et DC.

Danthonia cirrhata Hack.,

Herter, Flor. Urug. (1930) 35.

- 2712** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 19.11.1935, fl., det. Swallen (Chase, Parodi: *D. monata* Döll).
- 29254** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 10.10.1945, fr., det. Swallen.
- 29314** RGS, ibidem, 26.10.1945, fr., det. Swallen.
- 29380** RGS, ibidem, 3.11.1945, fl., det. Swallen.
- 30720** RGS, Vila Oliva p. Caxias, NEH, in siccis graminosis, 28.1.1946, fr., det. Swallen.
- 36409** RGS, Cambará p. S. Francisco de Paula, NEH, in graminosis saxosis, 2.1948, fr., det. Swallen.
- 37853** RGS, ibidem, 4.11.1948, fr., det. Swallen.
- 43695** RGS, Morro de Sta Teresa p. P. Alegre, NEL, in campestribus siccis, 3.10.1949, fl.,
- 44033** Morro do Osório p. P. Alegre, NEL, in campestribus siccis, 21.10.1949, fr.
- 55964** RGS, Taimbesinho p. S. Francisco de Paula, NEH, in campestribus saxosis, 3.11.1954, fl.
- 55988** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus dense graminosis, 19.11.1954, fl.

56003 RGS, ibidem, 19.11.1954, fl.

57205 RGS, ibidem, 22.10.1955, fl.

57235 RGS, Esteio p. S. Leopoldo, NEL, in graminosis, 11.11.1955, fr.

Habit: Elegant bunches, 40 cm high.

Habitat: Dry, grassy or rocky Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: RGS, Uruguay.

Danthonia dusenii Ekman,

54566 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in rivo ad rupes humidas, 13.11.1953, fl., det. Swallen.

Habit: Elegant bunches, 60 cm high.

Habitat: At wet rocks in a stream.

Distribution in RGS: Northeastern Highlands only.

General distribution: From Paraná to RGS.

Danthonia secundiflora Presl, Rel. Haenck. 1 () 255; Reitz, Sellowia 7 (1956) 100.

48303 SC, Campo dos Padres p. Bom Retiro, in campestribus uliginosis, 22.12.1948, fr., leg. R. Reitz 3418, det. Swallen.

50851 Paraná, p. Piraquara, in campestribus uliginosis, 13.11.1949, fr., leg. G. Hatschbach 1619, det. Parodi.

52606 RGS, Serra da Rocinha p. Bom Jesus, NEH, in campestribus uliginosis, 18.1.1950, fr.

Habit: Thin bunches, 50 cm high.

Habitat: Wet peaty Campos.

Distribution in RGS: Northeastern Highlands only.

General distribution: Peru; from Paraná to RGS.

Danthonia tenuifolia Döll, FB 2,3 (1878) 102, tab. 30, fig. 1.

2709 RGS, Portão, p. S. Leopoldo, NEL, in paludosis alte graminosis, 27.11.1935, fr., leg. C. Orth SJ, det. A. Chase.

8746 RGS, Serra da Rocinha p. Bom Jesus, NEH, in turfosis humidis, 14.1.1942, fr., det. Parodi.

54568 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in rivulo ad rupes humidas, 13.11.1953, fl., det. Swallen.

Habit: Elegant bunches, somewhat wiry, 40 cm high.

Habitat: Wet, rocky or peaty places.

Distribution in RGS: Northeastern Highlands.

General distribution: From Minas Gerais to RGS.

DESCHAMPSIA Beauv.

Deschampsia caespitosa (L.) Beauv., Essay Agrost. (1812) 91, 149, 160, tab. 18, fig. 3; Parodi, Darw. 8 (1949) 435, fig. 5; Reitz, Sellowia 7 (1956) 100.

2507 SC, S. Miguel p. Porto União, in humidis alte graminosis, 27.12.1936, fr., leg. C. Orth SJ, det. Chase.

50821 SC, Campo dos Padres p. Bom Retiro, in paludosis alte graminosis, 16.12.1948, fr., leg. R. Reitz 2363, det. Swallen.

54554 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in rivulo ad rupes humidas, 13.11.1953, fl., det. Swallen.

Habit: Elegant bunches, 45 cm high.

Habitat: Wet Campos and rocky, more or less flooded places.

Distribution in RGS: Northeastern Highlands.

General distribution: Temperate regions of both Hemispheres; in Brazil from Paraná to RGS.

DIGITARIA Heister

Digitaria aequiglumis (Hack. et Ar.) Parodi,

Herter, Flor. Ur. (1930) 28; — *Panicum aequiglume* Hack. et Arech., Gram. Urug. (1894) 93.

40741 RGS, Cristal p. P. Alegre, NEL, in subpaludosis, 31.3.1949, fr., det. Swallen.

Habit: Prostrate to ascending, branches 40 cm long.

Habitat: Wet places.

Distribution in RGS: Only findplace.

General distribution: From Paraná to Uruguay.

Digitaria horizontalis Willd., Enum. 1 () 92; Herter, Flor. Urug. (1930) 28. - *Panicum sanguinale* L, Spec. Plant. (1753) 57; Döll, FB2,2 (1877) 134.

52273 Paraná, Matinhos p. Paranaguá, in arenosis, 1.1.1950, fr., leg. G. Hatschbach 1691, det. Parodi.

55383 SC, Rodeio, in incultis, 11.6.1953, leg. R. Reitz 7138, det. Swallen.

58673 RGS, Estação de Silvicultura p. Sta Maria, CL, in incultis, 10.1.1956, fl., leg. O. Camargo 47.

Habit: Prostrate to ascending, branches up to 50 cm long.

Habitat: Dry, disturbed soil (always?)

Distribution in RGS: Uncertain.

General distribution: From Paraná to Uruguay.

Digitaria sacchariflora (Raddi) Henr.,

46499 RGS, Lagoa da Pinguela p. Osório, NEC, in subpaludosis alte graminosis, 27.3.1950, fr., det. Parodi.

Habit: Slender, up to 1 m high.

Habitat: Wet to swampy, shrubby, places.

Distribution in RGS: So far, no other findplace.

General distribution:

Digitaria sanguinalis (L.) Scop. Fl. Carn. 1 () 52; Herter, Flor. Urug. (1930) 28; Ekman, Ark. f. Bot. 11, n.4 (1912) 17; idem, Ark. f. Bot. 13, n.10 (1913) 22; Reitz, Sellowia 7 (1956) 100. — *Panicum sanguinale* L., Spec. Plant. (1753) 57; Döll, FB 2,2 (1877) 131.

737 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 6.12.1932, fr., leg. C. Orth SJ, det. Chase.

6995 SC, Lages, ad viam in incultis, 1935, leg. A. Bruxel SJ, det. Swallen.

29614 RGS, Pareci p. Montenegro, NEL, in agro inculto, 3.10.1945, fr., leg. E. Henz SJ.

29938 RGS, Kappesberg p. Montenegro, NES, in agro, 25.3.1945, fr., leg. E. Friderichs SJ.

31435 SC, Sombrio p. Araranguá, in incultis, 31.1.1946, fr.

32981 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 12.12.1945, fr.

37095 Chile, Cajón p. Cautín, in prato, 25.3.1946, fl., leg. et det. H. Gunkel 15873.

39904 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in agro, 12.1.1949, fl.

39086 RGS, Itapoan p. P. Alegre, NEL, in arenosis siccis, 22.12.1948, fl.

40660 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 26.3.1949, fr., det. Swallen.

40979 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in ruderatis, 10.4.1940, fr., det. Swallen.

45908 RGS, Lagoa dos Quadros p. Torres, NEC, in arenosis dumosis, 21.2.1950, fr.

51185 Paraná, Serra S. Luís p. Campo Largo, ad viam in tritis, 12.3.1950, fr., leg. G. Hatschbach 1905, det. Parodi.

52291 ad montem Anhangava p. Piraquara, Paraná, in incultis, 8.4.1951, fr., leg. G. Hatschbach 2223, det. Parodi.

55357 SC, Palhoça, p. litus in arenosis, 13.5.1953, fl., leg. R. Reitz 7169, det. Swallen.

56477 RGS, Torres, NEC, in arenosis tritis, 19.1.1955, fl.

59459 RGS, Farroupilha, NEH, in incultis, 25.11.1956, fl., leg. O. Camargo 1011.

59826 RGS, ibidem, 25.3.1957, fl., leg. O. Camargo 1176.

60378 RGS, Estação de Silvicultura p. Sta Maria, CL, in tritis arenosis, 8.4.1957, fr., leg. O. Camargo 614.

60379 RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 561.

60380 RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 606.

63096 RGS, Pelotas, SEC, in incultis, 10.3.1956, fl., leg. Schlichting.

64214 SC, S. Joaquim, in incultis, 10.1.1958, fl., leg. J. Mattos.

Habit: Widely spreading, rooting at the lower joints, prostrate-ascending.

Habitat: On disturbed soil.

Distribution in RGS: Molest weed throughout the whole of the cultivated area.

General distribution: Native to Europe; over the most of South America.

Digitaria violascens Linck, Hort. Berol. 1 () 229; Döll, FB 2,2 (1877) 130; Reitz, Sellowia 7 (1956) 100.

1023 RGS, S. Leopoldo, NEL, in humidis graminosis, 20.9.1934, fr., leg. C. Orth SJ, det. Chase.

33437 RGS, ibidem, 10.8.1946, fr., leg. E. Henz SJ.

33497 RGS, ibidem, 17.7.1946, fr., leg. E. Henz SJ.

38525 RGS, ibidem, 4.14.1948, fr.

38678 RGS, Rio dos Sinos p. S. Leopoldo, NEL, in humidis graminosis, 10.12.1948, fr.

39755 RGS, Parecí p. Montenegro, NEL, in humidis graminosis, 14.1.1949, fr., det. Swallen.

40930 RGS, ad montem das Cabras p. S. Leopoldo, NEL, in humidis alte graminosis, 8.4.1949, fr., det. Swallen.

41020 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in humidis subnudis, 10.4.1949, fr., det. Swallen.

41365 RGS, Morretes p. P. Alegre, NEL, in humidis graminosis, 2.5.1949, fr., det. Swallen.

41480 RGS, Estação Azevedo p. Montenegro, NEL, ad viam ferream in humidis, 6.5.1949, fr., det. Swallen.

42392 RGS, ad montem Ferrabraz p. N. Hamburgo, NEL, in humidis graminosis, 5.7.1949, fr.

42480 RGS, Parecí p. Montenegro, NEL, ad viam ferream in humidis, 7.7.1949, fr., det. Swallen.

42850 RGS, Morretes p. P. Alegre, NEL, in humidis, 10.8.1949, fr., det. Swallen.

45909 RGS, Lagoa dos Quadros p. Torres, NEC, in humidis graminosis, 21.2.1950, fr.

52752 RGS, Montenegro, NEL, in incultis, 3.7.1952, fr., leg. et det. Beetle.

64223 SC, S. Joaquim, in incultis, 10.1.1958, fl., leg. J. Mattos.

Habit: Erect, 50 cm high.

Habitat: Wet to swampy places.

Distribution in RGS: Apparently over the whole of the area.

General distribution: Widely distributed through South America.

ECHINOCHLOA Beauv.

Echinochloa colonum (L.) Linck, Hort. Berol. 2 (1833) 209; Herter, Flor. Urug. (1930) 28 — *Panicum colonum* L., Syst. Nat. ed. 10, 2 (1759) 870; Döll, FB 2,2 (1877) 140. Cabrera, Fl. Alred. B. Aires (1953) 94, fig. 27.

41019 RGS, ad fl. Gravataí inferius p. P. Alegre, in humidis graminosis, 10.4.1949, fr., det. Swallen.

Habit: Erect, 50 cm high.

Habitat: Wet, disturbed soil.

Distribution in RGS: So far, no other findplace.

General distribution: Cosmopolitan weed.

Echinochloa crus-galli (L.) Beauv., Ess. Agrost. (1812) 161; — *Panicum crusgalli* L., Spec. Plant. (1753) 56; Döll, FB 2,2 (1877) 140.

38295 RGS, Esteio p. S. Leopoldo, NEL, in agro humido, 24.11.1948, fl.

40624 RGS, Esteio p. S. Leopoldo, NEL, in agro humido, 23.3.1949, fr.

40744 RGS, Cristal p. P. Alegre, NEL, in subhumidis dumosis, 31.3.1949, fr.

41143 RGS, Ilha da Pintada p. P. Alegre, NEL, in subpaludosis, 20.4.1949, fr.

58993 RGS, Estação de Silvicultura p. Sta Maria, CL, in incultis humidis, 5.1.1956, fr. leg. O. Camargo 145.

59825 RGS, Farroupilha, NEH, in incultis humidis, 25.3.1957, fl., leg. O. Camargo 1202.

63083 RGS, Pelotas, SEC, in incultis, 11.3.1956, fl., leg. Schlichting.

63348 RGS, ibidem, 5.4.1955, fr., leg. J. Sacco, det. Parodi.

Habit: Erect, up to 1 m high.

Habitat: Wet to swampy soil (rice fields).

Distribution in RGS: Widely distributed weed.

General distribution: Native to Europe; adventitious in America.

Echinochloa crus-pavonis (H.B.K.) Schultes, Mant. 2 (1824) 269; Cabrera, Fl. Alred. B. Aires (1953) 96. — *Oplismenus Cruspavonis* H.B.K.,

Nov. Gen. et Spec. Plant. 1 (1815) 88. — *Panicum crusgalli* L., var. *echinatum* (Willd.) Döll, FB 2,2 (1877) 143.

1011 RGS, S. Leopoldo, NEL, in incultis humidis, 5.10.1934, fl., leg. C. Orth SJ, det. Chase.

3084 RGS, Granja Boa Vista p. P. Alegre, NEL, in plantatione Oryzae, 27.2.1938, fr., leg. C. Orth SJ, det. Parodi.

6431 SC, S. Ludgero p. Tubarão, in incultis humidis, 4.1.1941, fr., leg. C. Orth SJ, det. Swallen.

9847 RGS, Tupanciretan, NWH, in humidis dumosis, 29.1.1942, fl.

9886 RGS, ibidem, 29.1.1942, fr.

28398 RGS, Nonoai, NWU, in humidis alte graminosis, 3.1945, fr.

28892 SC, Tindade p. Florianópolis, in paludosis, 17.3.1945, fr., leg. A. Rohr SJ.

32778 RGS, Parecí p. Montenegro, NEL, in agro humido, 25.11.1945, fr., leg. E. Henz SJ.

33038 RGS, ibidem, in agro, 15.1.1946, fr., leg. A. Strider SJ.

40676 RGS, Vila Manresa p. P. Alegre, NEL, in dumosis ad rivum, 26.3.1949, fr., det. Swallen.

40745 RGS, Cristal p. P. Alegre, NEL, in humidis subdumosis, 31.3.1949, fr.

40965 RGS, ad fl. Gravataí p. P. Alegre, NEL, in humidis dumosis, 10.4.1949, fr., det. Swallen.

41211 RGS, Morro Santana p. P. Alegre, NEL, in dumosis ad rivum, 4.1949, fr., leg. G. Leal.

41456 RGS, Estação Azevedo p. Montenegro, NEL, in humidis dumosis, 6.5.1949, fr., det. Swallen.

45862 RGS, Lagoa dos Quadros p. Torres, NEC, ad lacum in paludosis, 21.2.1950, fr.

47023 RGS, Lagoa da Pinguela p. Osório, NEC, in humidis dumosis, 8.5.1950, fr.

49720 RGS, Lagoa dos Quadros p. Torres, NEC, in paludosis dumosis, 18.1.1951, fr.

49918 SC, Itapiranga ad fl. Uruguay superius, ad flumen, in alte dumosis, 6.2.1951, fr.

50252 RGS, Vila Manresa p. P. Alegre, NEL, in agro humido, 26.3.1951, fr.

Habit: Erect, up to 1 m high.

Habitat: Wet, disturbed soil.

Distribution in RGS: Common weed in rice fields.

General distribution: Native to tropical America, widely scattered over all tropical and subtropical regions.

Echinochloa spectabilis (Nees) Linck,

Herter, Flor. Urug. (1930) 28. — *Panicum spectabile* Nees, Agrost. Bras. (1829) 262; Döll, FB 2,2 (1877) 262, tab. 22.

31457 SC, Sombrio p. Araranguá, in paludosis alte dumosis, 1.2.1946, fr., det. Swallen.

41000 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in paludosis alte dumosis, 10.4.1949, fl., det. Swallen.

Habit: Very vigorous, several meter high.

Habitat: Dense, swampy thickets.

Distribution in RGS: Northeastern Lowlands to sea coast.

General distribution: From the West Indies to RGS.

ELEUSINE Gaertn.

Eleusine indica (L.) Gaertn., Fruct. et Sem. 1 (1788) 8; Döll, FB 2,3 (1878) 86, tab. 24; Ekman, Ark. f. Bot. 11,4(1912) 39; idem, Ark. f. Bot. 13, n.10 (1913) 47; Herter, Flor. Urug. (1930) 38; Reitz, Sellowia 7 (1956) 100. — *Cynosurus indicus* L., Spec. Plant. ed. 2 (1762) 106.

7456 RGS, S. Leopoldo, NEL, in incultis graminosis, 1907, fr., leg. F. Theissen SJ. det. Swallen.

28476 Uruguay, Tacuarembó, ad silvulam in umbrosis, 12.3.1945, fr., leg. et det. B. Rosengurtt B-4761.

32657 RGS, Parecí p. Montenegro, NEL, in agro inculto, 31.10.1945, fr., leg. E. Henz SJ.

33024 RGS, ibidem, 20.11.1945, fr., leg. C. Ritter SJ.

35998 RGS, P. Alegre, NEL, in incultis, 1946, fr., leg. P. Capparelli, det. Swallen.

36984 RGS, S. Leopoldo, NEL, in incultis, 10.3.1947, fr., leg. E. Henz SJ.

40128 RGS, Guaíba p. P. Alegre, NEL, in arenosis humidis, 24.1.1949, fr.

40623 RGS, Esteio p. S. Leopoldo, NEL, in silvula aperta, 23.3.1949, fr., det. Swallen.

40713 RGS, Cristal p. P. Alegre, NEL, in arenosis humidis, 31.3.1949, fl., det. Swallen.

55352 SC, Rio Tavares in insula Florianópolis, in arenosis, 13.5.1953, fr., leg. R. Reitz 7176, det. Swallen.

57484 RGS, Estação de silvicultura p. Sta Maria, CL, in incultis, 23.11.1955, fl., leg. O. Camargo 47.

58042 SC, Caixad'Águap. Itajaí, in incultis, 18.3.1954, fl., leg. R. Reitz 8306, det. Swallen.

60377 RGS, Estação de Silvicultura p. Sta Maria, CL, in incultis, 8.4.1956, fr., leg. O. Camargo 605.

Habit: Moderate bunches, prostrate to erect.

Habitat: Dry to moderately wet, disturbed soil.

Distribution in RGS: Common weed over the whole of the cultivated area.

General distribution: Native to Asia; adventitious in all warm regions.

Eleusine tristachya (Lam.) Kunth, Illustr. Gen. 1 (1791) 203; Ekman, Ark. f. Bot. 11, n. 4 (1912) 39; Herter, Flor. Urug. (1930) 38; Cabrera, Fl. Alred. B. Aires (1953) 87; Reitz, Sellowia 7 (1956) 100.

736 RGS, Canoas p. P. Alegre, NEL, in incultis graminosis, 6.11.1931, fr., leg. C. Orth SJ, det. Parodi, Swallen.

2910 RGS, S. Borja, SW, in incultis subhumidis, 1943, fr., leg. Fr. Bagione.

3875 RGS, Morro Vigia p. Livramento, SW, in campestribus tritis, 12.1.1941, fr.

7440 RGS, S. Leopoldo, NEL, in arenosis siccis, 1907, fr., leg. F. Theissen SJ.

10844 RGS, Sta Maria, CL, in incultis, 1943, fl., leg. A. Heidler SJ.

25572 RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus graminosis, 1.1944, fr., det. Swallen.

25580 RGS, ibidem, 1.1944, fr.

29247 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 10.10.1945, fr.

32526 RGS, Parecí p. Montenegro, NEL, in agro, 24.10.1945, fr., leg. E. Henz SJ.

35169 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus graminosis, 15.1.1947, fr.

37074 Chile, Temuco p. Cautín, in arenosis, 14.2.1941, fr., leg. H. Gunkel, det. Parodi.

38314 RGS, Esteio p. S. Leopoldo, NEL, in graminosis, 24.11.1948, fr.

38768 RGS, Conceição p. Caí, NEL, in arenosis graminosis, 14.12.1948, fr.

40681 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 23.3.1949, fr.

55382 SC, Ascurra p. Rodeio, ad viam, 11.6.1953, fr., leg. R. Reitz 7139, det. Swallen.

60015 RGS, Farroupilha, NEH, in incultis, 25.3.1957, fr., leg. O. Camargo 1318.

60009 RGS, ibidem, 15.2.1957, fr., leg. O. Camargo 1150.

61781 RGS, Taquarí, NEL, in incultis, 10.12.1957, fl., leg. O. Camargo 2856.

62852 RGS, ibidem, 10.12.1957, fl., leg. O. Camargo 3066.

Habit: Loose bunches, more or less erect.

Habitat: Dry to moderately wet, disturbed soil.

Distribution in RGS: Common weed over the whole of the area.

General distribution: Native to Africa, common in warm America.

ELIONORUS H.B.

Elionorus adustus (Trin.) Ekman, Ark. f. Bot. 13, n.10 (1913) 6. — E. latiflorus Nees var. adustus (Trin.) Hack.: FB 2,3 (1883) 307. — Andropogon adustus Trin., Mém. Ac. Pétersb. ser. 6,2 () 259.

50867 Paraná, Piraquara, in campestribus siccis, 30.10.1949, fr., leg. G. Hatschbach 1549, det. Parodi.

Habit: Up to 75 cm high.

Habitat: Dry, grassy Campos.

Distribution in RGS: Probable on the Highlands.

General distribution: Minas Gerais, Paraná.

Elionorus candidus (Trin.) Hack., FB 2,3 (1883) 306; Herter, Flor. Urug. (1930) 27. — Andropogon candidus Trin., Mém. Ac. Pétersb. ser. 6,2 (1849) 260.

854 RGS, S. Leopoldo, NEL, in siccis alte graminosis, 5.6.1933, fr., leg. C. Orth SJ, det. Chase, Parodi.

29212 RGS, Vila Manresa p. P. Alegre, NEL, in siccis alte graminosis, 3.10.1945, fr.

31923 SC, Sombrio p. Araranguá, in arenosis siccis alte graminosis, 9.10.1945, fr., leg. R. Reitz 1305.

32760 RGS, Parecí p. Montenegro, NEL, in campestribus siccis alte graminosis, 17.10.1945, fr., leg. E. Henz SJ.

37418 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 5.7.1948, fr.

38102 RGS, ibidem, 17.11.1948, fr.

43569 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis dumosis, 28.19.1949, fr.

54553 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in humidis alte graminosis, 13.11.1953, fr., det. Swallen.

57217 RGS, Vila Manresa p. P. Alegre, in alte graminosis, 22.10.1955, fl.

Habit: Bunches up to 75 cm high, rigid, ears blueish.

Habitat: Dry, grassy, Campos.

Distribution in RGS: Northeastern Lowlands, sea coast and Highlands.

General distribution: South Brazil to Uruguay.

Elionorus rostratus Nees, Agrost. Bras. (1829) 357; Hackel, FB 2,3 (1883) 308, tab. 70; Ekman, Ark. f. Bot. 11,n.4 (1912) 5; Herter, Flor. Urug. (1930) 27.

880 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 5.10.1933, fr., leg. C. Orth SJ, det. Parodi.

2726 RGS, ibidem, leg. C. Orth, SJ, det. Parodi, 2.10.1935, fr.

29252 RGS, ibidem, 10.10.1945, fr.

55889 RGS, Fazenda do Arroio p. Osório, NEC, in alte graminosis, 3.10.1954, fl., det. Parodi (Swallen: E. bisetosus Hack. et Lindm.).

63364 RGS, Pelotas, SEC, in campestribus siccis graminosis, 14.10.1954, leg. J. Sacco 197, det. Parodi, Swallen.

Habit: Loose dense bunches, 30 cm high.

Habitat: Dry Campos among shrubs and other grasses.

Distribution in RGS: Northeastern Lowlands and sea coast.

General distribution: Misiones, RGS, Uruguay.

ERAGROSTIS Beauv.

Eragrostis airoides Nees, Agrost. Bras. (1829) 509; Döll, FB 2,3 (1878) 137; Ekman, Ark. f. Bot. 11,n.4 (1912) 40; idem, Ark. f. Bot. 13, n. 10 (1913) 47; Cabrera, Fl. Alred. B. Aires (1953) 60.

4645 RGS, Estreito p. Marcelino Ramos, NWU, in rupestribus dumosis, 16.2.1941, fr., det. Parodi.

30724 RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis, 28.1.1946, fr.

32218 RGS, Azulega p. S. Francisco de Paula, NEH, in campestribus siccis, 15.2.1946, fr., det. Parodi.

32967 RGS, Parecí p. Montenegro, NEL, in dumosis secundariis, 15.1.1946, fr., leg. A. Strieder SJ.

35170 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 15.1.1947, fr.

40001 RGS, Linha Bonita p. Montenegro, NES, ad viam ferream in secundariis 19.1.1949, fr., det. Swallen.

40215 RGS, Farroupilha, NEH, in secundariis, 27.1.1949, fr., det. Swallen.

45383 RGS, Serra da Rocinha p. Bom Jesus, NEH, in campestribus subhumidis, 18.1.1950, fr.

45410 RGS, ibidem, 18.1.1950, fr.

45727 RGS, Sta Rita p. Farroupilha, NEH, in incultis, 7.2.1950, fl., det. Swallen.

45851 RGS, Lagoa dos Quadros p. Torres, NEC, in arenosis alte graminosis, 21.2.1950, fr.

54965 RGS, Vila Oliva p. Caxias, NEH, in humidis alte graminosis, 24.2.1954, fr., det. Swallen.

62065 RGS, Bento Gonçalves, NEH, in ruderatis, 6.10.1957, fl., leg. O. Camargo 1960.

62768 RGS, Farroupilha, NEH, in secundariis, 10.11.1957, fl., leg. O. Camargo 2496.

Habit: Stout bunches, up to 50 cm high.

Habitat: Dry Campos, apparently over the whole of the Campos area.

Distribution in RGS:

General distribution: From southern Brazil and Misiones to Uruguay and B. Aires.

Eragrostis articulata Nees, Agrost. Bras. (1829) 502; Döll, FB 2,3 (1878) 141; Herter, Flor. Urug. (1930) 36.

6985 SC, Lages, in campestribus, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

36436 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus graminosis, 2.1948, fr., det. Swallen.

Habit: Dense bunches, up to 40 cm high.

Habitat: Dry Campos.

Distribution in RGS: Northeastern Highlands.

General distribution: From Bahia to Uruguay.

Eragrostis bahiensis(Schrad.)Schult., Mant. 2 (1824) 318, Döll, FB 2,3 (1878) 150, Ekman, Ark. f. Bot. 11,n.4 (1912) 44; idem, Ark. f. Bot. 13, n.10 (1913) 52; Malme, Sv. Bot. Tidskr. 30 (1936) 11; Herter, Flor. Urug. (1930) 36; Cabrera, Fl. Alred. B. Aires (1953) 60; Reitz, Sellowia 7 (1956) 100.

1061 RGS, S. Leopoldo, NEL, in campestribus arenosis, 10.10.1934, leg. C. Orth SJ, det. Chase.

3068 RGS, Itapuan p. P. Alegre, NEL, in arenosis siccis, 28.1.1938, fr., leg. C. Orth SJ, det. Swallen. (10)

7038 SC, Florianópolis, in arenosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen. (10).

7040 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen. (10)

31398 SC, Sombrio p. Araranguá, in arenosis siccis, 31.1.1946, fr., det. Swallen. (10)

31685 SC, ibidem, 6.2.1946, fr., det. Swallen.

31757 SC, ibidem, 7.2.1946, fr., det. Swallen.

- 38006** RGS, Barra do Ribeiro p. P. Alegre, NEL, in arenosis submobili-
bus, 14.11.1948, fr., det. Swallen.
- 38766** RGS, Conceição p. Caí, NEL, in arenosis siccis, 14.12.1948, fr.,
det. Beetle.
- 39113** RGS, Itapoan p. P. Alegre, NEL, in arenosis siccis, 22.12.1948,
fr., det. Swallen.
- 39338** RGS, Belém Novo p. P. Alegre, NEL, in arenosis siccis,
31.12.1948, fr.
- 39450** RGS, Teresópolis p. P. Alegre, NEL, in arenosis siccis,
4.1.1949, fr., det. Swallen.
- 44561** RGS, Osório, NEC, ad lacum in arenosis subhumidis,
24.11.1949, fr. (10)
- 45225** RGS, Fazenda do Arroio p. Osório, NEC, in arenosis submobili-
bus, 4.1.1950, fr. (10)
- 45880** RGS, Lagoa dos Quadros p. Torres, NEC, ad lacum in arenosis
siccis 21.2.1950, fr.
- 51729** Palmares p. Lagoa dos Patos, NEC, in arenosis, 8.1.1952, fr.
- 52744** RGS, Torres, NEC, in arenosis, 6.8.1952, fr., leg. et det. Beetle,
(10)
- 54347** SC, Laguna, in arenosis, 22.12.1951, fr., leg. R. Reitz 5045, det.
Swallen. (10)
- 54348** SC, Ibidem, 29.2.1952, fr., leg. R. Reitz 4056, det. Swallen. (10)
- 54772** RGS, Torres, NEC, prope litus in arenosis, 11.2.1954, fr. (10)
- 55355** SC, Palhoça, prope litus in arenosis, 13.5.1953, leg. R. Reitz
7172, det. Swallen.
- 55367** SC, Barra do Sul p. Araquari, in arenosis, 8.4.1953, fr., leg. R.
Reitz 7159, det. Swallen. (10)
- 55376** SC, Massiambú p. Palhoça, in arenosis, 19.12.1952, fr., leg. R.
Reitz 7150, det. Swallen. (10)
- 55402** SC, Praia Braba p. Itajaí, in arenosis, 5.3.1953, fr., leg. R. Reitz
7108, det. Swallen. (10)
- 55408** SC, Massiambú p. Palhoça, in arenosis, 5.2.1953, fr., leg. R.
Reitz 7091, det. Swallen.
- 55426** SC, ibidem, 12.3.1953, fr., leg. R. Reitz 7065, det. Swallen.
- 56453** RGS, Cidreira, NEC, in arenosis, 17.12.1958, fr. (10)
- 56469** RGS, Torres, NEC, p. litus in arenosis, 19.1.1955, fr. (10)
- 56848** SC, Arroio do Silva p. Araranguá, in arenosis submobilibus,
19.2.1955, fr. (10)
- 59824** RGS, Farroupilha, NEH, in incultis siccis, 18.3.1957, fr., leg. O.
Camargo 1162. (10)

Habit: Thin, somewhat wiry bunches, 40 cm high.

Habitat: Preferably sandy, dry places.

Distribution in RGS: With few exceptions near the sea coast.

General distribution: From Rio de Janeiro to Uruguay and B. Aires.

Eragrostis brasiliensis Hack.,

1089 RGS, S. Leopoldo, NEL, in arenosis incultis, 10.8.1934, fr., leg. C. Orth SJ, det. Chase.

9871 RGS, Tupanciretan, NWH, in campestribus siccis, 29.1.1942, fr., det. Swallen.

59732 RGS, Farroupilha, NEH, in siccis graminosis, 28.5.1956, fr., leg. O. Camargo 733. (11)

Habit: Thick bunches, up to 60 cm high.

Habitat: Dry Campos.

Distribution in RGS: Apparently widely scattered.

General distribution: RGS.

Eragrostis ciliaris (All.) Linck, Malpighia 18 (1904) 386; Herter, Flor. Urug. (1930) 36; Cabrera, Fl. Alred. B. Aires (1953) 61. — *Poa ciliaris* All., Flor. Pedem. 2 (1785) 246.

60709 RGS, Pelotas, SEC, in incultis, 10.3.1956, fr., leg. Schlichting.

Habit: Erect, middle-sized bunches.

Habitat: Disturbed soil.

Distribution in RGS: Apparently extremely rare.

General distribution: Native to Europe; RGS, Uruguay, B. Aires.

Eragrostis ciliaris (L.) Linck, Hort. Berol. () 192; Dil, FB 2,3 (1878) 155; Ekman, Ark. f. Bot. 13, n.10 (1913) 52; Herter, flor. Urug. (1930) 36; Reitz, Sellowia 7 (1956) 101.

6429 RGS, sine loco, sine die, fr., det. Swallen.

31340 RGS, sine loco, sine die, fr., det. Swallen.

45938 RGS, Lagoa dos Quadros p. Torres, NEC, in arenosis siccis, fr.

52274 Paraná, Paranaguá, in arenosis, 1.1.1950, fr., leg. G. Hatschbach 1692, det. Parodi.

58050 SC, Praia Braba p. Itajaí, in arenosis, 1.10.1953, fr., leg. R. Reitz 8326, det. Swallen.

Habit: Dense bunches, up to 40 cm high.

Habitat: Dry, sandy places.

Distribution in RGS: Northeastern sea coast.

General distribution: From Mexico to Uruguay.

Eragrostis expansa Linck, Hort. Berol. 1 () 190; Döll, FB 2,3 (1878) 147; Herter, Flor. Urug. (1878) 36; Reitz, Sellowia 7 (1956) 101.

9848 Tupanciretan, NEH, in campestribus alte graminosis, 29.1.1942, fr., det. Swallen.

- 25568** RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus siccis alte graminosis, 1.1944, fr., det. Swallen.
- 25697** RGS, ibidem, 1.1944, fr., det. Swallen.
- 32991** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 12.12.1945, fr., det. Beetle.
- 35168** RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus dense graminosis, 15.1.1947, fr., det. Beetle.
- 39332** RGS, Belém Novo p. P. Alegre, NEL, in campestribus arenosis, 31.12.1948, fr., det. Swallen.
- 39906** RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in campestribus arenosis, 12.1.1949, fr.
- 40987** RGS, ad fl. Gravataí p. P. Alegre, NWH, in campestribus siccis, 10.4.1949, fr., det. Swallen.
- 41075** RGS, ibidem, 18.4.1949, fr., det. Swallen.
- 41091** RGS, ibidem, 18.4.1949, fr., det. Swallen.
- 41487** RGS, Estação Azevedo p. Montenegro, NEL, in campestribus alte graminosis, 6.5.1949, fr., det. Swallen.
- 44064** RGS, Barreto Vianna p. S. Leopoldo, NEL, in campestribus alte graminosis, 24.10.1949, fr., det. Swallen.
- 46782** RGS, Fazenda do Arroio p. Osório, NEC, in arenosis subsiccis, 14.4.1950, fr.
- 55391** SC, Anitápolis p. Palhoça, in dumosis, 4.4.1953, fr., leg. R. Reitz 7123, det. Swallen.
- 59012** RGS, Sta Maria, CL, in campestribus siccis, 2.1.1956, fr., leg. O. Camargo 126.
- 59024** RGS, ibidem, 12.2.1956, fr., leg. O. Camargo 167.
- 59314** RGS, ibidem, 10.3.1956, fr., leg. O. Camargo 21.
- 60396** RGS, ibidem, 8.4.1957, fr., leg. O. Camargo 616.

Habit: Thin, somewhat stiff, bunches, stalks up to 60 cm high.

Habitat: Dry Campos.

Distribution in RGS: Over the whole of the Campos area.

General distribution: From Paraná to Uruguay.

Eragrostis flaccida Lindman,
Herter, Flor. Urug. Addit.

- 4135** RGS, Rio Garopá p. Quaraí, in campestribus siccis, 14.1.1941, fr., det. Swallen.
- 28427** RGS, Nonoai, NWU, in campestribus siccis, 3.1945, fr.
- 29190** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 12.9.1945, fr., det. Swallen.
- 32753** RGS, Pareci p. Montenegro, NEL, in agro derelicto, 31.10.1945, fr.
- 38535** RGS, S. Leopoldo, NEL, in arenosis siccis, 4.12.1948, fr., det. Swallen.

40849 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis, 3.4.1949, fr.

52753 RGS, Caxias, NEH, ad viam, 8.7.1952, fr., leg. et det. Beetle 1523.

Habit: Dense bunches, up to 40 cm high.

Habitat: Dry Campos.

Distribution in RGS: Widely scattered, apparently over the whole of the Campos area.

General distribution: RGS and Uruguay.

Eragrostis hypnoides (Lam.) B.S.P., Prelim. Cat. N. Y. (1888) 69; Ekman, Ark. f. Bot. 11, n.4 (1912) 44; Herter, Flor. Urug. Addit.; Reitz, Sellowia 7 (1956) 101. — *E. reptans* Nees, Agrost. Bras. (1829) 514; Döll, FB 2,3 (1878) 148.

1907 RGS, Rio dos Sinos p. S. Leopoldo, NEL, in arenosis humidis repens, 13.3.1935, fr., leg. C. Orth SJ, det. Chase.

7441 RGS, ibidem, 1907, fr., leg. F. Theissen SJ.

26128 RGS, Fazenda do Jaraú p. Quaraí, SW, in humidis repens, 1.1945, fl., det. Swallen.

40081 RGS, Vila Manresa p. P. Alegre, NEL, in humidis repens, 22.1.1949, fr., det. Swallen.

40994 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in humidis repens, 10.4.1949, fr., det. Swallen.

41156 RGS, Ilha da Pintada p. P. Alegre, NEL, in humidis repens, 20.4.1949, fr., det. Swallen.

45737 RGS, Sta Rita p. Farroupilha, NEH, in humidis ad lacum, 7.2.1950, fr.

49729 RGS, Lagoa dos Quadros p. Torres, NEC, in arenosis ad lacum, 18.1.1951, fr.

Habit: Prostrate, rooting at the lower joints.

Habitat: Wet, sandy places.

Distribution in RGS: Over the whole of the Campos area, nowhere common.

General distribution: From Texas and Central America to Uruguay and B. Aires.

Eragrostis lugens Nees, Agrost. Bras. (1829) 505; Döll, FB 2,3 (1878) 140; Ekman, Ark. f. Bot. 11 (1912) 43; Herter, Flor. Urug. (1930) 36; Cabrera, Fl. Alred. B. Aires (1953) 61.

735 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 5.10.1932, fr., leg. C. Orth SJ, det. Parodi.

1069 RGS, ibidem, 5.11.1932, fr., leg. C. Orth SJ, det. Parodi.

- 30228** Uruguay, Juan Jackson p. Soriano, in campestribus siccis, 4.1942, fr., leg. et det. B. Rosengurtt, PE-1226.
- 52750** RGS, Dois Irmãos p. S. Leopoldo, NES, ad viam, 8.7.1952, fr., leg. et det. Beetle 1529.
- 57226** RGS, Vila Manresa p. P. Alegre, NEL, in dense graminosis, 15.10.1955, fl.
- 60395** RGS, Estação de Silvicultura p. Sta. Maria, CL, in campestribus siccis, 8.4.1946, fr., leg. O. Camargo 579.
- 60394** RGS, Estação de Silvicultura p. Sta. Maria, CL, in campestribus siccis, 8.4.1957, fr., leg. O. Camargo.
- 61999** RGS, Taquarí, NEL, in alte graminosis, 8.12.1957, fl., leg. O. Camargo 2729.

Habit. Dense bunches, up to 40 cm high.

Habitat: Dry Campos.

Distribution in RGS: Apparently over the whole of the Campos area.

General distribution: From Minas Gerais to Paraguay and Uruguay.

Eragrostis megapotamica (Spreng.) R. et Sc.,

- 6999** SC, Lages, in campestribus alte graminosis, 1935, leg. A. Bruxel SJ, det. Swallen.
- 41485** RGS, Estação Azevedo p. Montenegro, NEL, in arenosis alte dumosis 6.5.1949, fr.
- 59167** RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus parce humosis, 3.2.1956, fr.

Habit: Dense bunches, somewhat wiry, up to 50 cm high.

Habitat: Dry Campos and rocky places.

Distribution in RGS: Northeastern Lowlands.

General distribution: SC and RGS.

Eragrostis neesii Trin., Acta Petrop. 1831 405; Döll, FB 2,3 (1878) 150; Ekman, Ark. f. Bot. 13, n.10 (1913) 50; Herter, Flor. Urug. (1930) 36; Cabrera, Fl. Alred. B. Aires (1953) 61;

- 734** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus tritis, 5.10.1933, fr., leg. C. Orth SJ, det. Chase.
- 3704** RGS, Hortênsio p. Caí, NEL, in agro, 3.1.1941, fr.
- 7459** RGS, S. Leopoldo, NEL, in campestribus arenosis, 1907, fr., leg. F. Theissen SJ.
- 8443** RGS, Arroio dos Ratos p. S. Jerônimo, NEL, in campestribus siccis, 8.1.1942, fr.
- 25592** RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus siccis, 1.1944, fr.
- 25604** RGS, ibidem, 1.1944, fr.

- 26537** RGS, Parecí p. Montenegro, NEL, in campestribus siccis, 1944, fr., leg. E. Friderichs SJ.
- 26742** RGS, Cérro Largo p. S. Luiz, NWH, ad viam in incultis, 8.1944, fr., Leg. E. Friderichs SJ.
- 26948** RGS, P. Alegre, NEL, in campestribus siccis, 10.1944, fr., leg. P. Capparelli, det. Swallen (var. lindmani (Hack.) Ekm.).
- 29567** RGS, Parecí p. Montenegro, NEL, in campestribus siccis, 3.10.1945, fr.
- 29038** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 25.8.1945, fr.
- 30722** RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis, 28.1.1946, fr.
- 32765** RGS, Parecí p. Montenegro, NEL, in campestribus arenosis, 17.10.1945, fr.
- 33444** RGS, S. Leopoldo, NEL, in arenosis siccis, 10.7.1946, fr., leg. E. Henz SJ.
- 33509** RGS, ibidem, 17.7.1946, fr., leg. L. Stahl SJ.
- 35162** RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 15.1.1947, fr.
- 38061** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 17.11.1948, fr., det. Swallen.
- 38785** RGS, Conceição p. Caí, NEL, in campestribus arenosis, 14.12.1948 fr., det. Swallen.
- 54318** RGS, Fachinal Preto p. Bom Jesus, NEH, in campestribus dense graminosis, 29.1.1950, leg. R. Reitz 4996, det. Swallen.
- 54952** RGS, Vila Oliva p. Caxias, NEH, ad viam in incultis, 24.2.1954, fr.
- 57026** RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in siccis graminosis, 26.10.1955, fl.
- 57483** RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus, 23.11.1955, fl., leg. O. Camargo 48.
- 57519** RGS, ibidem, 30.11.1955, fl., leg. O. Camargo 71.
- 59371** RGS, Farroupilha, NEH, in incultis, 25.11.1956, fr., leg. O. Camargo 1028.
- 56002** RGS, Vila Manresa p. P. Alegre, NEL, in dense graminosis, 19.11.1954, fr.
- 61641** RGS, Taquarí, NEL, in campestribus, 8.12.1957, fl., leg. O. Camargo 2763.
- 61690** RGS, Farroupilha, NEH, in campestribus, 18.11.1957, fl., leg. O. Camargo 2584.
- 61847** RGS, Garibaldi, NEH, in graminosis, 29.10.1957, fl., leg. O. Camargo 2292.
- 63022** Paraguay, Primavera, in graminosis, 25.9.1955, fl., leg. A. Woolston 72.

63475 SC, S. Joaquim, in campestribus, 10.1.1958, fr., leg. J. Mattos.

Habit: Loose, rather small bunches, 10-30 cm high.

Habitat: Dry Campos and cultivated soil.

Distribution in RGS: Over the whole of the Campos area.

General distribution: From Paraná to Uruguay and B. Aires.

Eragrostis pilosa (L.) Beauv., Ess. Agrost. (1812) 71; Döll, FB 2,3 (1878) 141, tab. 42; Ekman, Ark. f. Bot. 11, n.4 (1912) 44; idem, Ark. f. Bot. 13, n.10 (1913) 49; Herter, Flor. Urug. (1930) 36; Cabrera, Fl. Alred. B. Aires (1953) 61; Reitz, Sellowia 7 (1956) 101. — **Poa ilosa** L., Spec. Plant. (1753) 68.

1007 RGS, S. Leopoldo, NEL, in tritis humidis, 2.7.1934, fr., leg. C. Orth SJ, det. Chase, Parodi (Swallen: *E. tephrosanthos* Schult.?).

2909 RGS, S. Borja, SW, in dense graminosis, 1943, fr., leg. Fr. Bagione, det. Swallen. (11)

38679 RGS, Rio dos Sinos p. S. Leopoldo, NEL, in campestribus dense graminosis, 10.12.1948, fr., det. Swallen.

41430 RGS, Estação Azevedo p. Montenegro, NEL, in campestribus siccis, 6.5.1949, fr., det. Swallen.

40746 RGS, Cristal p. P. Alegre, NEL, in arenosis humidis, 31.3.1949, fr., det. Swallen.

43734 RGS, Morro do Sabiá p. P. Alegre, NEL, in humidis alte graminosis, 5.10.1949, fr., det. Swallen.

57355 RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus alte graminosis, 23.11.1955, fl., leg. O. Camargo 45.

61998 RGS, Taquarí, NEL, in alte graminosis. 8.12.1957, fl., leg. O. Camargo 2702.

Habit: Dense bunches, up to 40 cm high.

Habitat: Dry to moderately wet Campos.

Distribution in RGS: Apparently over the whole of the Campos area.

General distribution: From Mexico to Uruguay and B. Aires.

Eragrostis polytricha Nees, Agrost. Bras. (1829) 507; Döll, FB 2,3 (1878) 139; Ekman, Ark. f. Bot. 13, n.10 (1913) 47; Herter, Flor. Urug. (1930) 37; Reitz Sellowia 7 (1956) 61.

1070 RGS, Vila Manresa p. P. Alegre, NEL in campestribus siccis, 5.11.1932, fr., det. Chase.

33350 RGS, sine loco, sine die, fr., det. Beetle.

52749 RGS, Estação Agronômica p. P. Alegre, NEL, in incultis, 11.7.1952, fr., leg. et det. Beetle.

Habit: Bunches up to 40 cm high.

Habitat: Dry Campos.

Distribution in RGS: Apparently northeastern Lowlands only.

General distribution: From S. Paulo to Uruguay.

Eragrostis trichocolea Hack. et Arch., Gram. Urug. (1894) 384; Herter, Flor. Urug. (1930) 37; Reitz, Sellowia 7 (1956) 101.

54776 RGS, Torres, NEC, prope litus in arenosis, 11.2.1954, fr., det. Swallen.

55370 SC, Barra do Sul p. Araraquá, in arenosis, 8.4.1953, fr., leg. R. Reitz 7156, det. Swallen.

63092 RGS, Pelotas, SEC, in arenosis, 10.3.1956, fl., leg. Schlichting.

Habit: Thin, rather wiry bunches from a rhizome.

Habitat: Sandy sea coast.

Distribution in RGS: Sea coast only.

General distribution: Sta. Catarina, RGS, Uruguay.

Eragrostis virescens Presl, Rel. Haenk. 1 () 276; Herter, Flor. Urug. (1930) 37.

858 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 5.10.1933, fr., leg. C. Ovth SJ, det. Parodi, Swallen.

32738 RGS, Parecí p. Montenegro, NEL, in agro derelicto, 17.10.1945, fr., leg. E. Henz SJ, det. Swallen.

Habit: Moderately thick bunches, up to 80 cm high.

Habitat: Dry Campos; cultivated soil.

Distribution in RGS: Northeastern Lowlands.

General distribution: Chile, Uruguay, RGS.

ERIANTHECIUM

Erianthecium bulbosum Parodi,

63392 RGS, Pelotas, SEC, in campestribus, 5.10.1954, fl., leg. J. Sacco 185, det. Parodi, Swallen.

Habit:

Habitat: In campestribus siccis alte graminosis.

Distribution in RGS: Near southern sea coast.

General distribution:

ERIANTHUS Michx.

Erianthus angustifolius Nees, Agrost. Bras. (1829) 316; E. saccharoides Michx., var. neesii Hack., FB 2,3 (1883) 258. — E. divaricatus (L.) Hitchc., ssp. *angustifolius* (Nees) Hack., Ekman, Ark. f. Bot. 13,n.10 (1913) 4.

1008 RGS, S. Leopoldo, NEL, in campestribus alte dumosis, 20.7.1934, fr., leg. C. Orth SJ, det. Chase.

7029 SC, Lages, in campestribus alte dumosis, 1935, fl., leg. A. Bruxel SJ, det. Swallen.

35451 RGS, S. Leopoldo, NEL, in dumosis graminosis, 6.11.1946, fl., leg. H. Simas SJ.

35765 RGS, Kappesberg p. Montenegro, NES, in alte dumosis, 20.12.1946, fr., leg. E. Henz SJ.

40004 RGS, Linha Bonita p. Montenegro, NES, in dumosis secundariis, 19.1.1949, fl.

41390 RGS, Morretes p. P. Alegre, NEL, in campestribus subhumidis alte graminosis, 2.5.1945, fl., det. Swallen.

41573 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in secundariis, 16.5.1949, fl., det. Swallen.

42451 RGS, Pareci p. Montenegro, NEL, in campestribus alte dumosis, 7.7.1949, fr., det. Swallen.

48354 SC, Figueiredo p. Bom Retiro, in campestribus alte dumosis, 28.12.1948, fl., leg. R. Reitz 3464, det. Swallen.

54998 RGS, Vila Oliva p. Caxias, NEH, in subpaludosis alte graminosis, 24.2.1954, fr., det. Swallen.

60376 RGS, Estação de Silvicultura p. Sta Maria, CL, in dumosis alte graminosis, 8.4.1956, fr., leg. O. Camargo 589.

Habit: Moderately thick bunches, up to 1,5 m high.

Habitat: More or less dry, shrubby Campos and secondary growth.

Distribution in RGS: Northern half of the State.

General distribution: From Paraguay and South Brazil do RGS.

Erianthus asper Nees, Agrost. Bras. (1829) 315; Hackel, FB 2,3 (1883) 259, tab. 61, fig. 1; Ekman, Ark. f. Bot. 11, n.4 (1912) 3; idem, Ark. f. Bot. 13, n.10 (1913) 5; Reitz, Sellowia 7 (1956) 101.

1052 RGS, S. Leopoldo, NEL, in alte dumosis, 4.3.1933, fr., leg. C. Orth SJ, det. Swallen.

31718 SC, Sombrio p. Araranguá, in paludosis alte graminosis, 7.2.1946, fr.

36464 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus dumosis 2.1948, fl., det. Swallen.

41388 RGS, Morretes p. P. Alegre, NEL, in alte dumosis, 2.5.1949, fr.

45936 RGS, Lagoa dos Quadros p. Torres, NEC, in paludosis alte dumosis, 21.2.1950, fr.

50925 Paraná, Paranaguá, prope litus in dumosis, 1.1.1950, fr., leg. G. Hatschbach 1674, det. Parodi.

54298 SC, Brusque, in dumosis secundariis, 4.1.1950, fl., leg. R. Reitz 4953, det. Swallen.

55420 SC Mafra, in paludosis alte graminosis, 26.1.1953, fr., leg. R. Reitz 7072, det. Swallen.

58863 RGS, Estação de Silvicultura p. Sta. Maria, CL, in *humidis alte graminosis*, 3.2.1956, fr., leg. O. Camargo 163.

Habit: Bunches up to 2m high.

Habitat: Moderately wet to swampy thickets and secondary growth.

Distribution in RGS: Northern half of the State.

General distribution: From Minas Gerais and Paraguay to Misiones and RGS.

Erianthus trinii Hack., Androp. in DC. Mongr. Phan. (1889) 135; Ekman, Ark. f. Bot. 11,n.4 (1912) 3; idem, Ark. f. bot. 13,n.11 (1913) 5; Herter, Flor. Urug. (1930) 27; Reitz, Sellowia 7 (1956) 101. — E. *saccharoides* Michx., var. *trinii* Hack., FB 2,3 (1883) 258.

2737 RGS, Dois Irmãos p. S. Leopoldo, in secundariis siccis, 3.7.1935, leg. C. Orth SJ, det. Chase.

7036 SC, Lages, in dumosis secundariis, 1935, fr., leg. A. Bruxel, det. Swallen.

30736 RGS, Vila Oliva p. Caxias, NEH, in dumosis secundariis, 28.1.1946, fl., det. Parodi.

30743 RGS, ibidem, in dumosis secundariis, 28.1.1946, fr., det. Parodi.

35160 RGS, Fazenda da Ronda p. Vacaria, NEH, in alte dumosis, 15.1.1947, fr.

56787 RGS, Vila Oliva p. Caxias, NEH, in paludosis alte graminosis, 8.2.1955, fr., det. Parodi.

59688 RGS, Farroupilha, NEH, in dumosis, 28.5.1956, fr., leg. O. Camargo 689.

59820 RGS, ibidem, 25.3.1957, fl., leg. O. Camargo 1205.

60807 RGS, ibidem, in dumosis, 18.6.1957, fl., leg. O. Camargo 1617.

60819 RGS, ibidem, 18.6.1957, fl., leg. O. Camargo 1633.

60820 RGS, ibidem, 18.6.1957, fl., leg. O. Camargo 1636.

63095 RGS, Pelotas, SEC, in dumosis secundariis, 10.3.1956, fl., leg. Schlichting.

Habit: Bunches 1-1,5 m high.

Habitat: Moderately dry to swampy thickets and secondary growth.

Distribution in RGS: Widely scattered over the whole of the area.

General distribution: From Bahia to RGS.

ERIOCHLOA H.B.K.

Eriochloa polystachya H. B. K., Nov. Gen. et Spec. 1 (1815) 95.

35409 RGS, S. Leopoldo, NEL, in *humidis alte graminosis*, 28.10.1946, fr., leg. E. Henz SJ.

40683 RGS, Vila Manresa p. P. Alegre, NEL, in dumosis ad rivum, 26.3.1949, fr., det. Swallen.

40999 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in subpaludosis alte dumosis, 10.4.1949, fr., det. Swallen.

41817 RGS, Canoas p. P. Alegre, NEL, in paludosis dumosis, 3.6.1949, fr., det. Swallen.

45873 RGS, Lagoa dos Quadros p. Torres, NEC, in subpaludosis alte dumosis, 21.2.1950, fr.

Habit: Erect, up to 75 cm high.

Habitat: Wet to swampy places among high grasses.

Distribution in RGS: Northeastern Lowlands and sea coast.

General distribution: From Mexico to southern Brazil.

Eriochloa punctata (L.) Hamilt., Plant. Ind. Occ. () 5; Herter, Flor. Urug. (1930) 28; Ekman, Ark. f. Bot. 13,n.10(1913) 22; Reitz, Sellowia 7 (1956) 101. — *Helopus punctatus* Nees, Agrost. Bras. (1829) 16; Döll, FB 2,2 (1877) 125.

1058 RGS, S. Leopoldo, NEL, ad viam ferream in humidis, 10.10.1934, fr., leg. C. Orth SJ, det. Chase.

32768 RGS, Pareci p. Montenegro, NEL, in agro, 10.11.1945, fr., leg. E. Henz SJ.

38450 RGS, Vila Manresa p. P. Alegre, NEL, in humidis dumosis, 1.2.1948, fr., det. Beetle.

39496 RGS, Cristal p. P. Alegre, NEL, in humidis graminosis, 5.1.1949, fr., det. Swallen.

41077 RGS, ad fl. Gravataí p. P. Alegre, NEL, in humidis dumosis, 18.4.1949, fr., det. Beetle.

40707 RGS, Cristal p. P. Alegre, NEL, in arenosis humidis, 31.3.1949, fr., det. Swallen.

40977 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in subhumidis dumosis, 10.4.1949, fl. det. Swallen.

41131 RGS, Ilha da Pintada p. P. Alegre, NEL, in humidis dumosis, 20.4.1949, fr., det. Swallen.

41184 RGS, Ilha das Flores p. P. Alegre, NEL, in humidis graminosis, 22.4.1949, fr., det. Swallen.

41339 RGS, Morretes p. P. Alegre, NEL, in arenosis dumosis, 2.5.1949, fr., det. Swallen.

55350 SC, Rio Tavares in insula Sta Catarina, prope litus in dumosis humidis, 13.5.1953, leg. R. Reitz 7178, det. Swallen.

Habit: 50 cm high.

Habitat: Wet to more or less swampy places.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Mexico to RGS.

ERIOCHRYYSIS Beauv.

Eriochrysis cayennensis Beauv. Ess. Agr. (1812) 8; Reitz, Sellowia 7 (1956) 101. — *Saccharum cayennense* Benth., Journ. Linn. Soc. 19 () 66; Hackel, FB 2,3 (1883) 253; Ekman, Ark. f. Bot. 11, n.4 (1912) 3; idem, Ark. f. Bot. 13,n.10 (1913); Herter, Flor. Urug. (1930) 27.

1054 RGS, S. Leopoldo, NEL, ad viam ferream in paludosis alte graminosis, 15.4.1934, fr., leg. C. Orth SJ, det. Chase.

7004 SC, Lages, in paludosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

7015 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

8779 RGS, Bom Jesus, NEH, in subpaludosis alte graminosis, 14.1.1942, fr.

28475 RGS, Nonoai ad fl. Uruguay, NWH, in paludosis, 3.1945, fr.

35159 RGS, Fazenda da Ronda p. Vacaria, NEH, in paludosis alte dumosis, 15.1.1947, fl., fr.

36476 RGS, Cambará p. S. Francisco de Paula, NEH, in subpaludosis, 2.1948, fl., det. Swallen.

36965 RGS, S. Leopoldo, NEL, in paludosis alte graminosis, 10.3.1947, fr., leg. E. Henz SJ.

41745 RGS, Fião p. S. Leopoldo, NEL, in paludosis alte dumosis, 29.5.1949, fr., det. Swallen.

54574 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in paludosis alte graminosis, 13.11.1953, fr. (Swallen: *Eriochrysis* sp.).

55342 SC, Mafra, in humidis alte graminosis, 26.1.1953, leg. R. Reitz 7189, det. Swallen.

56789 RGS, Vila Oliva p. Caxias, NEH, in dumosis humidis, 8.2.1955, fr.

58056 SC, Massiambú p. Palhoça, in humidis alte graminosis, 4.11.1953, leg. R. Reitz 8337, det. Swallen.

58929 RGS, Estação de Silvicultura p. Sta Maria, CL, in paludosis alte graminosis, 28.2.1956, fr., leg. O. Camargo 30.

62769 RGS, Farroupilha, NEH, in paludosis, 10.11.1957, fl., leg. O. Camargo 2499.

62958 RGS, Lageadinho p. Bom Jesus, NEH, in paludosis, 20.1.1958, fl., leg. O. Camargo 3129.

64233 SC, S. Joaquim, in paludosis, 20.1.1958, fl., leg. J. Mattos.

64270 SC, ibidem, 20.1.1958, fl., leg. J. Mattos.

Habit: Single stalks, up to 1,5 m high.

Habitat: Wet to swampy thickets among high grasses and shrubs.

Distribution in RGS: Over the whole Campos area.

General distribution: From Mexico to Uruguay and northern Argentina.

FESTUCA L.

Festuca ampliflora Döll, FB 2,3 (1878) 116, tab. 34. Ekman, Ark. f. Bot. 13, n. 10 (1913) 62.

54704 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in paludosis alte dumosis, 1.1.1954, fr., det. Swallen.

59543 RGS, Farroupilha, NEH, in paludosis alte dumosis, 26.10.1956, fl., leg. O. Camargo 822. (12)

Habit: Up to 1,5 m high.

Habitat: Swampy thickets.

Distribution in RGS: Northeastern Highlands.

General distribution: From Minas Gerais to RGS.

Festuca australis Nees, ap. Steud. Syn. Plant. Gram. (1855) 304; Malme, Sv. Bot. Tidskr. 30 (1936) 11. — **F. tenella** Willd., Spec. Plant. 1 () 419; Döll, FB 2,3 (1878) 112; Herter, Flor. Urug. (1930) 37.

2727 RGS, Morro da Polícia, NEL, in humidis subhumidis, 2.10.1935, fl., det. Case. (13)

37985 RGS, Barra do Ribeiro p. P. Alegre, NEL, in arenosis humidis, 14.11.1948, fr., det. Swallen. (13)

38151 RGS, Morro da Polícia p. P. alegre, NEL, in humidis subhumidis, 20.11.1948, fr. (13)

43608 RGS, ibidem, 28.9.1949, fr., det. Swallen.

44069 RGS, Barreto Vianna p. S. Leopoldo, NEL, in arenosis, 24.10.1949, fr.

Habit: 15 cm high.

Habitat: Dense stands on more or less wet, bare places.

Distribution in RGS: P. Alegre, S. Leopoldo.

General distribution: RGS, Uruguay.

Festuca dertonensis (All.) Aschers. et Graebn., Syn. mitteleur. Flora 2 (1900) 588. — **Bromus dertonensis** All., Fl. Pedem. 2 (1785) 249. Cabrera, Fl. Alred. B. Aires (1953) 61.

29378 Vila Manresa p. P. Alegre, NEL, in humidis graminosis, 3.11.1945, fl., det. Swallen. (14)

57143 RGS, ibidem, 4.11.1955, fl. (14)

62766 RGS, Farroupilha, NEH, in incultis, 31.10.1957, fl., leg. O. Camargo 2347. (14)

(12) = **Bryza calotheca** (Trin.) Hack, det. H. Longhi, 20.11.1975.

(13) = **Vulpia australis** (Nees ex Stendel) Blom, det. H. Longhi, 1975.

(14) = **Vulpia bromoides** S. F. Gray, det. H. Longhi, 11.11.1975.

Habit: 30 cm high, often in dense stands.

Habitat: More or less wet places on rocky ground.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: Native to Europe; RGS, Uruguay, B. Aires.

Festuca fimbriata Nees, Agrost. Bras. (1829) 472; Döll, FB 2,3 (1878) 114; Herter, Flor. Urug. (1930) 37.

35173 RGS, Fazenda da Ronda p. Vacaria, NEH, in subhumidis alte graminosis, 15.1.1947, fr. (15)

45452 RGS, Serra da Rocinha p. Bom Jesus, NEH, in paludosis alte graminosis, 18.1.1950, fr., det. Swallen. (15)

45466 RGS, ibidem, 18.1.1950, fr., det. Beetle. (15)

60082 SC, Campo dos Padres p. Bom Retiro, in humidis alte graminosis, 22.1.1957, fl.

Habit: 75 cm high.

Habitat: Wet to swampy places among high shrubs and grasses.

Distribution in RGS: Northeastern Highlands.

General distribution: RGS and Uruguay.

Festuca myurus L., Spec. Plant. (1753) 74; Cabrera, Fl. Alred. B. Aires (1953) 61.

733 RGS, Vila Manresa p. P. Alegre, NEL, in subhumidis graminosis, 20.1.1933, fr., det. Chase (14).

37096 Chile, Volc. Villa Rica p. Cautín, in arenosis, 6.12.1945, fl., leg. H. Gunckel 15721, det. Parodi (16).

59539 RGS, Farroupilha, NEH, in incultis, 22.10.1956, fr., leg. O. Camargo 846. (14)

Habit: 30-50 cm high.

Habitat: Somewhat wet places with rocky underground.

Distribution in RGS: Widely scattered.

General distribution: Native to Europe; RGS, B. Aires.

(15) = *Festuca* sp. det. H. Longhi.

(16) = *Vulpia megalura* (Nuttall) Ridberg, det. H. Longhi, 11.11.1975.

- Festuca ulochaeta** (Nees) Döll, FB 2,3 (1878) 115, tab. 33; **Vulpia ulochaeta** Nees, ex Steud. Syn. Plant. Gram. (1855) 305.
- 4420** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in araucarieto, 8.2.1941, fr.
- 6996** SC, Lages, in araucarieto, 1935, fl., leg. A. Bruxel SJ, det. Swallen.
- 7003** SC, ibidem, 1935, fl., leg. Bruxel SJ, det. Swallen.
- 30740** RGS, Vila Oliva p. Caxias, NEH, in araucarieto, 28.1.1946, fr.
- 31319** RGS, ibidem, 24.2.1946, fr.
- 35176** RGS, Fazenda da Ronda p. Vacaria, NEH, in araucarieto, 15.1.1947, f. deciso.
- 35258** RGS, Serra da Rocinha p. Bom Jesus, NEH, in silvula nebulari, 14.2.1947, fr.
- 36446** RGS, Cambará p. S. Francisco de Paula, NEH, in araucarieto, 2.1948, fr., det. Swallen.
- 44891** RGS, S. Francisco de Paula, NEH, in araucarieto, 18.12.1949, fr.
- 54696** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in umbrosis subhumidis, 1.1.1954, fl.,
- 54989** RGS, Vila Oliva p. Caxias, NEH, in humidis subumbrosis, 24.2.1954, fr., det. Swallen.
- 54999** RGS, ibidem, 24.2.1954, fr., det. Swallen.
- 55394** SC, Campo dos Padres p. Bom Retiro, in dumosis subhumidis, 20.12.1953, fl., leg. R. Reitz 7118, det. Swallen.
- 56319** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in dumosis, 2.1.1955, fl.
- 56434** RGS, ibidem, 2.1.1955, fl.
- 56725** RGS, Vila Oliva p. Caxias, NEH, in dumosis humidis, 8.2.1955, fl.
- 56768** RGS, ibidem, 8.2.1955, fr. (17)
- 59827** RGS, Farroupilha, NEH, in araucarieto, 25.3.1957, fl., leg. O. Camargo 1208.
- 59992** RGS, ibidem, 15.2.1957, fl., leg. O. Camargo 1157.
- 60081** SC, Campo dos Padres, in umbrosis humidis, 22.1.1957, fl.
- 61855** RGS, Garibaldi, NEH, in silva aperta, 29.10.1957, fl., leg. O. Camargo 2283. (17)

62765 RGS, Farroupilha, NEH, in umbrosis humidis, 31.1.1957, fl., leg. O. Camargo 2354. (17)

Habit: Slender, up to 1 m high.

Habitat: More or less wet Araucaria forest and Campos woodlets.

Distribution in RGS: Northwestern Highlands.

General distribution: From Paraná to RGS.

GUADUA Kunth

Guadua trinii Nees, Linnéa 9 (1834) 469; Döll, FB 2,3 (1880) 178. — (G. riograndensis Dutra msc.).

879 RGS, Esteio p. S. Leopoldo, 7.9.1944, fl., in silva paludosa, leg. C. Orth SJ.

4668 RGS, Sto Angelo, NWH, in silvula subpaludosa, 18.2.1941, ster.

9174 RGS, Jarí p. Tupanciretan, NWH, in silva campestri subpaludosa, 26.1.1942, ster.

28547 RGS, Nonoai ad fl. Uruguay, NWU, in silva campestri paludosa, 3.1945, ster.

53315 Cérro largo p. S. Luiz, NWH, in silva campestri paludosa, 24.11.1952, ster.

58632 RGS, Estação de Silvicultura p. Sta Maria, CL, in silva paludosa, 14.1.1956, ster., leg. O. Camargo 1.

64196 RGS, Palmeira, NWH, in silvula campestri paludosa, 1957, fl., leg. R. Beltrão.

Habit: Up to 8 m high, very spiny.

Habitat: Wet to swampy places.

Distribution in RGS: Lowlands and northeastern Highlands.

General distribution: Minas Gerais to RGS.

GYMNOPOGON Beauv.

Gymnopogon biflorus Pilger,

41026 RGS, Morro Santana p. P. Alegre, NEL, in siccis dumosis, 13.4.1949, fl., det. Swallen.

46811 RGS, Fazenda do Arroio p. Osório, NEC, in arenosis, 14.4.1950, fr.

Habit: Thin bunches, 30 cm high.

Habitat: Dry, grassy Campos.

Distribution in RGS: Northeastern Lowlands and sea coast.

General distribution:

HETEROROPOGON Pers.

Heteropogon villosus Nees, Agrost. Bras. (1829) 362; Hackel, FB 2,3 (1883) 269, tab. 62, fig. 2. — *Andropogon villosus* (Nees) Ekman.; Herter, Flor. Urug. (1930) 27.

855 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 20.10.1933, leg. C. Orth SJ, det. Chase.

29133 RGS, ibidem, 12.9.1945, fl., det. Swallen.

32757 RGS, Parecí p. Montenegro, NEL, in campestribus siccis alte graminosis, 17.10.1945, fl., leg. E. Henz SJ, det. Beetle.

33877 RGS, Taquarí, NEL, in campestribus siccis, 22.12.1945, fr., leg. P. Capparelli.

33889 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis alte graminosis, 8.11.1948, fr., det. Swallen.

35142 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 14.1.1947, fr., det. Beetle.

35166 RGS, ibidem, 15.1.1947, fr.

35172 RGS, ibidem, in campestribus alte graminosis, 15.1.1947, fr.

43624 RGS, ibidem, 28.9.1949, fl., det. Swallen.

43630 RGS, ibidem, 28.9.1949, fr., det. Swallen.

43643 RGS, Morro das Abertas p. P. Alegre, NEL, in rupestribus alte graminosis, 30.9.1949, fr.

44437 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis, 19.11.1949, fr.

48866 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis, 27.9.1950, fl.

Habit: Dense bunches, up to 50 cm high.

Habitat: Dry Campos among high brasses.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: From Mato Grosso and Minas Gerais to Uruguay.

HOLCUS L.

Holcus lanatus L., Spec. Plant. (1753) 1048; Herter, Flor. Urug. (1930) 35.

2899 RGS, S. Borja, SW, in incultis, 1943, fr., leg. F. Baglione.

7035 SC, Lages, in incultis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

8721 RGS, Bom Jesus, NEH, ad viam, 15.1.1942, fr., det. Swallen.

34231 Chile, Gualpén p. Concepción, in pascuo, 1.1.1941, fr., leg. et. det. H. Gunckel.

32459 RGS, Serra da Rocinha p. Bom Jesus, NEH, ad viam, 28.2.1946, fr., det. Parodi.

- 35158** RGS, Fazenda da Ronda p. Vacaria, NEH, in graminosis, 15.1.1947, fr.
- 35211** RGS, Serra da Rocinha p. Bom Jesus, NEH, ad viam in tritis, 14.2.1947, fl.
- 36474** RGS, Cambará p. S. Francisco de Paula, NEH, ad viam, 2.1948, fr., det. Swallen.
- 37076** Chile, Cautín p. Cajón, in prato, 26.3.1946, fl., leg. et det. H. Gunckel 5867.
- 50455** RGS, Serra da Rocinha p. Bom Jesus, NEH, ad viam, 18.1.1950, fr., leg. A. Sehnem SJ 4250.
- 53893** RGS, ibidem, 3.2.1953, fr.
- 54701** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, ad viam, 1.1.1954, fr.
- 54332** RGS, Fachinal Preto p. Bom Jesus, NEH, ad viam, 29.1.1950, fl., leg. R. Reitz 5000, det. Swallen.
- 59586** RGS, Farroupilha, NEH, in incultis, 26.9.1956, fl., leg. O. Camargo 784.
- 62076** RGS, Bento Gonçalves, NEH, in incultis, 6.10.1957, fl., leg. O. Camargo, 1971.
- 62249** RGS, Garibaldi, NEH, in incultis, 13.10.1957, fl., leg. O. Camargo 2033.

Habit: Up to 50 cm high.

Habitat: Disturbed soil; roadside.

Distribution in RGS: Fairly common, especially on the northeastern Highlands.

General distribution: Native to Europe; RGS, Uruguay, B. Aires.

HORDEUM L.

Hordeum compressum Gris., Plant. Lor. (1974) 201; Döll, FB 2,3 (1880) 233; Cabrera, Fl. Alred. B. Aires (1953) 67.

1946 RGS, S. Leopoldo, NEL, in campestribus siccis graminosis, 8.9.1934, fl., leg. C. Orth SJ, det. Chase.

Habit: Up to 50 cm high.

Habitat: Sandy, dry Campos.

Distribution in RGS: No other findplace.

General distribution: South Brazil, Central Argentina.

Hordeum pusillum Nutt., Gen. Am. 1 (1818) 87; Cabrera, Fl. Alred. B. Aires (1953) 67.

53064 RGS, Cêrro Largo p. S. Luiz, NEW, in incultis, 10.1952, fr., det. Swallen.

Habit: 10-40 cm high.

Habitat: Disturbed soil.

Distribution in RGS: No other findplace.

General distribution: Native to North America; Argentina, RGS.

HYMENACHNE Beauv.

Hymenachne amplexicaulis (Rudge) Nees, Agrost. Bras. (1829) 276; Cabrera, Fl. Alred. B. Aires (1953) 105. — *Panicum amplexicaule* Rudge, Plant. Guiana 1 (1805) 21; Döll, FB 2,3 (1877) 234; Herter, Flor. Urug. (1930) 29.

7450 RGS, S. Leopoldo, NEL, in paludosis alte dumosis, 1907, fr., leg. F. Theissen SJ.

31456 S. Leopoldo, NEL, Sombrio p. Araranguá, ad lacum in alte dumosis, 1.2.1946, fr., det. Swallen.

47055 RGS, Lagoa da Pinguela p. Osório, NEC, in humidis alte dumosis, 8.5.1950, fr., det. Beetle.

49700 RGS, Lagoa dos Quadros p. Torres, NEC, in paludosis alte dumosis, 18.11.1951, fr., det. Beetle.

Habit: Up to 1,5 m high.

Habitat: Swampy thickets.

Distribution in RGS: Northeastern sea coast and Lowlands.

General distribution: From the Antilles to RGS.

HYPARRHENIA

Hyparrhenia hirta (L.) Stapf,

Andropogon hirtus L., Spec. Plant. (1753) 1046.

1106 RGS, Dois Irmãos p. S. Leopoldo, NES, in secundariis alte graminosis, 13.9.1934, fr., leg. C. Orth SJ, det. Chase, Parodi.

Habit: Bunches up to 1 m high.

Habitat: Dry secondary growth.

Distribution in RGS: No other findplace.

General distribution:

ICHNANTHUS Beauv.

Ichnanthus pallens (Swartz) Munro, ex Benth. Flor. Hongk. (1861) 414; Döll, FB 2,2 (1877) 290; Ekman, Ark. f. Bot. 11,n.4 (1912) 26; idem, Ark. f. Bot. 13,n.10 (1913) 32; Malme, Sv. Bot. Tidskr. 30 (1936) 10; Cabrera, Fl. Alred. B. Aires (1953) 105.

- 1082** RGS, S. Leopoldo, NEL, in silva campestri, 28.8.1934, fr., leg. C. Orth SJ, det. Chase.
- 35568** RGS, Kappesberg p. Montenegro, NES, in silva, 15.12.1946, fr., leg. E. Henz SJ.
- 36466** RGS, Cambará p. S. Francisco de Paula, NEH, in araucarieto, 2. 1948, fr.
- 38763** RGS, Conceição p. Caí, NEL, in silva campestri ad rivum, 12.14.1948, fr., det. Swallen.
- 39865** RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in silva, 12.1.1949, fr.
- 40476** RGS, Vila Manresa P. Alegre, NEL, ad rivum in silva, 14.3.1949, fr., det. Swallen.
- 40899** RGS, ad montem das Cabras p. S. Leopoldo, NEL ad rivum in silva, 8.4.1949, fr., det. Swallen.
- 41686** RGS, Schwabenschneis p. S. Leopoldo, NES, in silva, 25.5.1949, fr.
- 41783** RGS, Esteio p. S. Leopoldo, NEL, in silva campestri, 1.6.1949, fr., det. Swallen.
- 41878** RGS, Reutersberg p. Novo Hamburgo, NES, in silva, 6.6.1949, fr.
- 42070** RGS, ad montem Sapucaia p. S. Leopoldo, NEL, ad rivum in silva, 18.6.1949, fr., det. Swallen.
- 42158** RGS, Schwabenschneis p. Novo Hamburgo, in silva, 25.6.1949, fr.
- 45872** RGS, Lagoa dos Quadros p. Torres, NEC, in umbrosis, 21.2.1950, fr.
- 45902** RGS, ibidem, 21.2.1955, fr.
- 46240** RGS, S. Francisco de Paula, NEH, in araucarieto, 13.3.1950, fr.
- 47096** RGS, Vila Manresa p. P. Alegre, NEL, ad rivum in umbrosis, 15.5.1950, fr., det. Parodi.
- 49467** RGS, Mato Leitão p. Venâncio Aires, NES, in silva ad viam, 1.1.1951, fl.
- 57140** RGS, Vila Manresa p. P. Alegre, NEL, in silvula, 4.11.1955, fl.,
- 60443** RGS, Estação de Silvicultura p. Sta Maria, CL, in silva campestri, 10.4.1957, fl., leg. O. Camargo 515.
- 60610** RGS, Farroupilha, NEH, in silva, 7.5.1957, fr., fr., leg. O. Camargo 1360.
- 61977** RGS, Taquarí, NEL, in silva, 12.12.1957, fl., leg. O. Camargo 2898.

Habit: Prostrate to ascending, branches up to 50 cm long.

Habitat: Forest.

Distribution in RGS: Through the whole of the northern forested zone

General distribution: From the Antilles to RGS and B. Aires.

IMPERATA Cyrillo

Imperata brasiliensis Trin., Mém. Ac. St. Petersb. ser. 6, 2 (1832) 331; Hackel, FB 2,3 (1883) 251; Ekman, Ark. f. Bot. 13,n.10 (1913) 4; Malme, Sv. Bot. Tidskr. 30 (1936) 9; Cabrera, Fl. Alred. B. Aires (1953) 110; Reitz, Sellowia 7 (1956) 102.

52746 RGS, Torres, NEC, prope litus in arenosis, 7.8.1952, fl., leg. et det. Beetle.

Habit: Erect, up to 40 cm high.

Habitat: More or less dry dunes.

Distribution in RGS: Only findplace.

General distribution: From Mexico to RGS and B. Aires.

Imperata contracta (H.B.K.) Hitchc.,

I.caudata Trin., Act. Ac. St. Petersb. 8,6 (1833) 331, tab. 2; Hackel, FB 2,3 (1883) 231;

25457 SC, Antônio Carlos p. Biguassú, in incultis, 2.3.1943, fr., leg. R. Reitz, det. Swallen.

46488 RGS, Lagoa da Pinguela p. Osório, NEC, in paludosis alte dumosis, 27.3.1950, fr., det. Parodi.

Habit: Up to 2 m high.

Habitat: Wet places among shrubs.

Distribution in RGS: Northeastern sea coast only.

General distribution: From Mexico to RGS.

Imperata tenuis Hack.,

2708 RGS, Portão p. S. Leopoldo, NEL, in paludosis alte graminosis, 27.11.1935, fr., leg. C. Orth SJ, det. Chase.

Habit: Up to 1,5 high.

Habitat: Swampy thickets among high grasses.

Distribution in RGS: Northeastern Lowlands.

General distribution:

ISCHAEMUM L.

Ischaemum urvilleanum Kunth, Rev. Gram. 1 (1829) 157; Hackel, FB 2,3 (1883) 260, tab. 72, 1; Ekman, Ark. f. Bot. 13,n.10(1913) 6; Herter, Flor. Urug. (1930) 27; Cabrera, Fl. Alred. B. Aires (1953) 110; Reitz, Sellowia 7 (1956) 102.

732 RGS, S. Leopoldo, NEL, in graminosis humidis, 10.1.1933, fr., leg. C. Orth SJ, det. Chase.

- 37981** RGS, Barra do Ribeiro p. P. Alegre, NEL, in humidis graminosis, 14.11.1948, fr., det. Swallen.
- 38272** RGS, Esteio p. P. Alegre, NEL, in subpaludosis graminosis, 24.11.1948, fl., det. Swallen.
- 44308** RGS, ibidem, 8.11.1949, fr.
- 44487** RGS, Granja Neugebauer p. Itapoan, NEL, in subpaludosis graminosis, 19.11.1949, fl.
- 51146** Paraná, Matinho p. Paranaguá, in humidis graminosis, 2.11.1949, fl., leg. G. Hatschbach 1579, det. Parodi.
- 54345** SC, Laguna, in arenosis prope litus, 20.2.1952, fr., leg. R. Reitz 5049, det. Swallen.
- 55377** SC, Massiambú p. Palhoça, in arenosis, 18.12.1952, fr., leg. R. Reitz 7148, det. Swallen.
- 56220** RGS, Torres, NEC, prope litus in dense graminosis, 12.11.1954, fr.
- 57144** RGS, Vila Manresa p. P. Alegre, NEL, in humidis dense graminosis, 4.11.1955, fl.
- 58034** SC, Massiambú p. Palhoça, in arenosis humidis, 5.11.1953, fr., leg. Reitz 8282, det. Swallen.
- 58049** SC, ibidem, 24.9.1953, fr., leg. R. Reitz 8323, det. Swallen.
- 62714** RGS, Porto Alegre, NEL, in humidis dense graminosis, 2.11.1957, fr., leg. O. Camargo 2368.

Habit: Densely ces pitose.

Habitat: More or less wet meadows.

Distribution in RGS: Northeastern Lowlands and sea coast.

General distribution: From S. Paulo to Uruguay and B. Aires.

KOELERIA Pers.

Koeleria phleoides (Vill.) Pers., Syn. Pl. 1 (1805) 97; Herter, Flor. Urug. (1930) 37; Cabrera, Flor. Alred. B. Aires (1953) 70. — *Festuca phleoides* Vill., Flor. Delph. (1895) 7.

- 875** RGS, Vila Manresa p. P. Alegre, NEL, in agro, 5.10.1933, fr., leg. C. Orth SJ, det. Chase.
- 26978** RGS, ibidem, 10.1944, fr., det. Swallen.
- 38073** RGS, ibidem, 17.11.1948, fl., det. Swallen.
- 56009** RGS, ibidem, 19.11.1954, fl..
- 57191** RGS, P. Alegre, NEL, in via publica, 1955, fl.

Habit: Dense bunches, up to 40 cm high.

Habitat: Disturbed soil.

Distribution in RGS: Widely scattered: Pelotas, S. Leopoldo, P. Alegre.

General distribution: Native to Europe; RGS, Uruguay, B. Aires.

LASIACIS (Gris.) Hitchc.**Lasiacis divaricata (L.) Hitchc.**

Panicum divaricata L., Spec. Plant. ed. 2 (1762) 86. — *P. latifolium* L., Spec. Plant. (1753) 58; Döll, FB 2,2 (1877) 206; Ekman, Ark. f. Bot. 11, n.4 (1912) 18; idem, Ark. f. Bot. 13, n.10 (1913) 25; Reitz, Sellowia 7 (1956) 102.

2719 RGS, Kappesberg p. Montenegro, NES, in silva pluviali, 4.1.1936, fr., det. Chase.

3817 RGS, Bohnental p. Caí, in silva, 4.1.1941, ster.

7442 RGS, S. Leopoldo, NEL, ad silvam, fr., leg. F. Theissen SJ.

28348 RGS, Nonoai, NWU, in silva pluviali, 3.1945, fr.

39890 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in silva, 12.1.1949, fr.

40005 RGS, Linha Bonita p. Montenegro, NES, in silva, 19.1.1949, fr.

42192 RGS, Schwabenschneis p. Novo Hamburgo, NES, in silva pluviali, 25.6.1949, fr.

42200 RGS, ad montem Steinkopf p. S. Leopoldo, NEL, in silva pluviali, 28.6.1949, fr. deciso.

42419 RGS, Pareci p. Montenegro, NEL, in silva pluviali, 7.7.1949, fr.

42693 RGS, Vila Manresa p. P. Alegre, NEL, in silva pluviali, 25.7.1949, fr.

46452 RGS, Lagoa da Pinguela p. Osório, NEC, in silva secundaria, 27.3.1950, fr.

47132 RGS, Butterberg p. Montenegro, NES, in silva pluviali, 22.5.1950, fr.

49870 SC, Itapiranga ad fl. Uruguay superius, in silva pluviali, 6.2.1951, fr.

53628 Paraná, Salto Iguassú, in silva, 11.1.1953, fr. maturo.

58038 SC, Morro da Fazenda p. Itajaí, in silva, 4.3.1954, fl., leg. R. Reitz 8292, det. Swallen.

60808 RGS, Sta Rita p. Farroupilha, NEH, in silva, 18.5.1957, fl., leg. O. Camargo 1448.

Habit: Recumbent, up to 3 m high.

Habitat: Rain forest.

Distribution in RGS: Common through the whole of the northern rain forest belt.

General distribution: From Guyana to Misiones and RGS.

LEERSIA Swartz

Leersia hexandra Swartz, Prodr. (1788) 21; Ekman, Ark. f. Bot. 11,n.4 (1912) 36; Herter, Flor. Urug. (1930) 32; Cabrera, Flor. Alred. B. Aires

(1953) 91; Reitz, Sellowia 7 (1956) 102. — *Oryza hexandra* (Sw.) Döll, FB 2,2 (1871) 10.

9849 RGS, Tupanciretan, NWH, in paludosis, 29.1.1942, fr., det. Parodi.

33192 RGS, Vila Oliva p. Caxias, NEH, in paludosis, 16.1.1946, fr.

35141 RGS, Fazenda da Ronda p. Vacaria, NEH, in humidis, 15.1.1947, fr.

35666 RGS, S. Leopoldo, NEL, in paludosis, 20.12.1946, fr., leg. E. Henz SJ.

41495 RGS, Estação Azevedo p. Montenegro, NEL, in palude, 6.5.1949, fr.

44774 RGS, Glorinha p. Gravataí, NEL, in palude, 14.12.1949, fr.

45142 RGS, Fazenda do Arroio p. Osório, NEC, in palude, 4.1.1950, fr.

55066 RGS, Vila Oliva p. Caxias, NEH, in subpaludosis, 24.2.1954, fr., det. Swallen.

55071 RGS, ibidem, 24.2.1954, fr.

58829 RGS, Estação de Silvicultura p. Sta Maria, CL, in subpaludosis, 1.3.1956, fr., leg. O. Camargo 142.

58981 RGS, ibidem, 5.1.1956, fr., leg. O. Camargo 157.

Habit: 30 cm high.

Habitat: Wet to swampy places.

Distribution in RGS: Over the whole of the Campos area, rather inconspicuous.

General distribution: From Guyana to RGS, Uruguay and B. Aires.

LEPTOCORYPHIUM Nees

Laptocoryphium lanatum (H.B.K.) Nees, Agrost. Brasil. (1829) 84; Döll, FB 2,2 (1877) 120, tab. 17; Herter, Flor. Urug. (1930) 29; Cabrera, Flor. Alred. B. Aires (1953) 104; Reitz, Sellowia 7 (1956) 104. — *Paspalum lanatum* H.B.K., Nov. Gen. et Spec. 1 (1816) 94, tab. 29. — *Athaenantia lanata* (H.B.K.) Benth.; Ekman, Ark.f.Bot. 11,n.4 (1912) 17; idem, Ark.f.Bot. 13, n.10 (1913) 22.

867 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 15.10.1933, fr., leg. C. Orth SJ, det. Chase.

29045 RGS, ibidem, 25.18.1945, fr.

33921 RGS, ibidem, 18.9.1946, fl.

34128 RGS, ibidem, 9.10.1946, fr.

37391 RGS, ibidem, 5.7.1948, fr.

37392 RGS, ibidem, 5.7.1948, fr.

38663 RGS, Rio dos Sinos p. S. Leopoldo, NEL, in campestribus

siccis, 10.12.1948, fl., det. Swallen.

38684 RGS, ibidem, 10.12.1948, fr.

44094 RGS, Barreto Vianna p. S. Leopoldo, NEL, in siccis, 24.10.1949, fr.

60337 RGS, Estação de Silvicultura p. Sta Maria, CN, in campestribus alte graminosis, 8.4.1956, fr., leg. O. Camargo 557.

61984 RGS, Taquarí, NEL, in campestribus siccis alte graminosis, 10.12.1957, fl., leg. Camargo 2882.

Habit: Stout bunches, up to 60 cm high.

Habitat: Dry Campos among high grasses and shrubs.

Distribution in RGS: Central and northeastern Lowlands.

General distribution: From Guyana to Uruguay and B. Aires.

LUZIOLA Juss.

Luziola peruviana Gmel., Syst. Nat. 2 (1791) 637; Döll, FB 2,2 (1871) 17, tab. 5; Herter, Flor. Urug. (1930) 32; Cabrera, Fl. Alred. B. Aires (1953) 91; — **Luziola leiocarpa** Lindm., Kongl. Sv. Vet. Ak. Handl. 34, n.6 (1912) 29; Ekman, Ark. f. Bot. 11,n.4(1912) 29. — Reitz Sellowia 7 (1956) 106.

1051 RGS, S. Leopoldo, NEL, in palude, 20.9.1934, fr., leg. C. Orth SJ, det. Chase.

31458 SC, Sombrio p. Araranguá, in palude, 1.2.1946, fr.

40135 RGS, Guaíba, NEL, in paludosis, 24.1.1949, fl.

40325 RGS, Sta Maria p. Farroupilha, NEH, in paludosis, 29.1.1949, fl., (Swallen: L. leiocarpa Lindm.?).

38519 RGS, S. Leopoldo, NEL, in paludosis, 4.12.1948, fr.

38802 RGS, Conceição p. Caí, in stagno, 14.12.1948, fl. (Swallen: L. leiocarpa Lindman?).

39602 RGS, Cachoeirinha p. Gravataí, NEL, in stagno, 7.1.1949, fl.

44316 Esteio p. S. Leopoldo, NEL, in stagno, 8.11.1949, fl.

44461 RGS, Granja Neugebauer p. Itapoan, NEL, in stagno, 19.11.1949, fr.

45705 RGS, Farroupilha, NEH, in stagno, 7.2.1950, fl.

54926 RGS, Vila Oliva p. Caxias, NEH, in stagno, 24.2.1954, fl. (Swallen: L. leiocarpa Lindm.).

57137 RGS, Vila Manresa p. P. Alegre, NEL, in stagno, 4.11.1955, fl.

58057 SC, Massiambú p. Palhoça, in paludosis, 5.11.1953, fr., leg. R. Reitz 8349, (Swallen: L. leiocarpa Lindm.).

59402 RGS, Farroupilha, NEH, in stagno fuitans, 15.11.1956, fr., fl., leg. O. Camrgo 984.

61490 RGS, Taquarí, NEL, in stagno fuitans, 14.12.1957, fl., leg. O. Camargo 2912.

Habit: Submersed, leaves floating.

Habitat: Shallow, stagnant or more or less perennial ponds.

Distribution in RGS: Common all over the Campos area.

General distribution: From Mexico to Uruguay and B. Aires.

MANISURIS Swartz

Manisuris loricata (Trin.) Kuntze, *Rottboellia loricata* Trin., Mém. Ac. Petersb. ser. 6, 5 (1849) 250; Hackel, FB 2, 3 (1883) 311, tab. 21, fig. 1, 2; Ekman, Ark. f. Bot. 13, n. 10 (1913) 6.

1074 RGS, S. Leopoldo, NEL, in humidis alte dumosis, 4.5.1933, fr., det. Chase.

2710 RGS, Portão p. S. Leopoldo, NEL, in subpaludosis alte dumosis, 27.11.1935, fr., leg. C. Orth SJ, det. Chase.

31944 SC, Sombrio p. Araranguá, in paludosis alte dumosis, 5.12.1943, fr., leg. R. Reitz 1315, det. Swallen.

60438 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fr., leg. O. Camargo 590.

61664 RGS, Taquarí, NEL, in humidis alte graminosis, 10.12.1957, fl., leg. O. Camargo 2841.

Habit: Up to 1,5 m high.

Habitat: Densely vegetated swamps.

Distribution in RGS: Swamps of the northern half of the State.

General distribution: From Minas Gerais to RGS.

MELICA L.

Melica aurantiaca Lam., Encycl. Méth. 4 (1797) 70; Döll, FB 2, 3 (1878) 128; Ekman, Ark. f. Bot. 11 n. 4 (1912) 44; Herter, Flor. Urug. (1930) 44; Cabrera, Flor. Alred. B. Aires, (1953) 63.

1098 RGS, Sapucaia p. S. Leopoldo, NEL, ad viam in siccis dumosis, 17.10.1934, fr., leg. C. Orth SJ (Parodi: *M. papilionacea* L.).

2713 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis graminosis, 19.11.1935, fr., leg. C. Orth SJ, det. Chase, Parodi Swallen.

8285 RGS, Montserrat p. P. Alegre, NEL, in campestribus siccis, 13.10.1941, fr., leg. K. Emrich.

11772 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 1943, fl.

11764 RGS, ibidem, 1943, fr.

27120 RGS, ibidem, 1943, fr.

- 29204** RGS, *ibidem*, 3.10.1945, fr.
34137 RGS, *ibidem*, 9.10.1946, fr.
34198 RGS, *ibidem*, 16.10.1946, fl.
35463 RGS, Cambará p. S. Francisco de Paula, NEH, in *campestribus siccis*, 2.1948, fr.
37745 RGS, Vila Manresa p. P. Alegre, NEL, in *campestribus siccis*, 22.9.1948, fl.
43728 RGS, Morro de Sta Teresa p. P. Alegre, NEL, in *campestribus siccis*, 3.10.1949, fl.
43976 RGS, Vila Manresa p. P. Alegre, NEL, in *campestribus siccis*, 19.10.1949, fr., det. Swallen.
43998 RGS, Morro do Osso p. P. Alegre, NEL, in *campestribus siccis*, 21.10.1949, fr.
48987 RGS, Vila Manresa p. P. Alegre, in *campestribus siccis*, 18.10.1950, fr.
54468 RGS, *ibidem*, 12.11.1953, fr., det. Swallen.
55999 RGS, *ibidem*, 19.11.1954, fr.
57231 RGS, *ibidem*, 15.10.1955, fl.
57487 RGS, Estação de Silvicultura p. Sta Maria, CL, in *campestribus*, 21.11.1955, fl., leg. O. Camargo 40.

Habit: Up to 1 m high.

Habitat: Dry, grassy, Campos.

Distribution in RGS: Apparently over the whole of the Campos area.

General distribution: RGS, Uruguay, B. Aires.

Melica sarmentosa Nees, Agrost. Bras. (1829) 485; Döll, FB 2,3 (1878) 127; Ekman, Ark. f. Bot. 11,n.4 (1912) 44; Idem, Ark. f. Bot. 13,n.10 (1913) 52; Herter, Flor. Urug. (1930) 37; Cabrera, Fl. Alred. B. Aires (1953) 62.

2718 RGS, Kappesberg p. Montenegro, NES, in *dumosis scandens*, 4.1.1936, leg. C. Orth SJ, det. Chase.

6540 RGS, sine die, fr.

7021 SC, Lages, ad *silvulam scandens*, 1935, ster., Leg. A. Bruxel SJ.

26779 RGS, Cerro Largo p. S. Luiz, NWH, ad *silvam scandens*, 9.1944, fr., leg. E. Friderichs SJ.

29566 RGS, Pareci p. Montenegro, NEL, ad *silvam scandens*, 3.10.1945, fr., leg. E. Henz SJ, det. Swallen.

32603 RGS, *ibidem*, 22.10.1945, fr. deciso, leg. E. Henz SJ.

34344 RGS, Cérro Largo p. S. Luiz, NWH, ad *silvam scandens*, 31.10.1946, fr., leg. Irm. Augusto, det. Swallen.

35355 RGS, Dois Irmãos p. S. Leopoldo, NES, ad *silvam*, 10.10.1946, fr., leg. C. Ritter SJ.

- 38622** RGS, Morro da Polícia p. P. Alegre, NEL, ad silvulam scandens, 20.11.1948, fr.
- 39949** RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, ad silvam scandens, 12.1.1949, fr.
- 43423** RGS, Kappesberg p. Montenegro, NES, ad silvam pluvialem scandens, 11.9.1949, fl., det. Swallen.
- 43757** RGS, Morro do Sabiá p. P. Alegre, NEL, in ripariis scandens, 5.10.1949, fr.
- 43783** RGS, Parecí p. Montenegro, NEL, ad silvam scandens, 7.10.1949, fr.
- 44808** RGS, S. Francisco de Paula, NEH, ad araucarietum scandens, 18.12.1949, fr.
- 49109** RGS, Butterberg p. Montenegro, NES, ad silvam primaevam scandens, 13.11.1950, fr.
- 50865** Paraná, Morro Grande p. Cerro Azul, fr., leg. G. Hatschbach 1490, det. Parodi.
- 59587** RGS, Farroupilha, NEH, ad araucarietum scandens, 26.9.1956, fl., leg. O. Camargo 783,
- 61171** SC, Itapiranga ad fl. Uruguay, ad silvam scandens, 7.10.1957, fl.
- 61886** RGS, Garibaldi, NEH, in silva, 29.10.1957, fl., leg. O. Camargo 2272.
- 62147** RGS, Farroupilha, NEH, in dumosis scandens, 8.10.1957, fl., leg. O. Camargo 1991.
- 62154** RGS, Bento Gonçalves, NEH, ad silvam scandens, 6.10.1957, fl., leg. O. Camargo 1980.
- 62201** RGS, Garibaldi, NEH, ad silvam scandens, 13.10.1957, fl., leg. O. Camargo 2058.
- 62260** RGS, Farroupilha, NEH, in silva, 10.10.1957, fl., leg. O. Camargo 1903.
- 62334** RGS, Montenegro, NEL, in dumosis scandens, 19.9.1957, fl., leg. O. Camargo 1781.
- 62622** RGS, P. Alegre, NEL, ad silvulam campestrem scandens, 2.11.1957, fl., leg. O. Camargo 2424.

Habit: Climbing up to 5 m high.

Habitat: Edge of rain forest, secondary growth.

Distribution in RGS: Over the whole of the forested area.

General distribution: From Minas Gerais to Uruguay and B. Aires.

MELINIS Beauv.

Melinis minutiflora Beauv., Essay Agrost. (1812) 54, tab. 11, fig. 4; Ekman, Ark. f. Bot. 13,n.10 (1913) 10; Reitz, Sellowia 7 (1956) 102. —

Panicum melinis Trin., Acta Petrop. 1835, 291; Döll, FB 2,2 (1877) 241, tab. 33.

881 RGS, Vila Manresa p. P. Alegre, NEL, in dumosis subspontanea, 10.10.1933, fr., leg. C. Orth SJ, det. Chase.

4204 SC, Nova Trento p. Florianópolis, in dumosis secundariis, 6.1938, fr., det. Swallen.

10832 RGS, Marcelino Ramos, NWH, ad viam in dumosis, 1.1943, fr., leg. E. Friderichs SJ.

25466 RGS, S. Leopoldo, NEL, in dumosis, 1943, ster., leg. R. Reitz.

32063 SC, Sombrio p. Araranguá, ad viam in dumosis, 28.8.1945, fr., leg. R. Reitz 1130, fr.

42026 RGS, Morro Santana p. P. Alegre, NEL, in secundariis efferata, 15.6.1949, fr.

43557 RGS, Portão p. S. Leopoldo, NEL, in secundariis eferata, 23.9.1949, fr.

52748 RGS, Escola Agronômica p. P. Alegre, in dumosis, 11.7.1952, fr., leg. et det. Beetle.

55379 SC, Ascurra p. Rodeio, in dumosis, 11.6.1953, leg. R. Reitz 7142, det. Swallen.

55395 SC, Praia Braba p. Itajaí, in dumosis scandens, 28.5.1953, fr., leg. R. Reitz 7115, det. Swallen.

Habit: Prostrate to ascending or half-climbing, up to 2 m high.

Habitat: Cultivated, or subspontaneous in secondary growth

Distribution in RGS: Over the whole of the northern cultivated area.

General distribution: Tropical and subtropical America.

MEROSTACHYS Spreng.

Merostachys burchellii Munro, Monogr. () 51; Döll, FB 2,3 (1880) 218.

3239 RGS, Kappesberg p. Montenegro, NES, in silva subhumida, 1938, fr.

4880 RGS, Passo do Inferno p. Canela, NEH, ad flumen in dumosis, 10.2.1941, fr.

6431 RGS, sine loco, sine die, fr.

9021 RGS, Bom Jesus, NEH, in silva campestri ad rivum, 16.1.1942, fr.

47382 RGS, Campinas p. Sta Rosa, NWH, in silva pluviali, 2.1950, fr., leg. A. Spies.

56384 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in araucarieto, 2.1.1955, ster. (19)

56611 RGS, Vila Oliva p. Caxias, NEH, in araucarieto, 8.2.1955, ster. (19)

60611 RGS, Farroupilha, NEH, in araucarieto, 7.5.1957, ster., leg. O. Camargo 1340.

60880 RGS, ibidem, 8.7.1957, ster., leg. O. Camargo 1648. (19)

60984 RGS, ibidem, 8.5.1957, ster., leg. O. Camargo 1434.

Habit: Culms up to 8 m high.

Habitat: More or less wet forest, often dominant in the undergrowth.

Distribution in RGS: Northeastern Highlands.

General distribution: From S. Paulo to RGS.

MICROCHLOA Brown

Microchloa indica (L.) K.,

Herter, Flor. Urug. (1930) 39.

29040 RGS, Vila Manresa p. P. Alegre, NEL, in rupestribus humidis parce humosis, 25.8.1945, fr.

54964 RGS, Vila Oliva p. Caxias, NEH, in rupestribus humidis parce humosis, 24.2.1954, fr. (Swallen: Microchloa sp.).

MUEHLENBERGIA Schreb.

Muehlenbergia schreberi Gmel., Syst. Nat. 2 (1791) 171; Herter, Flor. Urug. (1930) 33; Cabrera, Flor. Alred. B. Aires (1953) 76, fig. 23 J. — **M. diffusa** Schreb., Gräser () tab. 51; Döll, FB 2,3 (1878) 40; Ekman, Ark. f. Bot. 11,n.4 (1912) 33; Malme, Sv. Bot. Tidskr. 30 (1936) 10.

2874 RGS, Schwabenschneis p. Novo Hamburgo, NES, in secundariis, 14.9.1937, fr., leg. C. Orth SJ, det Parodi: **M. diffusa** Schreb.

42314 RGS, Langschneis p. Montenegro, NES, ad viam ferream in dumosis, 2.7.1949, fr., det. Swallen.

55090 RGS, Vila Oliva p. Caxias, NEH, ad viam in dumosis, 24.2.1954, fr., det. Swallen.

59775 RGS, Farroupilha, NEH, in dumosis secundariis, 18.3.1957, fl., leg. O. Camargo 1168.

59778 RGS, ibidem, 25.3.1957, fl., leg. O. Camargo 1196.

64344 SC, S. Joaquim, in campestribus, 10.2.1958, fl., leg. J. Mattos.

64345 SC, ibidem, 1.2.1958, fl., leg. J. Mattos.

Habit: Prostrate to ascending, branches up to 1 m long.

Habitat: Disturbed soil, secondary growth.

Distribution in RGS: Northeastern Lowlands, rainforest belt and Highlands.

General distribution: Subtropical and temperate America; Misiones, RGS, Uruguay, B. Aires.

OLYRA L.

Olyra humilis NEES, Agrost. Bras. (1829) 403; Döll 2,2 (1877) 321; Reitz, Sellowia 7 (1956) 103.

840 RGS, Vila Manresa p. P. Alegre, NEL, in silva primaeva, 10.11.1931, fr., det. Chase, leg. C. Orth SJ.

37864 RGS, Morro da Polícia p. P. Alegre, NEL, in silva campestri, 11.11.1948, fr., det. Swallen.

43033 RGS, Schwabenschneis p. Novo Hamburgo, NES, in silva pluviali, 22.8.1949, ster.

43745 RGS, Morro do Sabiá p. P. Alegre, NEL, in silvula riparia, 5.10.1949, fr.

43790 RGS, Parecí p. Montenegro, NEL, in silva pluviali, 7.10.1949, fl.

44445 RGS, Granja Neugebauer p. Itapoan, NEL, in silva, 19.11.1949, fr.

48937 RGS, ibidem, 11.10.1950, fl.

50877 Paraná, Piraquara p. Florestal, in araucarieto, 19.12.1948, fr., leg. G. Hatschbach 1140, det. Parodi.

53077 RGS, Cêrro Largo p. S. Luiz, NWH, in silva, 20.11.1942, fr.

53713 SC, Itapiranga ad fl. Uruguay superius, in silva pluviali, 17.1.1953, fr.

61507 RGS, Farroupilha, NEH, in silva, 25.11.1957, fl., leg. O. Camargo 2446, fl.,

61833 RGS, Garibaldi, NEH, in silva, 29.10.1957, fl., leg. O. Camargo 2242.

Habit: Bambusoid, loose bunches, up to 1 m high.

Habitat: Interior of the rain forest.

Distribution in RGS: Northern rain forest area.

General distribution: From Goiás to RGS.

Olyra latifolia L., Amoen. Ac. 5 () 408; Döll, FB 2,2 (1877) 316; Reitz, Sellowia 7 (1956) 103.

6430 SC, Tubarão, S. Ludgero p., in silva pluviali, 4.1.1941, fr., leg. C. Orth SJ, det. Swallen.

3602 Rio de Janeiro, ad montem Corcovado, 5.2.1940, fr., det. Swallen.

31839 SC, Sombrio p. Araranguá, in silva pluviali, 9.2.1946, fr., det. Swallen.

42283 RGS, ad montem Steinkopf p. S. Leopoldo, NEL, in silva pluviali, 28.6.1949, fr., det. Swallen.

Habit: Bambusoid, loose bunches, culms up to 1,5 m high.

Habitat: Interior of rain forest.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Amazonia to RGS.

OPLISMENUS Beauv.

Oplismenus hirtellus (L.) Beauv.,

Panicum hirtellaum L., Spec. Plant. ed. 2 (1762) 83. — Panicum loliaceum Lam., III. 1 () 170; Döll, FB 2,2 (1877) 145. — Oplismenus hirtellus (L.) R. et S., Ekman, Ark. f. Bot. 13, n.10 (1913) 33.

26365 RGS, Fazenda do Jaraú p. Quaraí, Sv, in silva campestri, 1.1945, fr.

30518 SC, Sombrio p. Araranguá, in umbrosis, 12.4.1944, fr., det. Swallen.

33063 RGS, Parecí p. Montenegro, NEL, in silva, 15.1.1945, fr., leg. C. Ritter SJ, det. Swallen.

41351 RGS, Morretes p. P. Alegre, NEL, in silva campestri, 2.5.1949, fr., det. Swallen.

42184 RGS, Schwabenschneis p. Novo Hamburgo, NES, in silva, 25.6.1949, fr. deciso.

63692 RGS, Feliz p. Caí, ad flumen in umbrosis, 17.3.1958, fl., fr., leg. O. Camargo.

Habit: Prostrate to ascending, rooting at the lower joints, branches up to 40 cm long.

Habitat: Forest and woodlets.

Distribution in RGS: Northeastern Lowlands and rain forest belt.

General distribution: From the Antilles to RGS.

Oplismenus setarius (Lam.) R. et Schult., Syst. Veg. 2 (1817) 481; Ekman, Ark. f. Bot. 13, n.10 (1913) 33; Herter, Flor. Urug. (1930) 29. — Panicum setarium Lam., Illustr. 1 (1791) 170; Döll, FB 2,2 (1877) 147.

731 RGS, Vila Manresa p. P. Alegre, in silva umbrosa, 15.1.1933, fl, fr., leg. C. Orth SJ, det. Chase.

40598 RGS, Esteio p. S. Leopoldo, NEL, in silva campestri, 23.3.1949, fr., det. Swallen.

41578 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in silva, 16.5.1949, fr., det. Swallen.

41842 RGS, sine loco, 3.6.1949, fl.

42072 RGS, Sapucaia p. S. Leopoldo, NEL, in silva campestri, 18.6.1949, fr.

45997 RGS, Lagoa dos Quadros p. Torres, NEC, in silva umbrosa, 21.2.1950, fr.

55354 SC, Rio Tavares in Insula Florianópolis, in arenosis umbrosis, 13.5.1953, leg. R. Reitz, 7173, det. Swallen.

55424 SC, Massimbú p. Palhoça, in umbrosis arenosis, 12.3.1953, fr., leg. R. Reitz 7067, det. Swallen.

60404 RGS, Feliz p. Caí, in umbrosis, 26.4.1956, fr., leg. O. Camargo 622.

Habit: Prostrate to ascending, branches up to 40 cm long.

Habitat: Shady places, forest.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Texas and Mexico to Uruguay and B. Aires.

ORYZA L.

Oryza subulata Nees, Agrost. Bras. (1829) 518; Döll, FB 2,2 (1871) 8; Ekman, Ark. f. Bot. 11,n.4 (1912) 29; Herter, Flor. Urug. (1930) 32.

41082 RGS, ad fl. Gravataí inferius p. P. Alegre, in subpaludosis alte dumosis, 18.4.1949, fl., det. Swallen.

Habit: Thick grayish bunches up to 1,5 high.

Habitat: Swampy thickets

Distribution in RGS: Common in the southwestern plains, often in extense stands, used for thatching.

General distribution: From Paraguay to Western RGS, and Uruguay.

PANICUM L.

Panicum anomalum Trin.,

59234 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in humidis dumosis, 12.2.1956, fr.

Habit: Strong, erect, up to 5 m high.

Habitat: Wet to swampy thickets.

Distribution in RGS: Northeastern Highlands.

General distribution:

Panicum aquaticum Poir., Encycl. Supl. 4 () 281; Reitz, Sellowia 7 (1956) 103.

38590 RGS, ad montem das Cabras p. S. Leopoldo, NEL, in paludosis alte graminosis, 7.12.1948, fr., det. Swallen.

40542 RGS, Sapucaia p. S. Leopoldo, NEL, in subpaludosis alte graminosis, 17.3.1949, fr., det. Swallen.

40725 RGS, Cristal p. P. Alegre, NEL, in humidis alte graminosis, 31.3.1949, fr., det. Swallen.

41092 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in paludosis dense dumosis, 18.4.1949, fr., det. Swallen.

41798 RGS, Esteio, p. S. Leopoldo, NEL, in paludosis alte graminosis, 1.6.1949, fr.

41739 RGS, Fião p. S. Leopoldo, NEL, in subpaludosis alte graminosis, 25.9.1949, fr., det. Swallen.

55346 SC, Massiambú p. Palhoça, in palude, 19.12.1952, fr., leg. R. Reitz 7185, det. Swallen.

55398 SC, Rio Tavares in Insula Florianópolis, in paludosis, 11.3.1953, leg. R. Reitz 7112, det. Swallen.

58033 SC, ibidem, 5.11.1953, fr., leg. R. Reitz 8275, det. Swallen.

Habit: Culms vigorous, 1 m and more long.

Habitat: Swampy thickets.

Distribution in RGS: Northeastern Lowlands.

General distribution: From the Antilles to RGS.

Panicum aristellum Döll, FB 2,2 (1877) 221; Ekman, Ark. f. Bot. 13, n.10 (1913) 26.

9731 RGS, Tupanciretan, NWH, in campestribus dumosis, 29.1.1942, fr., det. Swallen.

Habit: Up to 1,5 m high.

Habitat: Shrubby Campos.

Distribution in RGS: Northeastern Highlands.

General distribution: From Minas Gerais to RGS.

Panicum bergii Arech., Gram. Urug. (1894) 127; Ekman, Ark. f. Bot. 11,n.4 (1912) 19; idem, Ark. f. Bot. 13,n.10 (1913) 25; Herter, Flor. Urug. (1930) 29; Cabrera, Flora. Alred. B. Aires (1953) 100.

9845 RGS, Tupanciretan, NWH, in campestribus siccis, 29.1.1942, fr., det. Swallen.

9998 RGS, Ijuisinho p. Tupanciretan, NWH, in campestribus siccis, 30.1.1942, fr., det. Swallen.

26670 RGS, Pareci p. Montenegro, NEL, in campestribus siccis, 1944, ster. leg. E. Henz SJ.

28337 RGS, Nonoai, NWU, in campestribus siccis, 3.1945, fr.

32956 Uruguay, Soriano, in campestribus siccis, 1.1944, fr., leg. et det. B. Rosengurtt.

44499 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus alte graminosis, 19.11.1949, fr., det. A. Beetle.

58607 RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus siccis, 10.1.1956, fr., leg. O. Camargo 26.

Habit: Vigorous bunches, up to 75 cm high.

Habitat: Dry, shrubby, Campos.

Distribution in RGS: Northwestern Highlands, northeastern Lowlands.

General distribution: From Paraná to RGS, Uruguay and B. Aires.

Panicum boliviense Hack..

Reitz, Sellowia 7 (1956) 104.

7455 RGS, S. Leopoldo, NEL, in campestribus dumosis, 1907, fr., leg. F. Theissen SJ, det. Swallen ("brasiliense").

38538 RGS, ibidem, 4.12.1948, fr.

38765 RGS, Conceição p. Caí, NEL, in campestribus dumosis, 14.12.1948, fr.,

41663 RGS, S. Leopoldo, NEL, in umbrosis dumosis, 21.5.1949, fr., det. Swallen.

41892 RGS, Reutersberg p. Novo Hamburgo, NES, in secundariis, 6.6.1949, fr.

41988 RGS, Morro Santana p. P. Alegre, NEL, in dumosis secundariis, 15.6.1949, fr.

46923 RGS, ad fl. Capivarí p. Palmares, NEC, in campestribus dumosis, 24.4.1950, fr.

58045 SC, Caixa d'Água p. Itajaí, in dumosis, 18.3.1954, fr., leg. R. Reitz 8310, det. Swallen.

Habit: Prostrate to ascending, rooting at the lower joints, branches up to 50 cm long.

Habitat: Edge of forest, Campos woodlets.

Distribution in RGS: Northeastern Lowlands and sea coast.

General distribution:

Panicum chloroticum Nees, in Trinii Diss. 2 () 236; Döll 2,2 (1877) 197; Herter, Flor. Urug. (1930) 29.

1063 RGS, S. Leopoldo, NEL, in paludosis, 6.4.1935, leg. C. Orth SJ, det. Chase.

1957 RGS, ibidem, in arenosis, 4.10.1934, fr., leg. C. Orth SJ, det. Swallen.

40614 RGS, Esteio p. P. Alegre, NEL, in humidis graminosis, 23.3.1949, fr., det. Swallen.

40742 RGS, Cristal p. P. Alegre, NEL, in paludosis, 31.3.1949, fr., det. Swallen.

40996 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in paludosis, 10.4.1949, fr., det. Swallen.

41130 RGS, Ilha da Pintada p. P. Alegre, NEL, in paludosis graminosis, 20.4.1949, fr., det. Swallen.

46924 RGS, ad fl. Capivari p. Palmares, NEL, in humidis graminosis, 24.4.1950, fr.

Habit: Moderately hick bunches, ascending to erect, 40 cm high.

Habitat: Wet to swampy places.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Amazonia to Uruguay.

Panicum cyanescens Nees, Agrost. Bras. (1829) 220; Döll, FB 2,2 (1877) 262; Ekman, Ark. f. Bot. 13,n.10 (1913) 29; Herter, Flor. Urug. (1930) 29; Reitz, Sellowia 7 (1956) 104.

31768 SC, Sombrio p. Araranguá, in arenosis subhumidis, 7.2.1946, fr., det. Swallen.

43896 RGS, Ad fl. Caí inferius p. P. Alegre, NEL, in arenosis dumosis, 12.10.1949, fr.

Habit: Loosely cespitose, 30 cm high.

Habitat: Moderately wet places.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Amazonia to Uruguay.

Panicum decipiens Nees, Agrost. Bras. (1829) 193; Döll, FB 2,2 (1877) 237; Ekman, Ark. f. Bot. 13,n.10 (1913) 27; Malme, Sv. Bot. Tidskr. 30 (1936) 10; Cabrera, Fl. Alred. B. Aires (1953) 99, fig. 28B; Reitz, Sellowia 7 (1956) 104.

1003 RGS, ad montem Steinkopf p. S. Leopoldo, NEL, in arenosis subhumidis, 8.8.1934, fr., leg. C. Orth SJ, det. Parodi (Chase: *P. milloides* Nees).

8886 RGS, Serra da Rocinha p. Bom Jesus, NEH, in campestribus subturfosis, 15.1.1942, fr., det. Parodi.

9846 RGS, Tupanciretan NWH, in campestribus alte graminosis, 29.1.1942, fr.

37969 RGS, Ponta Grossa p. P. Alegre, NEL, in arenosis humidis, 14.11.1948, fr.

38180 RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis subhumidis, 22.11.1948, fr., det. Swallen.

39914 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in dumosis secundariis, 12.1.1949, fr.

40990 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in campestribus subhumidis, 10.4.1949, fr., det. Swallen.

45750 RGS, Sta Rita p. Farroupilha, NEH, in campestribus dumosis, 7.2.1950, fr.

54323 RGS, Fachinal Preto p. Bom Jesus, NEH, in campestribus humidis, 29.1.1950, fr., leg. R. Reitz, det. Swallen.

57029 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fl.

57133 RGS, Esteio p. S. Leopoldo, NEL, in graminosis, 14.11.1955, fl.

57543 RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 19.11.1955, fl.

60384 RGS, Estação de Silvicultura p. Sta Maria, CL, in graminosis, 8.4.1956, fr., leg. O. Camargo 577.

60606 RGS, Farroupilha, NEH, in graminosis, 7.5.1957, fl., fr., leg. O. Camargo 1362.

Habit: Dense bunches, up to 40 cm high.

Habitat: Moderately dry Campos.

Distribution in RGS: Over the whole of the Campos area.

General distribution: From Goiás and Minas Gerais to Uruguay and B. Aires.

Panicum demissum Trin., Spec. Gram. 27 () tab. 319; Döll, FB 2,2 (1877) 256; Ekman, Ark. F. Bot. 11,n.4 (1912) 23; idem, Ark. f. Bot. 13, n.10 (1913) 28; Malme, Sv. Bot. Tidskr. 30 (1936) 10; Herter, Flor. Urug. (1930) 29.

6959 SC, Lages, in siccis incultis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

34131 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 15.1.1947, fr.

38162 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus dumosis, 20.11.1948, fr.

39116 RGS, Itapoan p. P. Alegre, NEL, in arenosis siccis graminosis, 22.12.1948, fr., det. Swallen.

41003 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in campestribus subsiccis, 10.4.1949, fr.

44342 RGS, ibidem, 11.11.1949, fr.

59380 RGS, Farroupilha, NEH, in graminosis siccis, 25.11.1956, fr., leg. O. Camargo 1006.

Habit: Prostrate to ascending, culms up to 40 cm long.

Habitat: Dry to moderately wet Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: From Minas Gerais to Uruguay.

Panicum dusenii Hack., Fedde Rep. 8 (1910) 513; Ekman, Ark. f. Bot. 13, n.10 (1913) 26.

41695 RGS, Schwabenschneis p. Novo Hamburgo, NES, in dumosis secundariis, 25.5.1949, fr., det. Swallen.

59730 RGS, Farroupilha, NEH, ad silvam, 28.5.1956, fr., leg. O. Camargo 685.

Habit: Vigorous, recumbent, up to 1,5 m high.

Habitat: Edge of forest, secondary growth.

Distribution in RGS: Northeastern rainforest belt and Highlands.

General distribution: Paraná, RGS.

Panicum elephantipes Nees, Agrost. Bras. (1829) 165; Döll, FB 2,2 (1877) 196; Cabrera, Fl. Alred. B. Aires (1953) 98, frg. 28H.

2870 RGS, Granja Boa Vista p. P. Alegre, NEL, in paludosis, 16.12.1937, fr. leg. C. Orth SJ, det. Parodi.

2901 RGS, S. Borja, SW, in humidis alte graminosis, 1943, fr., leg. Fr. Baglione.

33539 RGS, S. Leopoldo, NEL, in humidis dumosis, 10.4.1946, fr., leg. E. Henz SJ.

39759 RGS, Parecí p. Montenegro, NEL, in subpaludosis dumodis, 14.1.1949, fr.

40982 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in paludosis dumosis, 10.4.1949, fr. det. Swallen.

43299 RGS, Estação Azevedo p. Montenegro, NEL, in paludosis alte dumosis, 5.9.1949, fr.

Habit: Vigorous, culms floating, up to 2 m long.

Habitat: Swampy thickets.

Distribution in RGS: Central and northeastern Lowlands.

General distribution: From Pará to RGS.

Panicum fultum Hack., Fedde Rep. 6 (1909) 342; Herter, Flor. Urug. (1930) 29; Cabrera, Fl. Alred. B. Aires (1953) 98; Reitz, Sellowia 7 (1956) 104.

849 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 5.10.1933, leg. C. Orth SJ, det. Chase.

Habit: Erect, 20-30 cm high.

Habitat: Dry Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Uruguay to RGS, Uruguay and B. Aires.

Panicum glutinosum Swartz, Prodromus () 24; Döll, FB 2,2 (1877) 226; Ekman, Ark. f. Bot. 13,n.10 (1913) 26; Malme, Sv. Bot. Tidskr. 30 (1936) 10; Reitz, Sellowia 7 (1956) 104.

6988 SC, Lages, in alte dumosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

7027 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

28343 RGS, Nonoai ad fl. Uruguay, NEU, in silva campestris, 3.1945, fr., det. Swallen.

40548 RGS, Sapucaia p. S. Leopoldo, NEL, in dumosis, 17.3.1949, fr.

40950 RGS, ad montem das Cabras p. S. Leopoldo, in silva campestris, 8.4.1949, fr., det. Swallen.

41040 RGS, Morro Agudo p. P. Alegre, NEL, in campestribus alte dumosis, 13.4.1949, fr., det. Swallen.

41438 RGS, Estação Azevedo p. Montenegro, NEL, in dumosis, 6.5.1949, fr. det. Swallen.

46317 RGS, Cristo Rei p. S. Leopoldo, NEL, in silvula campestris, 17.3.1950, fr.

46404 RGS, Gramado p. Canela, NEH, in araucarieto, 20.3.1950, fr.

46471 RGS, Lagoa da Pinguela p. Osório, NEC, in secundariis, 27.3.1950, fr.

51144 Paraná, Tunas p. Bocaiuva, ad silvam, 4.2.1950, fr., leg. G. Hatschbach 1810, det. Parodi.

52294 Matinhos p. Paranaguá, in dumosis, 7.3.1951, fr., leg. G. Hatschbach 2138, det. Parodi.

55628 SC, in insula Florianópolis, in dumosis, 12.3.1952, fr., leg. R. Reitz 5239, det. Swallen.

58937 RGS, Estação de Silvicultura p. Sta Maria, CL, in silvicis semi-scandens, 3.2.1956, fl., leg. O. Camargo 205.

60383 RGS, ibidem, 8.4.1956, fr. leg. O. Camargo 602.

Habit: Vigorous, recumbent, branches up to 2 m long.

Habitat: Edge of the forest, secondary growth.

Distribution in RGS: Apparently through the whole of the northern forest area and sea coast.

General distribution: From Amazonia to RGS.

Panicum gouini Fourn., Mex, Pl. 2 (1886) 28: Herter, Flor. Urug. (1930) 29: Cabrera, Fl. Alred. B. Aires (1953) 100, fig. 28C, Reitz, Sellowia 7 (1956) 104.

842 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus subpaludosis, 15.10.1933, fr., leg. C. Orth SJ, det. L. Parodi.

Habit: Vigorous, erect, up to 50 cm high.

Habitat: Swampy places.

Distribution in RGS: No other findplaces.

General distribution: Southern USA to Uruguay and B. Aires.

Panicum grumosum Nees, Agrost. Bras. (1829) 182; Döll, FB 2,2 (1977) 208; Ekman, Ark.f. Bot. 11,n.4 (1912) 18; idem, Ark. f. Bot. 13,n.10 (1913) 25; Herter, Flor. Urug. (1930) 29; Cabrera, Fl. Alred. B. Aires (1953) 98, fig. 28D; Reitz Sellowia 7 (1956) 104.

869 RGS, Vila Manresa p. P. Alegre, NEL, in paludosis alte dumosis, 10.10.1933, fr., leg. C. Orth SJ, det. Parodi.

1099 RGS, S. Leopoldo, NEL, in paludosis alte dumosis, 13.10.1934, fr., leg. C. Orth SJ, det. Chase.

7022 SC, Lages, in paludosis herbosis, 1935, fr., leg. A. Bruxel SJ, det. R. Swallen.

39312 RGS, Campo do Varejão p. Itapoan, NEL, in paludosis alte dumosis, 29.12.1948, fr.

39336 RGS, Belém Novo p. P. Alegre, NEL, in paludosis alte graminosis, 31.12.1948, fr.

41748 RGS, Estação Azevedo p. Montenegro, NEL, in paludosis alte cyperaceis, 6.5.1949, fr.

44306 RGS, Esteio p. S. Leopoldo, NEL, in paludosis alte dumosis, 8.11.1949, fr.

49217 RGS, Parecí p. Montenegro, NEL, in paludosis alte dumosis, 26.11.1950, fr.

57243 RGS, Esteio p. S. Leopoldo, NEL, in humidis alte graminosis, 11.11.1955, fl.

Habit: Erect, up to 1,5 m high.

Habitat: Swampy plains, often dominant.

Distribution in RGS: Lowlands.

General distribution: South Brazil to Uruguay and B. Aires.

Panicum gynerioides (Gris.) Pilger,

1105 RGS, Novo Hamburgo, NEL, in paludosis alte dumosis, 20.10.1934, fr., leg. C. Orth SJ, det. Chase, Parodi.

57236 RGS, Esteio p. S. Leopoldo, NEL, ad rivum in alte dumosis, 11.11.1955, fl.

62627 RGS, P. Alegre, NEL, in paludosis, 2.11.1957, fl., leg. O. Camargo 2417.

Habit: Large grayish-green bunches, culms up to 2 m high.

Habitat: Swampy thickets.

Distribution in RGS: Northeastern Lowlands.

General distribution:

Paspalum hebotes Trin., Act. Petrop. () 301; Döll, FB 2,2 (1877) 252.

1938 RGS, Amaral Ribeiro p. Novo Hamburgo, NEL, in arenosis dumosis, 4.10.1934, fr., leg. C. Orth SJ, det. Chase.

Habit: Slender, recumbent, branches up to 1 m long.

Habitat: Secondary growth.

Distribution in RGS: Not other findplace.

General distribution: From Minas Gerais to RGS.

Panicum helobium Mez, ap. Ekman, Ark. f. Bot. 11,n.4 (1912) tab.1; idem, Ark. f. Bot. 13., n.10 (1913) 30; Malme, Sv. Bot. Tidskr. 30 (1936) 10; Cabrera, Fl. Alred. B. Aires (1953) 99; Reitz, Sellowia 7 (1956) 104.

1925 RGS, S. Leopoldo, NEL, in stagno, 4.1.1934, fr., leg. C. Orth SJ, det. Swallen.

6960 SC, Lages, in stagnosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

8772 RGS, Bom Jesus, in stagno, 15.1.1942, fr., det. Swallen.

31452 SC, Sombrio p. Araranguá, in stagnosis, 31.1.1946, fr., det. Swallen.

32349 RGS, Tainhas p. S. Francisco de Paula, NEH, in paludosis, 17.2.1946, fl., det. Swallen.

32419 RGS, Serra da Rocinha p. Bom Jesus, NEH, in paludosis, 28.2.1946, fr., det. Swallen.

35163 RGS, Fazenda da Ronda p. Vacaria, NEH, in stagno, 15.1.1947, fr.

40362 RGS, Sta Rita p. Farroupilha, NEH, in stagnosis, 28.1.1949, fr.

44892 RGS, S. Francisco de Paula, NEH, in paludosis, 18.12.1949, fr.

45567 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in sphagno-to, 30.1.1950. fr.

54921 Vila Oliva p. Caxias, NEH, in stagnosis, 24.2.1954, fr.

55362 SC, Massiambú p. Palhoça, in stagnosis, 5.2.1953, fr., leg. R. Reitz 7164, det. Swallen.

55363 SC, Serra da Boa Vista p. S. José, 2.2.1953, fr., leg. R. Reitz 7163, det. Swallen.

56931 RGS, Caracol p. Canela, NEH, in stagno, 19.11.1955, fl., fr., leg. K. Emrich.

58566 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in stagno, 12.2.1956, fr.

58940 RGS, Estação de Silvicultura p. Sta Maria, CL, in stagnosis, 28.2.1956, fr., leg. O. Camargo 40.

62854 RGS, Bom Jesus, NEH, in stagno, 6.1.1958, fl., leg. O. Camargo 3112.

64560 RGS, Fazenda do Arroio p. Osório, NEC, in stagno, 23.1.1958, fl., fr.

Habit: Erect to floating.

Habitat: Stagnant, more or less perennial ponds, often dominant.

Distribution in RGS: Over the whole of the Campos area.

General distribution: From Paraná to RGS, Uruguay and B. Aires.

Panicum latiglume Döll, FB 2,2 (1877) 257; Herter, Flor. Urug. (1930) 29; Reitz, Sellowia 7 (1956) 104.

38181 RGS, Sapucaia p. S. Leopoldo, NEL, in alte dumosis, 22.11.1948, fr., det. Swallen.

44638 RGS, Vila Oliva p. Caxias, NEH, in dumosis secundariis, 3.12.1949, fr.

54532 RGS, Taimbesinho p. S. Francisco de Paula, NEH, ad rivum in rupestribus humidis, 13.11.1953, fr., det. Swallen (aff. latiglume).

55891 RGS, Fazenda do Arroio p. Osório, NEC, in arenosis dumosis, 3.10.1954, fl., det. Parodi, Swallen.

56050 RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 19.11.1954, fr., det. Parodi (Swallen: P. sabulorum Lam.).

57130 RGS, Esteio p. S. Leopoldo, NEL, in graminosis, 14.11.1955, fr.

57223 RGS, Vila Manresa p. P. Alegre, NEL, in graminosis, 15.10.1955, fl.

57444 RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in dumosis, 9.10.1955, fr.

59525 RGS, Farroupilha, NEH, in siccis graminosis, 26.9.1956, fr., leg. O. Camargo 808.

62006 RGS, Taquarí, NEL, in incultis, 14.12.1957, fl., leg. O. Camargo 2973.

62064 RGS, Bento Gonçalves, NEH, in dumosis, 6.10.1957, fl., leg. O. Camargo 1970.

Habit: Prostrate to ascending, grayish-green, up to 50 cm high.

Habitat: Dry, more or less open, places.

Distribution in RGS: Apparently over the whole of the open area.

General distribution: SC and RGS.

Panicum laxum Swartz, Prodr. (1788) 23; Döll, FB 2,2 (1877) 212; Ekman, Ark.f. Bot. 11,n.4 (1912) 19; idem, Ark.f. Bot. 13,n.10 (1913) 25; Herter, Flor.Urug. (1930) 29; Cabrera, Flor.Alred. B. Aires (1953) 98, Fig. 28L; Reitz, Sellowia 7 (1956) 104.

1924 RGS, S. Leopoldo, NEL, in campestribus dumosis, 30.3.1935, fr., leg. C. Orth SJ, det. Chasse, Parodi, Beetle.

6952 SC, Lages, in campestribus graminosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

33055 RGS, Pareci p. Montenegro, NEL, in campestribus arenosis, 28.12.1945, fr., leg. A. Strieder SJ, det. Swallen.

39941 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in dumosis, 12.1.1949, fr., det. Swallen.

40116 RGS, Guaíba p. P. Alegre, NEL, in arenosis, 24.1.1949, fr., det. Beetle.

40131 RGS, ibidem, 24.1.1949, fr., det. Swallen.

40981 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in campestribus subsiccis, 10.4.1949, fr., det. Swallen.

41069 RGS, in humidis alte graminosis, 18.4.1949, fr., det. Swallen.

55383 SC, Ascurra p. Rodeio, ad viam, 11.6.1953, fr., leg. R. Reitz 7136, det. Swallen.

Habit: Prostrate to ascending, slender, branches up to 50 cm long.

Habitat: Dry shady places among shrubs and trees.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Amazonia to Uruguay and B. Aires.

Panicum maximum Jacq., Collect. 1 () 76; Döll, FB 2,2 (1877) 202; Ekman, Ark.f.Bot. 13, n.10 (1913) 25; Reitz, Sellowia 7 (1956) 104.

1016 RGS, Sapucaia p. S. Leopoldo, NEL, in incultis dumosis, 15.6.1934, fr., leg. C. Orth SJ, det. Chase, Parodi.

32662 RGS, Parecí p. Montenegro, NEL, in dumosis secundariis, 22.10.1945, fr.

40916 RGS, ad montem das Cabras p. S. Leopoldo, in silva secundaria, 8.4.1949, fr., det. Swallen.

41210 RGS, Instituto Borges de Medeiros p. P. Alegre, NEL, ad rivum in dumosis, 4.1949, fr., leg. G.C.Leal.

52617 SC, Cedro p. Criciuma, ad flumen in dumosis, 30.11.1943, fr., leg. R. Reitz 1018, det. Serviço Florestal.

Habit: Culms up to 1,5 m high.

Habitat: Among shrubs and in secondary growth.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Amazonia to RGS.

Panicum milloides Nees, Agrost.Bras. (1829) 175; Herter, Flor. Urug. (1930) 29; Cabrera, Flor. Alred. B. Aires (1953) 99. — **P. hians** Elliott Bot. Carol. et Georg. 1 () 118; Döll, FB 2,2 (1877) 240; Ekman, Ark.f.Bot. 11,n.4 (1912) 21.

6993 SC, Lages, in campestribus siccis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

28279 RGS, Nonoai ad fl. Uruguay, NWU, in campestribus siccis graminosis, 3.1945, fr.

33196 RGS, Vila Oliva p. Caxias, NEH, in campestribus subhumidis, 16.1.1946, fr.

33428 RGS, S. Leopoldo, NEL, in campestribus siccis graminosis, 17.6.1946, fr., det. Beetle.

37387 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 5.7.1948, fr.

44341 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in campestribus siccis, 11.11.1949, fr.

55994 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus graminosis, 19.11.1954, fr., det. Parodi (Swallen: *P. schenckii* Hack.)

59381 RGS, Farroupilha, NEH, in dense graminosis, 15.11.1956, fl., leg. O. Camargo 932.

Habit: Culms slender, ascending to erect, up to 50 cm high.

Habitat: Dry, more or less open Campos.

Distribution in RGS: Apparently over the whole of the Campos area.

General distribution: From Paraná to RGS, Uruguay and B. Aires.

Panicum millegrana Poir., Encycl. Méth. 4 () 278; Ekman, Ark. f. Bot. 11,n.4 (1912) 23; idem, Ark.f.Bot. 13, n.10 (1913) 29; Reitz, Sellowia 7 (1956) 104. — *Panicum rugulosum* Trin., Diss. 2 () 194; Döll, FB 2,2 (1877) 258.

2717 RGS, Kappesberg p. Montenegro, NES, in dumosis, 4.1.1936, fr., leg. C. Orth SJ, det. Chase.

28637 RGS, Nonoai, ad fl. Uruguay, NWU, in silva campestri, 3.1945, fr., det. Swallen.

30233 Uruguay, Treinta y Tres, ad silvulam in umbrosis, 3.1945, fr. leg. B. Rosengurtt B-4819, det. Parodi.

46480 RGS, Lagoa da Pinguela p. Osório, NEC, in silva secundaria, 27.3.1950, fr., det. Parodi.

55084 RGS, Vila Oliva p. Caxias, NEH, in dumosis, 24.2.1954, fr., det. Swallen.

56613 RGS, ibidem, 8.2.1955, fr., det. Parodi.

56746 RGS, ibidem, 8.2.1955, fr., det. Parodi.

60275 RGS, Farroupilha, NEH, in dumosis, 14.1.1957, fr., leg. O. Camargo 1082.

63085 RGS, Pelotas, SEC, in alte graminosis, 11.3.1956, fl., leg. Schlichting.

Habit: Vigorous, recumbent, branches up to 1,5 long.

Habitat: Edge of forest, secondary growth.

Distribution in RGS: Northeastern quadrant.

General distribution: Amazonia to RGS.

Panicum missionum Mez,

857 RGS, Vila Manresa p. P. Alegre, NEL, ad silvam in umbrosis, 16.10.1933, fr., leg. C. Orth SJ, det. Chase.

8738 RGS, Serra da Rocinha p. Bom Jesus, NEH, in campestribus alte graminosis, 14.1.1942, fr., det. Swallen.

9808 RGS, Tupanciretan, NWH, in dumosis, 29.1.1942, fr., det. Swallen.

27809 São Paulo, Campos do Jordão, in campestribus, 1.1944, fr., leg. E. Friderichs SJ, det. Swallen.

36459 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus alte graminosis, 2.1948, fr., det. Swallen.

55059 RGS, Vila Oliva p. Caxias, NEH, in dumosis, 24.2.1954, fr., det. Swallen.

55088 RGS, ibidem, 24.2.1954, fr., det. Swallen.

Habit: Ascending to erect, rooting at the lower joints, branches up to 40 cm long.

Habitat: Edge of the forest, secondary growth.

Distribution in RGS: Apparently over the whole of the northern half of the State.

General distribution: From Paraná and Misiones to RGS.

Panicum numidianum Lam.,

1062 RGS, S. Leopoldo, NEL, in incultis, 10.8.1934, fr., leg. C. Orth SJ, det. Chase.

Habit: Culms up to 2 m high.

Habitat: On formerly cultivated soil.

Distribution in RGS: Only findplace.

General distribution:

Panicum olyroides H.B.K., Nov. Gen. et Spec. 1 () 102; Döll, FB 2,2 (1877) 229; Ekman, Ark.f.Bot. 11,n.4 (1912) 21; Ekman, Ark.f.Bot. 13, n.10 (1914) 27.

41286 RGS, Morro da Sta Teresa p. P. Alegre, NEL, in campestribus siccis, 29.4.1949, fr., det. Swallen.

44477 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis alte graminosis, 19.11.1949, fr., det. Swallen.

Habit: Erect, stiff, 50 cm high.

Habitat: Dry Campos.

Distribution in RGS: Granitic hills of P. Alegre.

General distribution: From Goiás to RGS.

Panicum ovuliferum Trin., Diss. 2 () 191; Döll, FB 2,2 (1877) 226; Ekman, Ark. f. Bot. 13,n.10 (1913) 26.

2722 RGS, Kappesberg p. Montenegro, NES, in secundariis, 4.1.1936, fr., leg. C. Orth SJ, det. Chase.

28796 RGS, Caracol p. Canela, NEH, in dumosis, 2.1945, fr., leg. K. Emrich.

42302 Langschneis p. Montenegro, NES, in dumosis, 2.7.1949, fr.

60825 RGS, Sta Rita p. Farroupilha, NEH, in dumosis, 18.5.1957, fr., leg. O. Camargo 1472.

64225 SC, S. Joaquim, 10.1.1958, fl., leg. J. Mattos.

64226 SC, ibidem, 10.1.1958, fl., leg. J. Mattos.

Habit: Vigorous, recumbent, branches up to 1,5 high.

Habitat: Edge of forest, secondary growth.

Distribution in RGS: Northeastern forest belt and Highlands.

General distribution: From Minas Gerais to RGS.

Panicum paludivagum Hitchc. et Chase,

730 RGS, Lagoa dos Barros p. Osório, NEC, in lacu fluitans, 3.2.1933, fr., leg. C. Orth SJ, det. Swallen.

Habit: Branches long, floating.

Habitat: Shallow freshwater lake.

Distribution in RGS: Northeastern sea coast.

General distribution: RGS, Uruguay.

Panicum pantrichum Hack.,

Reitz, Sellowia 7 (1956) 104.

1018 RGS, S. Leopoldo, NEL, in agro, 8.8.1934, fr., leg. C. Orth SJ, det. Beetle.

7452 RGS, ibidem, 1907, fr., leg. F. Theissen SJ, det. Swallen.

41664 RGS, ibidem, in arenosis dumosis, 21.5.1949, fr., det. Swallen.

Habit: Slender, prostrate to ascending, branches up to 75 cm long.

Habitat: Dry, shrubby Campos.

Distribution in RGS: S. Leopoldo.

General distribution:

Panicum pilosum Swartz, Prodr. (1788) 22; Döll, FB 2,2 (1877) 210; Ekman, Ark. f. Bot. 11,n.4 (1912) 19; idem, Ark.f.Bot. 13, n.10 (1913) 25; Herter, Flor. Urug. (1930) 29; reitz, Sellowia 7 (1956) 105.

1022 SC, Itapiranga ad fl. Uruguay superius, in agro, 15.2.1934, fr., det. Chase.

55338 SC, Anitápolis p. Palhoça, in dumosis, 2.4.1953, fr., leg. R. Reitz 7194, det. Swallen.

58043 SC, Caixa d'Água p. Itajaí, in humidis, 18.3.1954, fr., leg. R. Reitz 8308, det. Swallen.

58053 SC, Ibirama, ad silvam, 1.3.1954, fr., leg. R. Reitz 8331, det. Swallen.

Habit: Erect, 40 cm high.

Habitat: Edge of forest, secondary growth.

Distribution in RGS: Near the northeasternmost frontier.

General distribution: From Piauí to Uruguay.

Panicum polycladum Ekman, Ark. f. Bot. 11,n.4 (1912) 24, tab. 3. fig. 2.

29194 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 12.9.1945, fr., det. Swallen.

30733 RGS, Vila Oliva p. Caxias, NEH, in dumosis, 28.1.1946, fr.

43981 RGS, Sapucaia p. S. Leopoldo, NEL, in campestribus alte dumosis, 19.10.1949, fr., det. Swallen.

Habit: Erect, 50 cm high.

Habitat: Dry, shrubby Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: Misiones, RGS.

Panicum polygonatum Schrad., in Schultes, Mant. 2 () 256. — **Panicum pilosum** Swartz, var. **polygonatum** (Schrad.) Döll, FB 2,2 (1877) 211.

1796 RGS, Vila Manresa p. P. Alegre, NEL, in humidis alte dumosis, 16.1.1935, fr., leg. C. Orth SJ, det. Chase.

46329 RGS, Cristo Rei p. S. Leopoldo, NEL, in arenosis alte dumosis, 17.3.1950, fr.

55347 SC, Massiambú p. Palhoça, in umbrosis dumosis, 12.3.1953, fr., leg. R. Reitz 7184, det. Swallen.

Habit: Prostrate to ascending and recumbent, branches up to 75 cm long.

Habitat: Shrubby Campos and secondary growth.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Amazonia to RGS.

Panicum procurrens Nees, Agrost. Bras. (1829) 130; Döll, FB 2,2 (1877) 271; Ekman, Ark.f.Bot. 11,n.4 (1912) 25; idem, Ark. f. Bot. 13,n.10 (1913) 30.

6997 SC, Lages, in campestribus siccis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

8771 RGS, Bom Jesus, NEH, in campestribus siccis, 15.1.1942, fr.

8980 RGS, ibidem, 16.1.1942, fr.

9466 RGS, Jarí p. Tupanciretan, NWH, in campestribus siccis, 27.1.1942, fr.

10000 RGS, Ijuisinho p. Tupanciretan, NWH, in campestribus siccis, 30.1.1942, fr., det. Parodi.

30734 RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis, 28.1.1946, fr., det. swallen.

35139 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 15.1.1947 kr.

35179 RGS, ibidem, 15.1.1947, fr.

59023 RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus siccis, 12.2.1956, fr., leg. O. Camargo 168.

Habit: Erect, very elegant, 30 cm high.

Habitat: Dry Campos.

Distribution in RGS: Over the whole of the Highlands, not frequent.

General distribution: From Piauí to RGS.

Panicum racemosum Spreng., Syst.Veg. 1 (1825) 313; Herter, Flor. Urug. (1930) 30; Reitz, Sellowia 7 (1956) 105. — Döll, FB 2,2 (1877) 259; P. reptans Kunth, Rev. () tab. 21; Malme, Sv. Bot. Tidskr. 30 (1936) 10.

31890 SC, Sombrio p. Araranguá, in arenosis submobilibus, 9.10.1945, fr., leg. R. Reitz 1307.

37988 RGS, Barra do Ribeiro p. P. Alegre, NEL, in arenosis dumosis, 14.11.1948, fr.

44552 RGS, Osório, NEC, in arenosis submobilibus dumosis, 24.11.1948, fr.

56452 RGS, Cidreira, NEC, in arenosis, 17.12.1954, fl.

58037 SC, Barra do Sul p. Araquari, in arenosis, 29.10.1953, fr., det. leg. R. Reitz 8289, det. Swallen.

Habit: Culms from a rhizome, up to 75 cm high.

Habitat: More or less mobile sand dunes.

Distribution in RGS: Along sea coast.

General distribution: From Rio de Janeiro to Uruguay and B. Aires.

Panicum repens L., Spec.Plant. 2 ed. (1762) 24; Ekman, Ark.f.Bot. 13,n.10 (1913) 24; Reitz, Sellowia 7 (1956) 105.

30979 RGS, sine loco, sine die, fr.

46320 RGS, Cristo Rei p. S. Leopoldo, NEL, in subpaludosis graminosis, 17.3.1950, fr., det. Parodi (cum?).

Habit: Culms 10 cm high.

Habitat: Wet sandy places.

Distribution in RGS: Northeastern Lowlands.

General distribution: Paraná to RGS.

Panicum rhizogonum Hack.,

Herter, Flor. Urug. (1930) 30; Reitz, Sellowia 7 (1956) 105.

922 RGS, S. Leopoldo, NEL, ad silvam pluviale, 5.11.1934, fr., leg. C. Orth SJ, det. Chase, Parodi.

35175 RGS, Fazenda da Ronda p. Vacaria, in dumosis, 15.1.1947, fr.
41883 RGS, Reutersberg p. S. Leopoldo, NES, in secundariis, 6.6.1949, fr., det. Swallen.

42280 RGS, ad montem Steinkopf p. S. Leopoldo, NEL, in silva rupestri, 28.6.1949, det. Swallen.

55421 SC, Massiambú p. Florianópolis, in arenosis, 12.3.1953, fr., leg. R. Reitz 7070, det. Swallen.

59675 RGS, Farroupilha, NEH, in dumosis, 16.5.1956, fr., leg. O. Camargo 662.

Habit: Prostrate to ascending, rooting at the lower joints, branches up to 75 cm long.

Habitat: Edge of rain forest, secondary growth.

Distribution in RGS: Northeastern Lowlands, rain forest belt, and Highlands.

General distribution: From Paraná to RGS.

Panicum rivulare Trin., Diss 2 () 213; Döll, FB 2,2 (1877) 208; Ekamn, Ark, f. Bot. 11,n.4 (1912) 18; Ark.f.Bot. 13, n.10 (1913) 25; Herter, Flor. Urug. (1930) 30.

10839 RGS, Cerro Largo p. S. Luiz, NWH, in paludosis alte dumosis, 1.1943, fr., leg. P. Buck SJ, det. Swallen.

32772 RGS, Parecí p. Montenegro, ad flumen in alte dumosis, 3.11.1945, fr., leg. E. Henz SJ.

38317 RGS, Esteio p. S. Leopoldo, NEL, in paludosis alte dumosis, 24.11.1948, fr., det. Swallen.

44089 RGS, Barreto Vianna p. S. Leopoldo, NEL, in paludosis alte dumosis 24.10.1949, fr.

62005 RGS, Taquarí, NEL, in humidis alte graminosis, 14.12.1957, fl., leg. O. Camargo 2968.

Habit: Robust, erect, up to 1,5 m high.

Habitat: Swampy thickets, often dominant.

Distribution in RGS: Apparently over the whole of the Campos area.

General distribution: From Minas Gerais to Uruguay.

Panicum sabulorum Lam., Encycl. Méth. 4 (1798) 744; Herter, Flor. Urug. (1930) 30; Cabrera, Fl. Alred. B. Aires (1953) 99, fig. 28J; Reitz; Sellowia 7 (1956) 105.

1014 RGS, S. Leopoldo, NEL, in arenosis siccis graminosis, 5.10.1934, fr., leg. C. Orth SJ, det. Parodi (Chase: P. latiglume Döll).

6748 SC, Florianópolis, in arenosis submobiliibus, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

26512 RGS, Parecí p. Montenegro, NEL, in arenosis alte graminosis, 1944, fr., leg. E. Henz SJ.

- 29528** RGS, Sapucaia p. S. Leopoldo, NEL, in dumosis umbrosis, 5.9.1945, fr., det. Swallen.
- 29573** RGS, Pareci p. Montenegro, NEL, in arenosis graminosis, 3.10.1956, fr., leg. E. Henz SJ.
- 31893** SC, Sombrio p. Araranguá, in arenosis graminosis, 9.10.1945, fr., leg. R. Reitz 1292, det. Swallen.
- 33501** RGS, S. Leopoldo, NEL, in campestribus alte graminosis, 17.7.1946, fl., leg. E. Henz SJ.
- 36460** RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus graminosis, 2.1948, fr., det. Swallen.
- 38536** RGS, S. Leopoldo, NEL, in campestribus, 4.12.1948, fr.
- 39854** RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in arenosis dumosis, 12.1.1949, fr.
- 41342** RGS, Morretes p. P. Alegre, NEL, in arenosis graminosis, 2.5.1949, fr.
- 44279** RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis graminosis, 5.11.1949, fr.
- 44387** RGS, Fião p. S. Leopoldo, NEL, in arenosis dumosis, 14.11.1949, fr.
- 45382** RGS, Serra da Rocinha p. Bom Jesus, NEH, in rupestribus subhumidis, 18.1.1950, fr., det. Parodi (cum?).
- 46152** RGS, Fazenda do Arroio p. Osório, NEC, in arenosis graminosis, 6.3.1950, fr.
- 46162** RGS, ibidem, 6.3.1950, fr., det. Parodi.
- 56468** RGS, Torres, NEC, prope litus in arenosis, 19.1.1955, fr.
- 57011** RGS, in summo monte Sapucaia, p. S. Leopoldo, NEL, in rupestribus graminosis, 26.10.1955, fr.
- 57015** RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fr.
- 57028** RGS, ibidem, 26.10.1955, fr.
- 57136** RGS, Vila Manresa p. P. Alegre, NEL, in graminosis, 4.11.1955, fr.
- 57209** RGS, ibidem, 22.10.1955, fl.
- 57210** RGS, ibidem, 22.10.1955, fr.
- 57211** RGS, ibidem, 22.10.1955, fr.
- 57213** RGS, ibidem, 15.10.1955, fr.
- 57241** RGS, Esteio p. P. Alegre, NEL, in alte graminosis, 11.11.1955, fr.
- 58939** RGS, Estação de Silvicultura p. Sta Maria, CL, in siccis graminosis, 1.3.1956, fr., leg. O. Camargo 103.
- 62578** RGS, Passo Fundo, CH, in campestribus tritis, 20.10.1957, fl., leg. O. Camargo 2160.
- 62579** RGS, ibidem, 20.10.1957, fl., leg. O. Camargo 2253.

63380 RGS, Pelotas, SEC, in arenosis, 19.10.1954, fl., leg. J. Sacco 202.

63383 RGS, Pelotas, SC, in arenosis, 5.10.1954, fl., leg. J. Sacco 187, det. Parodi.

Habit: Prostrate to ascending, rooting at the lower joints, branches up to 50 cm long.

Habitat: Dry, preferably sandy places.

Distribution in RGS: Common over the whole of the Campos area.

General distribution: From Paraná to Uruguay.

Panicum schenckii Hack., österr. Bot. Zeitschr. 51 (1901) 426; Ekman, Ark.f.Bot. 11,n.4 (1912); 22; Reitz, Sellowia 7 (1956) 105.

29376 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus dumosis, 3.11.1945, fr., det. Swallen (vel aff.).

39261 RGS, Morro do Sabiá p. P. Alegre, NEL, ad flumen in paludosis, 28.12.1948, fr., det. Swallen.

35143 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 15.1.1947, fr.

41186 RGS, Ilha das Flores p. P. Alegre, NEL, in humidis graminosis, 22.4.1949, fr., det. Swallen.

43761 RGS, Morro do Sabiá p. P. Alegre, NEL, ad flumen in humidis, 5.10.1949, fr., det. Swallen.

Habit: Thick bunches, up to 50 cm high.

Habitat: Dry to more or less wet places.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: From Paraná to Uruguay.

Panicum secundum Trin., Spec. Gram. () tab. 234; Döll, FB 2,2 (1877) 194; Reitz, Sellowia 7 (1956) 105.

2510 RGS, sine loco, sine die, det. Chase.

47605 SC, in summo Morro do Baú p. Itajaí, in dumosis, 27.10.1947, fr., leg. R. Reitz 3057, det. Parodi.

47708 SC, ibidem, 29.1.1948, leg. R. Reitz 2982, det. Parodi.

50787 SC, Brusque, in dumosis, 20.9.1948, fr., leg. R. Reitz 3547, det. Swallen.

50341 SC, ad montem Cambirela p. Florianópolis, in humidis dumosis, 18.7.1951, fr.

Habit: Culms erect, up to 50 cm high.

Habitat: Edge of forest, secondary growth.

Distribution in RGS: Unknown, probably northeastern Lowlands.

General distribution: From Rio de Janeiro to RGS.

Panicum sellowii Nees, Agrost. Bras. (1929) 153. — *P. rugulosum* Trin., Diss. 2 () 195; Döll, FB 2,2 (1877) 258.

1064 RGS, S. Leopoldo, NEL, in arenosis alte dumosis, 19.8.1934, fr., leg. C. Orth SJ, det. Swallen.

2521 RGS, Sapucaia p. S. Leopoldo, NEL, ad silvam campestrem, 13.4.1936, fr., leg. C. Orth SJ, det. Chase.

39805 RGS, Pareci p. Montenegro, NEL, in silvam campestrem, 14.1.1949, fr.

40214 RGS, Sta Rita p. Farroupilha, NEH, in alte dumosis, 27.1.1949, fr., det. Swallen.

40254 RGS, ibidem, 28.1.1949, fr., det. Swallen.

40852 RGS, Granja Neugebauer p. Itapoan, NEL, in silvula campestri, 3.4.1949, fr.

40922 RGS, ad montem das Cabras p. S. Leopoldo, NEL, in alte dumosis, 8.4.1949, fr., det. Swallen.

41425 RGS, Estação Azevedo p. Montenegro, NEL, in silvula campes- tri, 6.5.1949, fr.

41579 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in umbrosis, 16.5.1949, fr., det. Swallen.

41738 RGS, Esteio p. S. Leopoldo, NEL, in silva campestri, 29.5.1949, fr. det. Swallen.

41891 RGS, Reutersberg p. S. Leopoldo, NES, in silvula secundaria, 6.6.1949, fr., det. Swallen.

42015 RGS, Morro Santana p. P. Alegre, NEL, in silvula campestri, 15.6.1949, fr.

47049 RGS, Lagoa da Pinguela p. Osório, NEC, in umbrosis, 8.5.1950, fr.

47094 RGS, Vila Manresa p. P. Alegre, NEL, in humidis alte dumosis, 15.5.1950, fr., det. Parodi.

54961 RGS, Vila Oliva p. Caxias, NEH, in dumosis, 24.2.1954, fr., det. Swallen.

56779 RGS, ibidem, 8.2.1955, fr., det. Parodi, Swallen.

59696 RGS, Farroupilha, NEH, in alte dumosis, 21.5.1956, fr., leg. O. Camargo 683.

Habit: Robust, prostrate to ascending, recumbent, rooting at the lower joints, branches up to 1,5 m long.

Habitat: Edge of rain forest, secondary growth.

Distribution in RGS: Northeastern quadrant.

General distribution: From Amazonia to RGS.

Panicum stigmosum Trin., Diss. 2 () 194; Döll, FB 2,2 (1877) 225; Reitz, Sellowia 7 (1956) 105.

1104 RGS, S. Leopoldo, NEL, in arenosis siccis, 10.10.1934, fr., leg. C. Orth SJ, det. Swallen.

- 32574 RGS, Pareci p. Montenegro, NEL, in arenosis dumosis, 30.10.1945, fr., det. Swallen.
- 32712 RGS, Vila Manresa p. P. Alegre, NEL, in umbrosis dumosis, 22.9.1948, fr., det. Swallen.
- 33865 RGS, ibidem, 3.10.1946, fr.
- 34205 RGS, ibidem, 16.10.1946, fr.
- 38758 RGS, Conceição p. Caí, NEL, in campestribus alte graminosis, 14.12.1948, fr., det. Swallen.
- 41528 RGS, Morro do Osso p. P. Alegre, NEL, in campestribus dumosis, 13.5.1949, fr., det. Swallen.
- 43613 RGS, Morro da Polícia p. P. Alegre, NEL, in umbrosis dumosis, 28.9.1949, fr., det. Swallen.
- 50719 SC, Campo dos Padres p. Bom Retiro, in campestribus dumosis, 20.12.1948, fr., leg. R. Reitz 2595, det. Swallen.
- 54457 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 12.11.1953, fr., det. Swallen.
- 56342 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in dumosis, 2.1.1955, fr., det. Parodi.
- 56818 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in umbrosis, humidis, 16.2.1955, fr., det. Parodi.
- 57131 RGS, Esteio p. S. Leopoldo, NEL, in dumosis, 14.11.1955, fr.
- 59442 RGS, Farroupilha, NEH, in dumosis, 25.11.1956, fr., leg. O. Camargo 1008.
- 62762 RGS, P. Alegre, NEL, in campestribus, 1.11.1957, fl., leg. O. Camargo 2409.
- 62763 RGS, ibidem, 1.10.1957, fl., leg. O. Camargo 1837.

Habit: Prostrate to ascending, branches up to 50 cm high.

Habitat: Dry, shady places.

Distribution in RGS: Northeastern quadrant.

General distribution: From Minas Gerais to RGS.

Panicum stoloniferum Poir., Encycl. Méthod, Suppl. 4 () 274; Döll, FB, 2,2 (1877) 209; Herter, Flor. Urug. (1930) 30.

37751 RGS, sine loco, sine die, ster.

38269 RGS, Esteio p. S. Leopoldo, 1947, fr., in dumosis, det. Swallen.

Habit: Ascending to recumbent, branches up to 50 cm high.

Habitat: Dry places among shrubs.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Central America to Uruguay.

Panicum subjuncuum Ekman, Ark.f.Bot. 10,n.17 (1911) 16, tab. 5, fig. 2; tab. 6, fig. 8.

2732 RGS, S. Leopoldo, NEL, in dumosis siccis, 23.10.1935, fr., leg. C. Orth SJ, det. Chase.

63370 RGS, Pelotas, SEC, in alte graminosis, 14.10.1954, fl., leg. Sacco, det. Parodi.

Habit: Robust, up to 1 m high.

Habitat: Among high shrubs.

Distribution in RGS: Widely scattered.

General distribution: From Paraná to RGS.

Panicum superatum Hack.,

53841 RGS, Serra da Rocinha p. Bom Jesus, NEH, in humidis dense graminosis, fr., det. Swallen.

Habit: Moderately thick bunches, erect, 30 cm high.

Habitat: Wet, grassy, campos.

Distribution in RGS: Northeastern Highlands.

General distribution:

Panicum trichanthum Nees, Agrost. Bras, (1829) 210; Döll, FB 2,2 (1877) 247.

4642 RGS, Estreito p. Marcelino Ramos, NWU, ad flumen in rupestribus, 16.2.1942, fr., det. Parodi.

41350 RGS, Morretes p. P. Alegre, NEL, in silvula riparia, 2.5.1949, fr., det. Swallen.

42068 RGS, Sapucaia p. S. Leopoldo, NEL, in silva campestri, 18.6.1949, fr.

45722 RGS, Farroupilha, NEH, in alte dumosis, 7.2.1950, fr.

49843 SC, Itapiranga ad fl. Uruguay superius, ad silvam pluvialem, 6.2.1951, fr.

56764 RGS, Vila Oliva p. Caxias, NEH, in paludosis dumosis, 8.2.1955, fr., det. Parodi (Swallen: P. helobium Mez).

Habit: Slender, recumbent, branches up to 50 cm long.

Habitat: Edge of forest, secondary growth.

Distribution in RGS: Apparently over the whole of the northern half of the State.

General distribution: From the West Indies to RGS.

Panicum tricholaenoides Steud., Syn.Gram. (1855) 68; Herter, Flor. Urug. (1930) 30; Cabrera, Fl. Alred. B. Aires (1953) 99. — **P. junceum** Nees, Agrost.Bras. (1929) 159; Döll, FB 2,2 (1877) 216.

850 RGS, Pedra Redonda p. P. Alegre, NEL, in incultis alte dumosis, 5.1.1933, fr., leg. C. Orth, det. Parodi (Swallen: P. junceum Nees)

5740 RGS, Esteio p. S. Leopoldo, NEL, in humidis alte graminosis, 11.11.1955, fl.

Habit: Culms up to 1,5 m high.

Habitat: Densely vegetated swamps.

Distribution: Northeastern Lowlands.

General distribution: From Goiás to Uruguay.

PASPALUM L.

Paspalum acuminatum Raddi, Agrost. Bras. (1828) 25; Döll, FB 2,2 (1877) 97.

40355 RGS, Sta Rita p. Farroupilha, NEH, in paludosis dumosis, 29.1.1949, fl.

40625 RGS, Esteio p. S. Leopoldo, NEL, in paludosis alte dumosis, 23.3.1949, fr., det. Swallen.

40972 RGS, ad fl. Gravataí inferius, in subpaludosis dumosis, 10.4.1949, fr., det. Swallen.

41122 RGS, Ilha da Pintada p. P. Alegre, NEL, in paludosis dumosis, 20.4.1949, fr., det. Swallen.

Habit: Prostrate, rooting at most joints, branches up to 1 m long.

Habitat: Swampy thickets.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Rio de Janeiro to RGS.

Paspalum arechavaletae Hack.,

Herter, Flor. Urug. (1930) 30.

44124 RGS, Barreto Vianna p. S. Leopoldo, NEL, in humidis alte dumosis, 24.10.1949, fr., det. Swallen. (Barreto: P. exaltatum Presl).

Habit: Robust, erect, up to 1,5 m high.

Habitat: Swampy thickets.

Distribution in RGS: Northeastern Lowlands.

General distribution: RGS, Uruguay.

Paspalum arenarium Schrad., ap. Roem. et Schult. Mantissa 2 () 172; Döll, FB 2,2 (1877) 46; Herter, Flor. Urug. (1930) 30; Reitz, Sellowia 7 (1956) 106.

31405 SC, Sombrio p. Araranguá, in arenosis, 31.1.1946, fr., det. Swallen.

33053 RGS, Parecí p. Montenegro, NEL, in dumosis, 28.2.1945, fl., leg. A. Strieder SJ.

54351 SC, Laguna, in arenosis, 22.12.1951, fr., leg. R. Reitz 5053, det. Swallen.

54352 SC, ibidem, leg. R. Reitz 5054, det. Swallen.

54785 RGS, Torres, NEC, prope litus in arenosis, 11.2.1954, fr., det. Parodi.

54839 RGS, ibidem, 11.2.1954, fr., det. Parodi.

55378 SC, Massiambú p. Palhoça, in arenosis, 19.12.1953, fr., leg. R. Reitz 7144, det. Swallen.

56472 RGS, Torres, NEC, prope litus in arenosis, 19.1.1955, fr.

56947 RGS, Esteio p. S. Leopoldo, NEL, in humidis, 14.11.1955, fr.

Habit: Culms up to 40 cm high, ears mostly simple.

Habitat: Sandy, more or less wet places.

Distribution in RGS: Northeastern Lowlands and sea coast.

General distribution: From the Antilles to Uruguay.

Paspalum conjugatum Berg., Act. hevet. 7 () 129, tab. 8; Döll, FB 2,2 (1877) 55; Ekman, Ark. f. Bot. 11,n.4 (1912) 11; idem, Ark.f.Bot. 13,n.10 (1913) 11; Reitz, Sellowia 7 (1956) 106.

862 RGS, Vila Manresa p. P. Alegre, NEL, in umbrosis subhumidis, 5.10.1933, fr., leg. C. Orth SJ, det. Chase.

1055 RGS, S. Leopoldo, NEL, in umbrosis, 10.9.1934, fr., leg. C. Orth SJ, det. Beetle.

33054 RGS, Parecí p. Montenegro, NEL, in subhumidis umbrosis, 28.12.1945, fr., leg. A. Strieder SJ.

33533 RGS, S. Leopoldo, NEL, in subhumidis umbrosis, 14.4.1946, fr., leg. E. Henz SJ.

38531 RGS, S. Leopoldo, NEL, in humidis umbrosis, 4.12.1948, fr.

39942 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL in umbrosis, 12.1.1949, fr., det. Swallen.

40136 RGS, Guaíba p. P. Alegre, NEL, in humidis umbrosis, 24.1.1949, fr., det. Swallen.

41068 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in humidis umbrosis 18.4.1949, fl.

41148 RGS, Ilha da Pintada p. P. Alegre, NEL, in humidis umbrosis, 24.4.1949, fr.

41369 RGS, Morretes p. P. Alegre, NEL, in humidis umbrosis, 2.5.1949, fr.

41662 RGS, S. Leopoldo, NEL, in umbrosis subhumidis, 21.5.1949, fl., det. Swallen.

42048 RGS, Sapucaia p. S. leopoldo, NEL, in umbrosis, 18.6.1949, fr.

46473 RGS, Lagoa da Pinguela p. Osório, NEC, in umbrosis subhumidis, 27.3.1950, fr., det. Parodi.

49814 SC, Itapiranga, ad fl. Uruguay superius, ad viam in incultis, 6.2.1951, fr.

50924 Paraná, S. José dos Pinhais, ad flumen in arenosis, 6.1.1950, fr., leg. G. Hatschbach 1705, det. Parodi.

55339 SC, Massiambú p. Palhoça, ad viam in incultis, 12.3.1953, fr., leg. R. Reitz 17193, det. Swallen.

55381 SC, Ascurra p. Rodeio, ad viam, 11.6.1953, leg. R. Reitz 7140, det. Swallen.

55427 SC, Barra do Sul p. Araquari, in dumosis, 8.4.1953, fr., leg. R. Reitz 7061, det. Swallen.

58036 SC, Azambuja p. Brusque, in incultis, 18.11.1953, fl., leg. R. Reitz 8288, det. Swallen.

58051 SC, Caixad'Águap. Itajaí, in incultis, 18.3.1953, fl., leg. R. Reitz 8304, det. Swallen.

Habit: Densely cespitose, culms up to 50 cm long.

Habitat: Always on disturbed soil.

Distribution in RGS: Apparently not native; all over the cultivated area.

General distribution: From Amazonia to Uruguay.

Paspalum conspersum Schrad., in Schult. Mant. 2 () 174; Reitz, Sellowia 7 (1956) 106. — *P. virgatum* L. var. *conspersum* (Schrd.) Döll, FB 2,2 (1877) 89.

41601 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in subpaludosis dumosis, 16.5.1949, fr., det. Swallen.

Habit: Robust, up to 1,5 m high.

Habitat: Wet, shrubby places.

Distribution in RGS: No other findplace.

General distribution: From Minas Gerais to RGS.

Paspalum corcovadense Raddi, Agrost. Bras. (1828) 27; Reitz, Sellowia 7 (1956) 106. — *P. laxum* Lam., Encycl. Illustr. 176 () 931; Döll, FB 2,2 (1877) 85.

1102 RGS, ad montem Steinkopf p. S. Leopoldo, NEL, in umbrosis dumosis, 21.11.1934, fr., leg. C. Orth SJ, det. Chase, Parodi.

40799 RGS, Sapucaia p. S. Leopoldo, NEL, in umbrosis siccis dumosis, 1.4.1949, fr., det. Swallen.

41013 RGS, ad montem das Cabras p. S. Leopoldo, NEL, in siccis dumosis, 8.4.1949, fl. det. Swallen.

41539 RGS, Vila Manresa p. P. Alegre, NEL, in siccis umbrosis, 14.5.1949, fr.

41580 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in siccis dumosis, 16.5.1949, fr., det. Swallen.

41845 RGS, Canoas p. P. Alegre, NEL, in siccis umbrosis, 3.6.1949, fr.

42010 RGS, Morro Santana p. P. Alegre, NEL, in dumosis umbrosis, 16.6.1949, fl.

42495 RGS, Sta Rita p. Farroupilha, NEH, in umbrosis, 13.7.1949, fr.

46402 RGS, Gramado p. Canela, NEH, in umbrosis, 20.3.1950, fr.

46174 RGS, Fazenda do Arroio p. Osório, NEC, in umbrosis, 6.3.1950, fr.

49875 SC, Itapiranga ad fl. Uruguay superius, in silva pluviali ad viam, 6.2.1951, fr.

55416 SC, Laguna, prope litus in silvula, 22.12.1952, fr., leg. R. Reitz 7081, det. Swallen.

59728 RGS, Farroupilha, NEH, in dumosis umbrosis, 28.5.1956, fr., leg. O. Camargo 690.

59729 RGS, ibidem, 28.5.1956, fr., leg. O. Camargo 708.

Habit: Erect, loose bunches, leaves 30, ear 50 cm long.

Habitat: Shady places at forest edge and in secondary growth.

Distribution in RGS: Apparently over the whole of the forested area.

General distribution: From Bahia to RGS.

Paspalum cromyorrhizon Trin., ap. Döll, FB 2,2 (1877) 74;

2734 RGS, S. Leopoldo, NEL, in siccis dumosis, 23.10.1935, fr., leg. C. Orth SJ, det. Chase (Parodi: *P. guaraniticum* Par.).

38822 RGS, Conceição p. Caí, NEL, in campestribus alte graminosis, 14.12.1948, fl. det. Swallen.

43885 RGS, ad fl. Caí inferius p. P. Alegre, NEL, in subsiccis alte dumosis, 12.10.1949, fr., det. Swallen.

56953 RGS, Esteio p. S. Leopoldo, NEL, in humidis alte graminosis, 11.11.1955, fr.

Habit: Thin bunches, culms up to 80 cm high.

Habitat: More or less wet Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: South Brazil.

Paspalum dilatatum Poir., Encycl. Méth. 5 (1804) 35; Döll, FB 2,2 (1877) 64; Ark.f.Bot. 11,n.4 (1912) 11; Herter, Flor. Urug. (1930) 30; Reitz, Sellowia 7 (1956) 106.

25573 RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus dumosis, 1.1944, fr., det. Swallen.

35154 RGS, Fazenda da Ronda p. Vacaria, NEH, in subpaludosis dumosis, 15.1.1947, fr.

36452 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus dense graminosis, 2.1948, fr., det. Swallen.

39378 RGS, Belém Novo p. P. Alegre, NEL, in campestribus dumosis, 31.12.1948, fr.

40232 RGS, Sta Rita p. Farroupilha, NEH, in subpaludosis dumosis, 28.1.1949, fr.

54949 RGS, Vila Oliva p. Caxias, NEH, ad viam in humidis, 24.2.1954, fr., det. Parodi, Swallen.

55064 RGS, ibidem, 24.2.1954, fr., det. Parodi, Swallen.

59526 RGS, Farroupilha, NEH, in alte graminosis, 26.10.1956, fr., leg. O. Camargo 891.

59991 RGS, ibidem, 25.3.1957, fr., leg. O. Camargo 1198.

60405 RGS, ibidem, 26.4.1956, fr., leg. O. Camargo 623.

Habit: Erect, 40 cm high.

Habitat: More or less wet Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: From Pernambuco to Uruguay and B. Aires.

Paspalum distichum L., Amoen. Acad. 5 (1789) 301; Ekman, Ark.f.Bot. 11,n.4 (1912) 16; idem, Ark.f.Bot. 13,n.10 (1913) 19; Malme, Sv. Bot. Tidskr. 30 (1936) 9; Cabrera, Fl. Alred. B. Aires (1953) 102; Reitz, Sellowia 7 (1956) 106. — *P. vaginatum* Swartz, Prodr. () 21; Döll. FB, 2,2 (1877) 74.

2871 RGS, Ilha do Paiva p. P. Alegre, NEL, in paludosis, 26.12.1937, fr., leg. C. Orth SJ, det. Parodi.

3103 SC, Florianópolis, prope litus in arenosis humidis, 6.1938, ster., det. Swallen.

34251 Chile, Padre Las Casas p. Cautín, in pascuo, 4.12.1945, fl., leg. H. Gunckel 15703, det. Parodi.

48343 SC, Figueiredo p. Bom Retiro, ad lacum, 28.12.1948, fl., leg. R. Reitz 3442, det. Swallen.

Habit: Prostrate to ascending, branches up to 50 cm long.

Habitat: Wet to swampy places.

Distribution in RGS: Northeastern Lowlands.

General distribution: Over the whole of tropical and subtropical to temperate South Amerika, and in the Old World.

Paspalum exaltatum Presl, Rel. Haenck. 1 () 219.

54683 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in paludosis alte dumosis, 1.1.1954, fl., fr., det. Parodi (Swallen: *Paspalum* sp.).

56294 RGS, ibidem, 2.1.1955, fr., det. Parodi, Swallen, Barreto: *P. haumanii* Par.

59816 RGS, Farroupilha, NH, in subpaludosis dumosis, 28.12.1956, fl., leg. O. Camargo 1058.

Habit: Culms up to 1,5 m high.

Habitat: Swampy places among high shrub and grasses.

Distribution in RGS: No other findplace.

General distribution: Chile, RGS, Uruguay.

Paspalum guaraniticum Parodi,

2734 RGS, S. Leopoldo, NEL, in subhumidis, 1934, fr., det. Parodi
(Barreto: P. cromyorrhizone Trin.)

56948 RGS, Esteio p. S. Leopoldo, in subhumidis alte graminosis,
14.11.1955, fr.,

56949 RGS, ibidem, 14.11.1955, fr.

Habit: Erect, 75 cm high.

Habitat: Moderately wet places among high grasses.

Distribution in RGS: Northeastern Lowlands.

General distribution: RGS.

Paspalum guenoarum Arech., An. 1 ed. 1 (1894) 55; Herter, Flor. Urug.
(1930) 55.

56627 RGS, Vila Oliva p. Caxias, NEH, in campestribus subhumidis,
8.2.1955, fr., det. Parodi (Swallen, Barreto: P. maculosum Trin.).

60479 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte
graminosis, 8.4.1946, fr., leg. O. Camargo 607.

60481 RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 544.

60485 RGS, ibidem, 8.4.1957, fr., leg. O. Camargo 555.

60488 RGS, ibidem, 8.4.1956, fl., leg. O. Camargo 552.

Habit: Culms up to 100 cm high.

Habitat: Dry to somewhat wet Campos among high grasses.

Distribution in RGS: Widely scattered.

General distribution: Uruguay, RGS.

Paspalum haumanii Parodi, Com. Mus.Nac. B. Aires 2,n.21 (1925) 217;
Herter, Flor. Urug. (1930) 30; Cabrera, Fl. Alred. B. Aires (1953) 102.

1101 RGS, S. Leopoldo, NEL, in paludosis, 8.11.1934, fr., leg. O. Orth
SJ, det. Parodi (Chase: P. durifolium Mez).

8652 RGS, Serra da Rocinha p. Bom Jesus, NEH, in paludosis alte
dumosis, 16.1.1942, fl.

32773 RGS, Parecí p. Montenegro, NEL, in dumosis ad flumen,
10.11.1945, fr., leg. E. Henz SJ.

35176 RGS, Serra da Rocinha p. Bom Jesus, NEH, in humidis
dumosis 14.2.1947, fr., det. Beetle.

35686 RGS, S. Leopoldo, NEL, in paludosis alte dumosis, 22.2.1947,
fr., leg. E. Henz SJ, det. Beetle.

38345 RGS, Esteio p. S. Leopoldo, NEL, in paludosis alte graminosis,
24.11.1948.

49287 RGS, Gravataí p. P. Alegre, NEL, ad rivum in alte dumosis, 1.12.1950, fr.

55001 RGS, Vila Oliva p. Caxias, NEH, in paludosis alte dumosis, 24.2.1954, fr., det. Parodi.

57356 RGS, Estação de Silvicultura p. Sta Maria, CL, in subpaludosis dense dumosis, 18.11.1955, fr., leg. O. Camargo 15.

62004 RGS, Farroupilha, NEH, in humidis dense graminosis, 18.11.1957, fl., leg. O. Camargo.

62965 RGS, Bom Jesus, NEH, in subhumidis, 6.1.1958, fl., leg. O. Camargo 3123.

Habit: Culms up to 2 m high.

Habitat: Swampy places among high shrubs and grasses.

Distribution in RGS: Apparently over the whole of the swampy area.

General distribution: RGS, Uruguay, B. Aires.

Paspalum hyalinum Nees, Agrost. Bras. (1829) 49; Döll, FB 2,2 (1877) 46; Ekman, Ark.f.Bot. 13,n.10 (1913) 11; Reitz, Sellowia 7 (1956) 106.

2522 RGS, Portão p. S. Leopoldo, NEL, in campestribus subhumidis, 12.3.1936, fr., leg. C. Orth SJ, det. Chase, Parodi.

31402 SC, Sombrio, p. Araranguá, in arenosis graminosis, 31.1.1946, fr.

Habit: Small, culms up to 25 cm high, weak and slender.

Habitat: More or less wet Campos among other grasses.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Minas Gerais to RGS.

Paspalum inaequivalve Raddi, Agrost. Bras. (1828) 28; Döll, FB 2,2 (1877) 51; Ekman, Ark. f. Bot. 11,n.4 (1912) 11; Reitz, Sellowia 7 (1956) 107.

1020 RGS, S. Leopoldo, NEL, in incultis, 8.9.1934, fr., leg. C. Orth SJ, det. Chase.

30218 Uruguay, Tacuarembó, 12.3.1945, fr., leg. et det. B. Rosengurtt.

39858 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in umbrosis dumosis, 12.1.1949, fr.

41352 RGS, Morretes p. P. Alegre, NEL, in dumosis, 2.5.1949, fl.

41640 RGS, Esteio p. P. Alegre, NEL, in alte graminosis, 18.5.1949, fr., det. Swallen.

41791 RGS, ibidem, 1.6.1949, fl., det. Swallen.

42298 RGS, Langschneis p. Montenegro, NES, in umbrosis alte dumosis, 2.7.1949, fl., det. Swallen.

Habit: Slender, prostrate to ascending branches up to 50 cm long.

Habitat: Shady, more or less dry places.

Distribution in RGS: Northeastern Lowlands and rain forest belt.

General distribution: From Rio de Janeiro to RGS.

Paspalum juergensii Hack.,

4312 RGS, Taimbésinho p. S. Francisco, NEH, in subpaludosis alte dumosis, 7.2.1941, fr., det. Swallen.

41248 RGS, ad fl. Caí inferius p. P. Alegre, NEL, in subpaludosis alte dumosis, 27.4.1949, fr., det. Swallen.

43864 RGS, ibidem, 12.10.1949, fr., det. Swallen.

45943 RGS, Lagoa dos Quadros p. Torres, NEC, in humidis alte dumosis, 21.2.1950, fr.

46316 RGS, Cristo Rei p. S. Leopoldo, NEL, in dumosis, 17.3.1950, fr.

Habit: Robust, up to 75 cm high.

Habitat: Wet to swampy places among shrubs and other grasses.

Distribution in RGS: Northeastern quadrant.

General distribution: RGS.

Paspalum maculosum Trin., Spec. Gram. (1828) tab. 113; Döll, FB 2,2 (1877) 72; Ekman, Ark.f.Bot. 13,n.10 (1913) 14; Herter, Flor. Urug. (1930) 30; Reitz, Sellowia 7 (1956) 107.

7018 SC, Lages, in campestribus, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

35218 RGS, Serra da Rocinha, p. Bom Jesus, NEH, in campestribus subpaludosis, 14.2.1947, fr.

54888 RGS, Vila Oliva p. Caxias, NEH, in campestribus dense graminosis, 24.2.1954, fr., det. Parodi.

55441 SC, Mafra, in campestribus alte graminosis, 27.1.1953, fr., leg. R. Reitz 7040, det. Swallen.

60480 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fr., leg. O. Camargo. 603.

60484 RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 581.

Habit: Culms slender, up to 60 cm high.

Habitat: Dry Campos among high grasses.

Distribution in RGS: Campos of the northern half of the State.

General distribution: From Piauí to Uruguay.

Paspalum mandiocanum Trin., Dias 2 () 113; Döll, FB 2,2 (1877) 80; Ekman, Ark.f.Bot. 13, n.10 (1913) 15; Malme, Sv. Bot. Tidskr. 30 (1936) 10; Sellowia 7 (1956) 107.

866 RGS, Vila Manresa p. P. Alegre, NEL, in umbrosis dumosis, 10.1.1933, fr., leg. C. Orth SJ, det. Chase.

33027 RGS, Parecí p. Montenegro, NEL, in incultis, 10.10.1945, fr., leg. A. Strieder SJ.

- 38075** RGS, Vila Manresa p. P. Alegre, NEL, in umbrosis dumosis, 17.11.1948, fr., det. Swallen.
- 38161** RGS, Morro da Polícia p. P. Alegre, NEL, in umbrosis, 20.11.1948, fr., det. Swallen.
- 38217** RGS, Sapucaia p. S. Leopoldo, NEL, in umbrosis dumosis, 22.11.1948, fr.
- 38850** RGS, S. Leopoldo, NEL, in umbrosis, 17.12.1948, fr.
- 39806** RGS, Parecí p. Montenegro, NEL, in dumosis secundariis, 14.1.1949, fl., det. Swallen.
- 39998** RGS, Linha Bonita p. Montenegro, NES, in dumosis, 19.1.1949, fr., det. Swallen.
- 40477** RGS, Vila Manresa p. P. Alegre, NEL, ad rivum in dumosis, 14.3.1949, fr., det. Swallen.
- 40682** RGS, ibidem, 26.3.1949, fr., det. Swallen.
- 40851** RGS, Granja Neugebauer p. P. Alegre, NEL, ad silvam in dumosis, 3.4.1949, fr., det. Swallen.
- 40900** RGS, ad montem das Cabras p. S. Leopoldo, NEL, in umbrosis dumosis, 8.4.1949, fr., det. Swallen.
- 40917** RGS, ibidem, 8.4.1949, fr., det. Swallen.
- 41072** RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in humidis umbrosis, 18.4.1949, fr.
- 42073** RGS, Sapucaia p. S. Leopoldo, NEL, in dumosis umbrosis, 18.6.1949, fr.
- 54990** RGS, Vila Oliva p. Caxias, NEH, in humidis alte graminosis, 24.2.1954, fr., det. Parodi.
- 56945** RGS, Esteio p. S. Leopoldo, NEL, in alte dumosis, 15.11.1955, fr.
- 58055** SC, Ibirama, ad silvam, 1.3.1954, fr., leg. R. Reitz 8334, det. Swallen.
- 60478** RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fl., leg. O. Camargo.
- 60607** RGS, Farroupilha, NEH, in dumosis, 7.5.1957, fl., fr.
- 64230** SC, S. Joaquim, in dumosis, 10.1.1958, fl., leg. J. Mattos.
- 64231** SC, ibidem, 10.1.1958, fl., leg. J. Mattos.
- 64232** SC, ibidem, 10.1.1958, fl., leg. J. Mattos.

Habit: Vigorous, prostrate to ascending, rooting at the lower joints, branches up to 50 cm long.

Habitat: Forest edge, secondary growth, shady places.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Minas Gerais to RGS.

Paspalum multiflorum Döll, FB 2,2 (1877) 90; Ekman, Ark.f.Bot. 11, n.4 (1911) 14; idem, Ark.f.Bot. 13, n.10 (1913) 19; Herter, Flor. Urug. (1930) 30; Bornmüller, Rev. Sud. Bot. 1 (1934) 132.

55612 RGS, Fachinal Preto p. Bom Jesus, NEH, in paludosis, 29.1.1950, fr. deciso, det. Asplund. (Barreto: P. quadrifarium Lam.).

Habit: Robust, culms up to 1 m high.

Habitat: Wet thickets among high grasses and shrubs.

Distribution in RGS: Highlands.

General distribution: From S. Paulo to Uruguay.

Paspalum nicorae Parodi,

Reitz, Sellowia 7 (1956) 107.

37999 RGS, Barra do Ribeiro p. P. Alegre, NEL, in arenosis siccis, 14.11.1948, fr., det. Swallen.

39466 RGS, Teresópolis p. P. Alegre, NEL, in campestribus siccis, 4.1.1949, fr.

44424 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis, 19.11.1949, fr.

54339 SC, Laguna, in arenosis, 22.12.1951, fr., leg. R. Reitz 5033, det. Swallen.

54340 SC, ibidem, 10.11.1951, leg. R. Reitz, det. Swallen.

54830 RGS, Torres, NEC, in arenosis, 11.2.1954, fr., det. Parodi, Swallen.

56487 RGS, ibidem, 19.1.1955, fr.,

60483 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fr., leg. O. Camargo 578.

Habit: Cespitose, culms up to 30 cm high.

Habitat: Dry Campos.

Distribution in RGS: Northeastern Lowlands and sea coast.

General distribution: RGS, Uruguay and B. Aires.

Paspalum notatum Flügge, Monogr. () 106; Döll, FB 2,2 (1877) 72; Ekman, Ark. f. Bot. 11, n.4 (1912) 11; idem, Ark. f. Bot. 13, n.10 (1913) 14; Herter, Flor. Urug. (1930) 30; Reitz, Sellowia 7 (1956) 107.

727 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 20.12.1932, fr., det. Chase.

7444 RGS, S. Leopoldo, NEL, in campestribus siccis, 1907, fr., leg. F. Theissen SJ.

9707 RGS, Tupanciretan, NWH, in campestribus siccis, 29.1.1942, fl., det. Chase, Parodi.

11696 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 1943, fr., det. Swallen.

25582 RGS, Fazenda Sta Cecília p. S. Gabriel, in campestribus siccis, 1.1944, fl.

25590 RGS, ibidem, 1.1944, fl.

- 30721** RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis, 28.1.1946, fl.
- 31743** SC, Sombrio p. Araranguá, in arenosis siccis, 7.2.1946, fr., det. Swallen.
- 31952** SC, ibidem, 5.12.1945, fl., leg. R. Reitz 1343.
- 32816** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 12.12.1945, fl.
- 33035** RGS, Parecí p. Montenegro, NEL, in campestribus siccis, 10.10.1945, fl., leg. A. Strieder SJ.
- 35152** RGS, Serra da Rocinha p. Bom Jesus, NEH, in campestribus siccis, 14.2.1947, fr.
- 38527** RGS, S. Leopoldo, NEL, in campestribus siccis, 4.12.1948, fr.
- 39522** RGS, Cachoeirinha p. Gravataí, NEL, in campestribus siccis, 7.1.1949, fl.
- 40134** RGS, Guaíba P. Alegre, NEL, in campestribus siccis graminosis, 24.1.1949, fl.
- 48350** SC, Figueiredo p. Bom Retiro, in campestribus siccis, 28.12.1948, fr., leg. Reitz 3473, det. Swallen.
- 54336** SC, Laguna, prope litus in arenosis, 22.12.1952, leg. R. Reitz 5030, det. Swallen.
- 55083** RGS, Vila Oliva p. Caxias, NEH, in dense graminosis, 24.2.1954, fr.
- 55436** SC, Mafra, in campestribus alte graminosis, 26.1.1953, fl., leg. R. Reitz 7047, det. Swallen.
- 55437** SC, ibidem, 26.12.1953, fl., leg. R. Reitz 7046, det. Swallen.
- 56404** RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in dense graminosis, 2.1.1955, fr.
- 56694** RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis dense graminosis, 8.2.1955, fr.
- 58052** SC, Massiambú p. Palhoça, in arenosis, 4.11.1955, fr., leg. R. Reitz 8329, det. Swallen.
- 58602** RGS, Estação de Silvicultura p. Sta Maria, CL, in graminosis siccis, 10.1.1956, fl., leg. O. Camargo 31.
- 58997** SC, ibidem, 5.1.1956, leg. O. Camargo 141.
- 59856** RGS, Farroupilha, NEH, in pascuo, 18.3.1957, fl., leg. O. Camargo 1169.
- 61639** RGS, Taquarí, in campestribus siccis, 8.12.1957, fl., leg. O. Camargo 2758.
- 64210** SC, S. Joaquim, in campestribus, 10.1.1958, fl., leg. J. Mattos.

Habit: Cespiteous, culms up to 30 cm high.

Habitat: Dry Campos, often dominant.

Distribution in RGS: Common over the whole of the Campos area.

General distribution: From Goiás to Uruguay and B. Aires.

Paspalum orbiculatum Poir. Encycl. Méth. 5 () 32; *P. pusillum* Vent., ap. Döll, FB 2,2 (1877) 45.

1956 RGS, S. Leopoldo, NEL, in humidis, 9.10.1934, fl., leg. C. Orth SJ, det. Chase.

Habit: Prostrate, very small, branches 10 cm long.

Habitat: Wet places.

Distribution in RGS: No other findplace.

General distribution: From the West Indies to RGS.

Paspalum paniculatum L., Spec. Plant. ed. 2 (1762) 81; Döll, FB 2,2 (1877) 56; Ekamn, Ark. f. Bot. 11,n.4 (1912) 11; idem, Ark. f. Bot. 13, n.10 (1913) 11; Malme, Sv. Bot. Tidskr. 30 (1936) 10; Reitz, Sellowia 7 (1956) 107.

1004 RGS, S. Leopoldo, NEL, in incultis, 12.10.1934, fr., leg. C. Orth SJ, det. Chase, Parodi.

10836 RGS, Marcelino Ramos p. fl. Uruguay, NWU, in dumosis, 1.1943, fr., leg. E. Friederichs SJ.

36453 RGS, Cambará p. S. Francisco de Paula, NEH, in dumosis, 2.1948, fr., det. Swallen.

36454 RGS, ibidem, 1.1948, fr., det. Swallen.

40705 RGS, Cristal p. P. Alegre, NEL, in arenosis humidis dumosis, 31.3.1949, fr. det. Swallen.

54992 RGS, Vila Oliva p. Caxias, NEH, in paludosis, 24.2.1954, fr., det. Parodi.

55091 RGS, ibidem, 24.2.1954, fr., det. Parodi (Swallen: *P. juergensii* Hack.).

55392 SC, Anitápolis p. Palhoça, in dumosis, 2.4.1953, fr., leg. R. Reitz 7122, det. Swallen.

56770 RGS, Vila Oliva p. Caxias, NEH, in subhumidis dumosis, fr. det. Parodi (Swallen: *P. umbrosum* Trin; Barreto: *P. juergensii* Hack.).

56813 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in humidis dumosis, 16.2.1955, fr., det. Parodi, (Swallen, Barreto: *P. juergensii* Hack.).

58943 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 1.3.1956, fr., leg. O. Camargo 134.

59235 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in dumosis, 12.2.1956, fr.,

59814 RGS, Farroupilha, NEH, in dumosis, 25.3.1957, fr., leg. O. Camargo 1323.

59815 RGS, ibidem, 25.3.1957, fr., leg. O. Camargo 1320.

59817 RGS, ibidem, 18.3.1957, fr., leg. O. Camargo 1171.

59818 RGS, ibidem, 15.2.1957, fr., leg. O. Camargo 1142.

59819 RGS, ibidem, 25.3.1957, fr., leg. O. Camargo 1315.

- 60269** RGS, *ibidem*, 14.1.1957, fl., fr., leg. O. Camargo 1088.
- 60471** RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fr., leg. O. Camargo 568.
- 60472** RGS, *ibidem*, 8.4.1956, fr., leg. O. Camargo 569.
- 60473** RGS, *ibidem*, 8.4.1956, fr., leg. O. Camargo 576.
- 60474** RGS, *ibidem*, 8.4.1956, fr., leg. O. Camargo 596.
- 60475** RGS, *ibidem*, 8.4.1956, fr., leg. O. Camargo 598.
- 63086** RGS, Pelotas, SEC, in alte graminosis, 11.3.1956, fl., leg. Schlichting.
- 64227** SC, S. Joaquim, in dumosis, 10.1.1958, fl., leg. J. Mattos.
- 64228** SC, *ibidem*, 10.1958, fl., leg. J. Mattos.
- 64229** SC, *ibidem*, 10.1.1958, fl., leg. J. Mattos.

Habit: Erect, slender, up to 75 cm high.

Habitat: Dry to moderately wet brushwood.

Distribution in RGS: Common over the whole of the Campos area.

General distribution: From the West Indies to RGS.

Paspalum plicatum Michx., *Flor. bor. Am.* 1 (1803) 45; Döll, *FB* 2,2 (1877) 76; Ekman, *Ark.f.Bot.* 11,n.4 (1912) 11; idem, *Ark.f.Bot.* 13, n.10 (1913) 14; Herter, *Flor. Urug.* (1930) 30; Malme, *Sv. Bot. Tidskr.* 30 (1936) 10; Cabrera, *Flor. Alred. B. Aires* (1953) 102; Reitz, *Sellowia* 7 (1956) 107.

864 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 5.10.1933, fr., det. Chase.

25593 RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus subdumosis, 1.1944, fl.,

26185 RGS, Fazenda do Jarau p. Quaraí, SW, in campestribus siccis dumosis, 1.1945, fr.

29344 RGS, Vila Manresa p. P. Alegre, NEL, in siccis graminosis, 26.10.1945, fr., det. Parodi.

29375 RGS, *ibidem*, 3.11.1945, fr.

30723 RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis, 28.1.1946, fr., det. Swallen.

30729 RGS, *ibidem*, 28.1.1946, fr.

33213 RGS, Parecí p. Montenegro, NEL, in siccis dumosis, 20.10.1945, fr., leg. E. Henz SJ, det. Swallen.

33576 Uruguay, Punta Ballena p. Maldonado, in siccis dumosis, 14.4.1944, fr., leg. et det. B. Rosengurtt B-4676 (var. *vilosissimum* Pilger).

34157 RGS, Vila Manresa p. P. Alegre, NEL, in graminosis, 16.10.1946, fr.

35155 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis subdumosis, 15.1.1947, fr.

- 35452 RGS, S. Leopoldo, NEL, in arenosis siccis dumosis, 6.11.1946, fr., leg. E. Henz SJ.
- 36451 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus alte graminosis, 2.1948, fr., det. Swallen.
- 37720 RGS, Vila Manresa, p. P. Alegre, NEL, in campestribus siccis graminosis, 22.9.1948, fr., det. Swallen.
- 38455 RGS, ibidem, 1.12.1948, fr., det. Swallen.
- 38664 RGS, Rio dos Sinos p. S. Leopoldo, NEL, in arenosis alte graminosis, 10.12.1948, fr., det. Swallen.
- 38787 RGS, Conceição p. Caí, NEL, in siccis arenosis, 14.12.1948, fr., det. Swallen.
- 38941 RGS, Chácara Weber p. Itapoan, NEL, in siccis graminosis, 18.12.1948, fr.
- 40848 Granja Neugebauer p. Itapoan, NEL, in campestribus alte graminosis, 3.4.1949, fr., det. Swallen.
- 41212 RGS, ad fl. Caí inferius p. P. Alegre, NEL, in siccis arenosis, 26.4.1949, fr., det. Swallen.
- 41491 RGS, Estação Azevedo p. Montenegro, NEL, in campestribus siccis, 6.5.1949, fr., det. Swallen.
- 41807 RGS, Esteio p. S. Leopoldo, NEL, in siccis graminosis, 1.6.1949, fr.
- 43975 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus dumosis, 19.10.1949, fr. det. Swallen.
- 44381 RGS, Fião p. S. Leopoldo, NEL, in arenosis siccis, 14.11.1949, fr.
- 46220 RGS, Cristo Rei p. S. Leopoldo, NEL, in campestribus siccis graminosis, 10.3.1950, fr.
- 46784 RGS, Fazenda do Arroio p. Osório, NEC, in arenosis siccis dumosis, 14.4.1950, fr.
- 54724 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in graminosis, 1.1.1954, fr., det. Parodi (var. glabrum Ar.)
- 55073 RGS, Vila Oliva p. Caxias, NEH, in dense graminosis, 24.2.1954, fr., det. Parodi, Swallen.
- 55440 SC, Mafra, in campestribus dumosis, 16.1.1953, fr., leg. R. Reitz 7042, det. Swallen.
- 56051 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus dumosis, 19.11.1954, fr., det. Parodi, Swallen, Barreto.
- 56413 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in campestribus dense graminosis, 2.1.1955, fr.
- 56950 RGS, Esteio p. S. Leopoldo, NEL, in humidis alte graminosis, 14.11.1955, fr.
- 56951 RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 4.11.1955, fr.

- 56952** RGS, Esteio p. S. Leopoldo, NEL, in humidis alte graminosis, 11.11.1955, fr.
- 57007** RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fr.,
- 57500** RGS, Estação de Silvicultura, Sta Maria, CL, in alte graminosis, 15.11.1955, fl., leg. O. Camargo 24.
- 57545** RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 19.11.1955, fl.
- 57546** RGS, ibidem, 19.11.1955, fr.
- 58358** RGS, Taimbesinho p. S. Francisco de Paula, NEH, in graminosis, 12.2.1956, fr.
- 60486** RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fr., leg. O. Camargo 562.
- 60490** RGS, ibidem, 8.4.1956, fl., leg. O. Camargo 604.
- 60491** RGS, ibidem, 8.4.1957, fr., leg. O. Camargo 609.
- 60492** RGS, ibidem, 8.4.1956, fl., leg. O. Camargo 613
- 61727** RGS, Farroupilha, NEH, in alte graminosis, 18.11.1957, fl., leg. O. Camargo 2580.
- 61745** RGS, Taquarí, NEL, in campestribus, 8.12.1957, fl., leg. O. Camargo 2764.
- 61754** RGS, P. Alegre, NEL, in alte graminosis, 19.11.1957, fl., leg. O. Camargo 2592.

Habit: Culms erect, slender, up to 1 m high.

Habitat: Dry to moderately wet Campos.

Distribution in RGS: In several varieties over the whole of the Campos area.

General distribution: From Amazonia to Uruguay and B. Aires.

Paspalum plumosum

59335 RGS, Estação de Silvicultura p. Sta Maria, CL, in campestribus alte graminosis, fr., 10.1956, fr., leg. O. Camargo 45.

Habit:

Habitat: Dry, grassy Campos.

Distribution in RGS: No other findplace.

General distribution:

Paspalum polyphyllum Nees, Agrost. Bras. (1929) 41; Herter, Flor. Urug. (1930) 31; Reitz, Sellowia 7 (1956) 107. — *Paspalum distichophyllum* H.B.K., Nov. Gen. et Spec. 1 () 86; Döll, FB 2,2 (1877) 65.

1059 RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus dumosis, 8.9.1934, fr., leg. C. Orth SJ, det. Chase.

7017 SC, Lages, in subhumidis dumosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

7034 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

35188 RGS, Serra da Rocinha p. Bom Jesus, NEH, in rupestribus humidis, 14.2.1947, fr.

40511 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus subhumidis, 14.3.1949, fr., det. Swallen.

40690 RGS, ibidem, 26.3.1949, fr., det. Swallen.

40804 RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in rupestribus temporarie humidis, 1.4.1949, fr., det. Swallen.

50901 Paraná, Serra de S. Luiz p. Campo Largo, in rupestribus humidis, 12.3.1950, fr., leg. G. Hatschbach 1903, det. Parodi.

54991 RGS, Vila Oliva p. Caxias, in campestribus dense graminosis, 24.2.1954, fr. det. Parodi.

64343 SC, Lages, in campestribus, 17.2.1958, fl., leg. J. Mattos.

Habit: Erect, culms slender, up to 60 cm high.

Habitat: Slightly wet Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: From Venezuela to RGS.

Paspalum pumilum Nees, Agrost. Bras. (1829) 52; Döll, FB 2,2 (1877) 57; Ekman, Ark. f. Bot. 13, n.10 (1913) 12; Herter, Flor. Urug. (30) 31; Malme, Sv. Bot. Tidskr. 30 (1936) 10; Cabrera, Fl. Alred. B. Aires (1953) 101; Reitz, Sellowia 7 (1956) 107.

9610 RGS, Tupanciretan, NWH, in campestribus siccis, 28.1.1942, fl., det. Swallen.

31434 SC, Sombrio p. Araranguá, in arenosis siccis, 31.1.1946, fr., det. Swallen.

36455 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus dense graminosis, 2.1948, fr., det. Swallen.

36457 RGS, ibidem, 2.1948, fr., det. Swallen.

38518 RGS, S. Leopoldo, NEL, in arenosis siccis, 4.12.1948, fr.

54649 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in campestribus dense graminosis, 1.1.1954, fr., det. Parodi.

55070 RGS, Vila Oliva p. Caxias, NEH, in campestribus alte graminosis, 24.2.1954, fr., det. Swallen.

55412 SC, Serra da Boa Vista p. S. José, in campestribus dense graminosis, 2.2.1953, fr., leg. R. Reitz 7086, det. Swallen.

55415 SC, Palhoça, prope litus in arenosis, 22.12.1952, fr., leg. R. Reitz 7082, det. Swallen.

56765 RGS, Vila Oliva p. Caxias, NEH, in campestribus subhumidis, 8.2.1955, fr., det. Parodi, Swallen, Barreto.

56944 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus, 3.11.1955, fl.

62003 RGS, Farroupilha, NEH, in alte graminosis, 18.11.1957, fl., leg. O. Camargo 2656.

Habit: Erect, up to 60 cm high.

Habitat: Dry, grassy Campos.

Distribution in RGS: From Guyana to Uruguay and B. Aires.

General distribution: Northeastern Lowlands and Highlands.

Paspalum quadrifarium Lam., Illustr. 176 (1791) n.934; Döll, FB 2,2 (1877) 89; Herter, Flor. Urug. (1930) 31; Cabrera, Fl. Alred. B. Aires (1953) 102.

1009 RGS, S. Leopoldo, NEL, in subpaludosis dumosis, 20.9.1934, fr., leg. C. Orth SJ, det. Parodi (Chase: P. ferrugineum Trin.).

41450 RGS, Estação Azevedo p. S. Leopoldo, NEL, in subpaludosis alte dumosis, 6.5.1949, fr., det. Swallen.

46195 RGS, Cristo Rei p. S. Leopoldo, NEL, in humidis alte dumosis, 10.3.1950, fr., det. Swallen.

60477 RGS, Estação de Silvicultura p. Sta Maria, CN, in alte graminosis, 8.4.1956, fr., leg. O. Camargo 543.

62001 RGS, Taquarí, NEL, in alte graminosis, 8.12.1957, fl., leg. O. Camargo 2743.

62002 RGS, ibidem, 8.12.1957, fl., leg. O. Camargo 2704.

Habit: Robust, erect, culms up to 1 m high.

Habitat: More or less wet places, among high grasses and shrubs.

Distribution in RGS: Northeastern Lowlands.

General distribution: From S. Paulo to Uruguay and B. Aires.

Paspalum regnellii Mez,

Ekamn, Ark. f. Bot. 11,n.4 (1912) 14; Reitz, Sellowia 7 (1956) 107.

1056 RGS, Sapucaia p. S. Leopoldo, NEL, in incultis, 20.9.1934, fr., leg. C. Orth SJ, det. Chase (Parodi: P. conspersum Schrad.).

40915 RGS, ad montem das Cabras p. S. Leopoldo, NEL, ad rivum in alte dumosis, 8.4.1949, fl., det. Swallen.

47149 RGS, Butterberg p. Montenegro, NES, ad rivulum in dumosis, 22.5.1950, fr.

55399 SC, Rio Tavares in Insula Florianópolis, in dumosis arenosis, 11.3.1953, fl., leg. R. Reitz 7111, det. Swallen.

Habit: Robust, Culms up to 1,5 m high.

Habitat: Wet places, streamside.

Distribution in RGS: Northeastern Lowlands and rain forest belt.

General distribution: Misiones, RGS,

Paspalum repens Gerg., Ac. helv. 7 (1762) tab. 7; Döll, FB 2,2 (1877) 98; Herter Flor. Urug. (1930) 31; Cabrera, Fl. Alred. B. Aires (1953) 99.
2738 RGS, S. Leopoldo, NEL, in palude fluitans, 29.5.1935, fr., leg. C. Orth SJ, det. Chase, Parodi.

41084 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in palude fluitans, 18.4.1949, fr., det. Swallen.

55414 SC, Massiambú p. Palhoça, in arenosis, 5.2.1953, fr., leg. R. Reitz 7084, det. Swallen.

Habit: Robust, branches floating, up to several m long.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Guyana to Uruguay and B. aires.

Paspalum rojasii Hack.,

Reitz, Sellowia 7 (1956) 107.

7023 SC, Lages, in dumosis, 1935, fr., leg. A. Bruxel SJ. det. Swallen.

7025 SC, ibidem, leg. A. Bruxel SJ.

46098 RGS, Vila Manresa p. P. Alegre, NEL, in alte dumosis, 3.3.1950, fr.

Habit: Slender, culms up to 1 m high.

Habitat: Dry, shrubby, Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: Paraná to RGS.

Paspalum umbrosum Trin., Act. Petrop. () 153; *P. densiflorum* Döll FB 2,2 (1877) 52.

39133 RGS, Espírito Santo p. P. Alegre, NEL, in umbrosis dumosis, 24.12.1948, fr., det. Swallen.

41453 RGS, Estação Azevedo p. Montenegro, NEL, in humidis dumosis, 6.3.1949, fr., det. Swallen. (Barreto: *P. juergensii* Hack.).

Habit: Slender, 60 cm high.

Habitat: More or less wet, shady places.

Distribution in RGS: Northeastern Lowlands.

General distribution: From the Antilles to RGS.

Paspalum urvillei Steud., Syn. Gram. (1855) 24; Herter, Flor. Urug. (1930) 31; Cabrera, Fl. Alred. B. Aires (1953) 102; Reitz, Sellowia 7 (1956) 107.

868 RGS, Vila Manresa p. P. Alegre, NEL, in humidis subpaludosis, 5.12.1932, fr., det. Chase, Parodi.

7010 SC, Florianópolis, in humidis dumosis, 1935, fr., leg. A. Bruxel, SJ, det. Swallen.

7026 SC, ibidem, 1935, leg. A. Bruxel SJ, det. Swallen.

- 25602 RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in humidis alte dumosis, 1.1944, fr.
- 29384 RGS, Vila Manresa p. P. Alegre, NEL, in humidis alte dumosis, 3.11.1945, fr.
- 38218 RGS, Sapucaia p. S. Leopoldo, NEL, in paludosis alte dumosis, 22.11.1948, fr., det. Swallen.
- 38448 RGS, ibidem, 1.12.1948, fr.
- 40224 RGS, Sta Rita p. Farroupilha, NEH, in humidis alte dumosis, 28.1.1949, fr.
- 40964 RGS, ad fl. Gravataí, inferius p. P. Alegre, NEL, in humidis graminosis, 10.4.1949, fr., det. Swallen.
- 41218 RGS, ad fl. Caí inferius p. P. Alegre, NEL, in humidis alte graminosis, 22.4.1949, fr., det. Swallen.
- 44119 RGS, Barreto Vianna p. P. alegre, NEL, in paludosis alte graminosis, 24.10.1949, fl., fr.
- 45128 RGS, Fazenda do Arroio p. Osório, in humidis alte dumosis, 4.1.1950, fl., fr.
- 49254 RGS, Parecí p. Montenegro, NEL, in paludosis dumosis, 26.11.1950, fr.
- 55058 RGS, Vila Oliva p. Caxias, NEH, in humidis alte graminosis, 24.2.1954, fr., det. Parodi.
- 55374 SC, Rio Tavares in Insula Sta Catarina, in humidis alte graminosis, 23.12.1952, fr., leg. R. Reitz 7152, det. Swallen.
- 56049 RGS, Vila Manresa p. P. Alegre, NEL, in dumosis subhumidis, 15.11.1954, fr., det. Parodi.
- 56672 RGS, Vila Oliva p. Caxias, NWH, in dumosis, 8.2.1955, fr., det. Parodi.
- 56788 RGS, ibidem, 8.1.1955, fr., det. Parodi.
- 56946 RGS, Esteio p. S. Leopoldo, NEL, in humidis alte graminosis, 14.11.1955, fl.
- 59362 RGS, Farroupilha, NEH, in humidis graminosis, 15.11.1956, fl., leg. O. Camargo 963.
- 59391 RGS, ibidem, 25.11.1956, fl., leg. O. Camargo 1010.
- 60016 RGS, ibidem, 25.3.1957, fr., leg. O. Camargo 1322.
- 60482 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fr., leg. O. Camargo 567.
- 60487 RGS, Estação de Silvicultura p. Sta Maria, CN, in alte graminosis, 8.4.1956, fl., leg. O. Camargo.
- 60489 RGS, ibidem, 8.4.1956, fl., leg. O. Camargo 550.
- 61556 RGS, Farroupilha, NEH, in late graminosis, 22.11.1957, fl., leg. O. Camargo 2675.
- 62927 RGS, Taquarí, NEL, in campestribus, 10.12.1957, fl., leg. O. Camargo 3008.
- 63087 RGS, Pelotas, SEC, in in cultis, 11.3.1956, fl., leg. Schlichting.

64212 SC, S. Joaquim, in incultis, 10.1.1958, fl., leg. Mattos.

Habit: Robust, erect, in thick bunches, culms up to 1 m high.

Habitat: Dry to moderately wet places, often on disturbed soil.

Distribution in RGS: Over the whole of the Campos and cultivated area.

General distribution: From SC to Uruguay and B. Aires.

Paspalum vaginatum Swartz, Nov. Gen. et Spec. (1788) 21; Döll, FB 2,2 (1877) 74; Cabrera, Fl. Alred. B. Aires (1953) 102; Reitz, Sellowia 7 (1956) 107.

54337 SC, Laguna, prope litus in arenosis, 22.12.1951, fr., leg. R. Reitz 5031, det. Swallen.

54338 SC, ibidem, 22.12.1951, leg. R. Reitz 5032, det. Swallen.

54349 SC, ibidem, 5.6.1952, fr., leg. R. Reitz 5049, det. Swallen.

54350 SC, ibidem, in humidis salsuginosis, 28.2.1952, fr., leg. R. Reitz 5051, det. Swallen.

54774 RGS, Torres, NEC, prope litus in arenosis humidis, 11.2.1954, fr., det. Parodi.

55371 SC, Rio Tavares in Insula Sta Catarina, in humidis salsuginosis, 23.12.1952, fr., leg. R. Reitz 7155, det. Swallen.

55428 SC, ibidem, 23.12.1952, fr., leg. R. Reitz 7059, det. Swallen.

56454 RGS, Cidreira, NEC, in arenosis humidis, 17.12.1954, fr.

56852 SC, Arroio do Silva p. Araranguá, in arenosis late serpens, 19.2.1955, fr., 1.

58031 SC, Rio Tavares in Insula Sta Catarina, in arenosis, 5.11.1953, fr. det. Swallen.

Habit: Extense stands, deeply rooting in the sand, culms up to 30 cm high.

Habitat: Wet to salty sands near the sea coast.

Distribution in RGS: Sea coast only.

General distribution: From USA to B. Aires and Chile.

Paspalum yaguaronense Henr.,

Reitz, Sellowia 7 (1956) 108.

1936 RGS, p. Bom Jesus, NEH, in campestribus dumosis, 2.2.1934, fr., leg. J. Dutra, det. Swallen.

7000 SC, Lages, in dumosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

35157 RGS, Fazenda da Ronda p. Vacaria, NEH, in dumosis, 15.1.1947, fr.

39738 RGS, Pareci p. Montenegro, in dumosis, 14.1.1949, fr.

55359 SC, Massiambú p. Palhoça, in arenosis, 5.2.1953, fr., leg. R. Reitz 7167, det. Swallen.

60468 RGS, Sta Maria, CL, in campestribus alte graminosis, 8.4.1956, fr., leg. O. Camargo 566.

60469 RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 546.

60470 RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 545.

Habit: Robust, Culms up to 75 cm high.

Habitat: In more or less dry, shrubby, Campos.

Distribution in RGS: Widely scattered.

General distribution: From Paraná to Uruguay.

PENNISETUM L.C.Rich.

Pennisetum latifolium Spreng., Syst. 1 (1825) 302; Ekman, Ark. f. Bot. 13, n.10 (1913) 35; Herter, Flor. Urug. (1930) 31; Cabrera, Fl. Alred. B. Aires (1953) 108; Reitz, Sellowia 7 (1956) 108. — *Gymnothrix tristachya* H.B.K., Nov. Gen. et Spec. 1 () 113; Döll, FB 2,2 (1877) 303, tab. 61.

1068 RGS, Sapucaia p. S. Leopoldo, NEL, in dumosis ad rivum, 5.10.1934, fl., leg. C. Orth SJ, det. Parodi.

7013 SC, Lages, in dumosis ad viam, fl., leg. A. Bruxel SJ, det. Swallen.

29959 RGS, Kappesberg p. Montenegro, NES, ad rivum in alte dumosis, 25.3.1945, fl., leg. E. Friderichs SJ, det. Swallen.

31269 RGS, Vila Oliva p. Caxias, NEH, in dumosis, 21.2.1946, fl. fr.

33191 RGS, ibidem, 16.1.1946, fr.,

33292 RGS, Caracol p. Canela, NEH, ad viam in silva, 26.2.1946, fr., leg. K. Emrich.

59779 RGS, Farroupilha, NEH, in dumosis, 25.3.1957, fl., leg. O. Camargo 1317.

59780 RGS, ibidem, 15.3.1957, fl., leg. O. Camargo 1155.

60374 RGS, Estação de Silvicultura p. Sta Maria, CL, in dumosis, 8.3.1956, fr., leg. O. Camargo 595.

Habit: Culms up to 1,5 m high.

Habitat: More or less shrubby places among high grasses.

Distribution in RGS: Apparently over the whole of the area.

General distribution: From Mexico to Uruguay and B. Aires.

Pennisetum purpureum Schuman, Schum. et Thonn,m Beskr. () tab. 44.

882 RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 5.1.1933, fr., leg. C. Orth SJ, det. A. Chase.

25487 RGS, S. Leopoldo, NEL, in incultis, 1943, fr., leg. R. Reitz.

28989 RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 15.8.1945, fl., det. Swallen.

58766 RGS, Estação de Silvicultura p. Sta Maria, CL, culta, 1.3.1956, ster., leg. O. Camargo 101.

Habit: Large bunches, culms up to 3 m high.

Habitat: Cultivated and subspontaneous in secondary growth.

Distribution in RGS: Over the whole of the cultivated area.

General distribution: Native to tropical Africa, widely scattered in South America.

PHALARIS L.

Phalaris angusta Nees, Agrost. Bras. (1829) 391; Döll, FB 2,2 (1871) 28, tab. 9; Ekman, Ark. f. Bot. 11,n.4 (1912) 29; Herter, Flor, Urug. (1930) 32; Cabrera, Fl. Alred. B. Aires (1953) 89;

726 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 5.10.1932, fr., leg. C. Orth SJ, det. Chase.

2902 RGS, S. Borja, SW, in incultis, 1943, fr., leg. Fr. Baglione.

2903 RGS, ibidem, fr., leg. Fr. Baglione.

6652 RGS, sine die, sine loco, fr., det. Swallen.

10843 RGS, Sta Maria, CL, in incultis, 1943, fr., leg. A. Heidler SJ.

26647 RGS, Parecí p. Montenegro, NEL, in incultis, 1944, fr., leg. E. Henz SJ.

26714 RGS, Cérro Largo p. S. Luiz, NWH, in incultis, 8.1944, fr., det. E. Friderichs SJ.

32872 RGS, Kappesberg p. Montenegro, NES, in incultis, 24.10.1945, leg. E. Friderichs SJ.

33212 RGS, Parecí p. Montenegro, NEL, in campestribus graminosis, 22.10.1945, fr., leg. E. Henz SJ.

33983 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 10.9.1946, fr.

34334 RGS, Cerro Largo p. S. Luiz, NWH, in incultis, 31.10.1946, fl., leg. Irm. Augusto.

35525 RGS, S. Leopoldo, NEL, in incultis, 6.11.1946, fl., leg. E. Henz SJ.

37987 RGS, Barra do Ribeiro p. P. Alegre, NEL, in arenosis, 14.11.1948, fr., det. Swallen.

43738 RGS, Morro do Sabiá p. P. Alegre, NEL, ad viam in incultis, 5.10.1949, fr., det. Swallen.

54453 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 12.11.1953, fr.

56007 RGS, ibidem, in incultis, 19.11.1954, fr.

57227 RGS, ibidem, 15.10.1955, fl.

59627 RGS, Farroupilha, NEH, in incultis, 26.10.1956, fl., leg. O. Camargo 867.

- 60866** RGS, *ibidem*, in incultis, 8.7.1957, fl., leg. O. Camargo 1641.
61856 RGS, Garibaldi, NEH, in incultis, 29.10.1957, fl., leg. O. Camargo 2281.
62211 RGS, Farroupilha, NEH, in incultis, 10.10.1957, fl., leg. O. Camargo 1910.
63379 RGS, Pelotas, SEC, in incultis, 19.10.1954, fl., leg. J. Sacco 206, det. Swallen.

Habit: Culms up to 1 m high.

Habitat: Disturbed soil, often dominant.

Distribution in RGS: Frequent over the whole of the cultivated area.

General distribution: South Brazil, Uruguay, Argentina.

PHARUS L.

Pharus glaber H.B.K., Nov. Gen. et Spec. 1 (1815) 196; Döll, FB 2,2 (1871) 21, tab. 6; Ekman, Ark. f. Bot. 11,n.4 (1912) 28; idem, Ark. f. Bot. 13,n.10 (1913) 36; Herter, Flor. Urug. (1930) 32; Reitz, Sellowia 7 (1956) 108.

725 RGS, Vila Manresa p. P. Alegre, NEL, in silva pluviali, 5.1.1933, fr., leg. C. Orth SJ, det. Chase, Parodi. (20)

7454 RGS, S. Leopoldo, NEL, in silva campestri, 1907, fr., leg. F. Theissen SJ.

10840 RGS, Cerro Largo p. S. Luiz, NWH, in silva pluviali, 1.1943, fl., leg. P. Buck SJ.

29436 RGS, Vila Manresa p. P. Alegre, NEL, in silva pluviali, 15.11.1945, fr. (20).

33029 RGS, Pareci p. Montenegro, NEL, in silva pluviali, 15.1.1946, leg. A. Strieder SJ.

33245 RGS, *ibidem*, 10.12.1945, fr., leg. E. Henz SJ.

36083 RGS, Campinas p. Sta Rosa, NEH, in silva pluviali, 2.1947, leg. A. Spies.

39185 RGS, Morro da Polícia p. P. Alegre, NEL, in silva pluviali, 27.12.1948, fl.

40018 RGS, Linha Bonita p. Montenegro, NES, in silva pluviali, 19.1.1949, fl.

40104 RGS, Vila Manresa p. P. Alegre, NEL, in silva pluviali, 22.1.1949, fr.

41709 RGS, Schwabenschneis p. S. Leopoldo, NES, in silva pluviali, 25.5.1949, fr.

42151 RGS, *ibidem*, 25.6.1949, fr.

- 45901** RGS, Lagoa dos Quadros p. Torres, NEC, in silva pluviali, 21.2.1950, fr.
- 46099** RGS, Vila Manresa, p. P. Alegre, NEL, in silva pluviali, 3.3.1950, fl.
- 49942** SC, Itapiranga p. fl. Uruguay superius, in silva pluviali, 6.2.1951, fr.
- 51143** Paraná, Astorga, in silva pluviali, 29.1.1950, fr., leg. G. Hatschbach 1830, det. Parodi (20).
- 53524** Paraná, Cascavel, in silva pluviali, 10.1.1953, fr. deciso (20).
- 53688** SC, Itapiranga p. fl. Uruguay superius, in silva pluviali, 17.1.1953, fr.
- 53714** SC, ibidem, 17.1.1953, fr.
- 58687** RGS, Estação de Silvicultura p. Sta Maria, CL, in silva, 28.2.1956, fr., leg. O. Camargo 4.
- 63371** RGS, Pelotas, SEC, in silva, 11.3.1955, fl., leg. J. Sacco 287.

Habit: Culms up to 75 cm high.

Habitat: Dark rain forest.

Distribution in RGS: Through the whole of the forested area.

General distribution: From the Antilles to Uruguay.

PIPTOCHAETIUM Presl

Piptochaetium confusum Parodi,

- 29374** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis graminosis, 3.11.1945, fr., det. Swallen.

Habit: Dense bunches, up to 40 cm high.

Habitat: Dry Campos, among other bunchgrasses.

Distribution in RGS: Northeastern Lowlands.

General distribution: RGS.

Piptochaetium lasianthum Gris., Symb. (1879) 297; Herter, Flor. Urug. (1930) 34; Cabrera, Fl. Alred. B. Aires (1953) 77.

- 33083** Uruguay, Juan Jackson p. Soriano, im campestribus alte graminosis, 12.1942, fr., leg. B. Rosengurtt PE-5124, det. Chase.

37868 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus alte graminosis, 6.11.1948, fr., det. Swallen.

43713 RGS, Morro Sta Teresa p. P. Alegre, NEL, in campestribus siccis alte graminosis.

Habit: Bunches up to 1 m high.

Habitat: Dry, grassy Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: Northeastern Argentina, RGS, Uruguay.

- Piptochaetium montevidense** (Spreng.) Parodi, Rev. Fac. Agron. Vet. B. Aires, 7 (1930) 78; Herter, Flor. Urug. (1930) 34; Cabrera, Fl. Alred. B. Aires (1953) 78. — *Urachne panicoides* Trin., Act. Petrop. 1849, 23; Döll, FB 2,2 (1878) 3. — *Caryochloa montevidenses* Spreng., Syst. Veg. Cur. Post. (1827) 31; Ekman, Ark. f. Bot. 13,n.10 (1913) 40.
- 724 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 5.10.1932, fr. leg. C. Orth SJ, det. Parodi.
- 26819 RGS, P. Alegre, NEL, in campestribus siccis, 11.1944, fr., leg. P. Capparelli.
- 26991 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis, 10.1944, fr.
- 29162 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus dumosis, 12.9.1945, fr., det. Swallen.
- 29253 RGS, ibidem, 10.10.1945, fr., det. Swallen.
- 29373 RGS, ibidem, 3.11.1945, fr., det. Swallen.
- 32540 RGS, Pareci p. Montenegro, NEL, in campestribus siccis, 31.10.1945, fr., leg. E. Henz SJ, det. Swallen.
- 32572 RGS, ibidem, 30.10.1945, fr., leg. E. Henz SJ.
- 32758 RGS, ibidem, 17.10.1945, fr., det. Henz SJ, det. Swallen.
- 33888 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 25.9.1946, fr.
- 34202 RGS, ibidem, 16.10.1946, fr.
- 35367 RGS, S. Leopoldo, NEL, in campestribus siccis, 6.11.1946, fr., leg. E. Henz SJ.
- 35399 RGS, ibidem, 2.10.1946, fr., leg. E. Henz SJ.
- 36437 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus siccis, 2.1948, det. Swallen.
- 37850 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis, 4.11.1948, fr., det. Swallen.
- 37855 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 4.11.1948, fr., det. Swallen.
- 43601 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis, 28.9.1949, fr., det. Swallen.
- 43645 RGS, Morro das Abelhas p. P. Alegre, NEL, in campestribus siccis, 30.9.1949, fl.
- 48821 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 22.9.1950, fr.
- 52270 Paraná, Timoneira, in campestribus siccis, 26.11.1950, fr., leg. G. Hatschbach 2035, det. Parodi.
- 53774 RGS, Pareci p. Montenegro, NEL, in campestribus siccis dense graminosis, 7.10.1949, fr.
- 54421 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 12.11.1953, fl., det. Swallen.

56047 RGS, ibidem, 19.11.1954, fr.

57014 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fr.

57207 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 22.10.1955, fr.

57225 RGS, ibidem, 15.10.1955, fr.

57353 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 15.11.1955, fr. leg. O. Camargo 23.

59540 RGS, Farroupilha, NEH, in siccis graminosis, 22.10.1956, fr., leg. O. Camargo 843.

61842 RGS, Garibaldi, NEH, in alte graminosis, 29.10.1957, fl., fr., leg. O. Camargo 2257.

62219 RGS, Bento Gonçalves, NEH, in dense graminosis, 6.10.1957, fr., leg. O. Camargo 1932.

62269 RGS, P. Alegre, NEL, in campestribus, 1.10.1957, fr., leg. O. Camargo 1836.

62495 RGS, Passo Fundo, CH, in campestribus, 20.10.1957, fr., leg. O. Camargo 2139.

62770 RGS, Farroupilha, NEH, in graminosis, 21.10.1957, fl., leg. O. Camargo 2342.

63386 RGS, Pelotas, SEC, in campestribus, 5.10.1954, fl., leg. J. Sacco 78, det. Parodi.

Habit: Dense bunches, up to 40 cm high.

Habitat: Dry, grassy, Campos.

Distribution in RGS: Over the whole of the Campos area.

General distribution: Paraguay, South Brazil, Uruguay, Argentina, Chile.

Piptochaetium panicoides (Lam.) Desv., in Gay, Fl. Chil. 6 (1853) 270;

Herter, Flor. Urug. (1930) 34; Cabrera, Fl. Alred. B. Aires (1953) 78.

– *Stipa panicoides* Lam., Illustr. Gen. 1 (1791) 158.

63385 RGS, Pelotas, SEC, in campestribus, 5.10.1954, fl., fr., det. Swallen.

Habit: Up to 40 cm high.

Habitat: Dry Campos.

Distribution in RGS: Only findplace.

General distribution: RGS, Uruguay, B. Aires.

Piptochaetium ruprechtianum Desv.,

38117 RGS, Morro da Polícia p. P. Alegre, NEL, in alte graminosis, 20.11.1948, fr., det. Swallen.

38391 RGS, Sapucaia p. S. Leopoldo, NEL, in campestribus siccis alte dumosis, 29.11.1948, fr.

43574 RGS, Portão p. S. Leopoldo, NEL, in campestribus alte dumosis, 23.9.1949, fr., det. Swallen.

43652 RGS, Morro das Abertas p. P. Alegre, NEL, in campestribus alte graminosis, 30.9.1949, fr.

44038 RGS, Morro do Osso p. P. Alegre, NEL, in campestribus siccis alte graminosis, 21.10.1949, fr., det. Swallen.

44131 RGS, Barreto Vianna p. S. Leopoldo, NEL, in campestribus siccis alte dumosis, 24.10.1949, fr., leg. Swallen.

Habit: Bunches up to 50 cm high.

Habitat: Dry, densely grassy Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: RGS, Uruguay, B. Aires.

Piptochaetium stipoides (Trin. et Rupr.) Hack., ap. Arech. Gram. Urug. (1896) 268; Herter, Flor. Urug. (1930) 34; Cabrera, Fl. Alred. B. Aires (1953) 77. — *Urachne stipoides* Trin. et Rupr., Act. Ac. St. Petersb., ser. 6, 5 (1842) 25; Döll, FB, 2, 3 (1878) 5.

43697 RGS, Morro de Sta Teresa p. P. Alegre, NEL, in campestribus graminosis, 3.10.1949, fr., det. Swallen.

Habit: Bunches up to 50 cm high.

Habitat: Dry, densely grassy Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: South Brazil, Uruguay, B. Aires.

POA L.

Poa annua L., Spec. Plant. (1753) 68: Döll, FB 2,3 (1878) 121; Herter, Flor. Urug. (1930) 37; Cabrera, Fl. Alred. B. Aires (1953) 65; Reitz, Sellowia 7 (1956) 108.

1005 RGS, S. Leopoldo, NEL, in incultis, 12.10.1934, fr., leg. C. Orth SJ, det. Chase.

26818 RGS, P. Alegre, NEL, in incultis, 9.1944, fr., leg. P. Capparelli.

29049 RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 25.8.1945, fr., det. Parodi.

29621 RGS, Parecí p. Montenegro, NEL, in agro, 19.9.1945, fr.

30520 SC, Jacinto Machado p. Araranguá, in pascuo, 27.10.1943, fr., leg. R. Reitz 944.

33922 RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 18.9.1946, fr.

43086 RGS, Linha Bonita p. Montenegro, NEL, in agro, 24.8.1949, fr.

56855 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in incultis ad viam, 2.1.1955, fl.

57142 RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 4.11.1955, fl.

59654 RGS, Farroupilha, NEH, in incultis, 30.8.1956, fl., leg. O. Camargo 723.

60075 SC, Campo dos Padres p. Bom Retiro, in incultis, 22.1.1957, fr.

61193 SC, Itapiranga ad fl. Uruguay superius, in incultis, 7.10.1957, fl.

62272 RGS, Farroupilha, NEH, in incultis, 12.9.1957, fl., leg. O. Camargo, 1680.

62394 RGS, ibidem, 12.9.1957, fl., leg. O. Camargo 1707.

62404 RGS, ibidem, 12.9.1957, fl., leg. O. Camargo 1698.

62549 RGS, Garibaldi, NEH, in incultis, 13.10.1957, fl., leg. O. Camargo 2062.

Habit: Prostrate to ascending, often densely matted.

Habitat: Disturbed soil only.

Distribution in RGS: Over the whole of the cultivated area.

General distribution: Native to Europe, widely scattered over America.

Poa bonariensis (Lam.) Kunth, Rev. Gram. 1 (1829) 115; Herter, Flor. Urug. (1930) 37; Cabrera, Fl. Alred. B. aires (1953) 65. — *Festuca bonariensis* Lam., Ill. Gen. 1 (1791) 192.

873 RGS, Vila Manresa p. P. alegre, NEL, in campestribus saxosis, 5.10.1933, fr., leg. C. Orth SJ, det. Chase.

Habit: Bunchy, erect, up to 50 cm high.

Habitat: More or less wet, grassy, places.

Distribution in RGS: No other findplace.

General distribution: RGS, Uruguay, B. Aires.

Poa bradei Pilger,

56414 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in subhumidis graminosis, 2.1.1955, fl., det. Parodi.

60045 SC, Campo dos Padres p. Bom Retiro, in humidis graminosis, 24.1.1957, fl.

60553 SC, S. Joaquim, in campestribus alte graminosis, leg. J. Mattos.

Habit: Slender, 50 cm high.

Habitat: Wet, grassy places.

Distribution in RGS: Northeastern Highlands.

General distribution SC, RGS.

Poa lanigera Nees, Agrost. Bras. (1829) 491; Döll, FB 2,3 (1878) 120; Herter, Flor. Urug. (1930) 38; Cabrera, Fl. Alred. B. Aires (1953) 65.

34143 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 15.1.1947, fr.

- 37710** RGS, Vila Manresa p. P. Alegre, NEH, in rupestribus alte graminosis, 22.9.1948, fr.
- 37759** RGS, ibidem, 1.10.1948, fr.
- 37761** RGS, ibidem, 1.10.1948, fr.
- 43489** RGS, ibidem, 20.9.1949, fr., det. Swallen.
- 43607** RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis, 28.9.1949, fr.
- 43617** RGS, ibidem, 28.9.1949, fr.
- 43642** RGS, Morro das Abertas p. P. Alegre, NEL, in campestribus alte graminosis, 30.9.1949, fr.
- 48743** RGS, Osório, NEC, in arenosis subsiccis, 11.9.1950, fr.
- 48826** RGS, Vila Manresa p. P. Alegre, NEL, in saxosis alte graminosis, 22.9.1950, fr.
- 48988** RGS, ibidem, 18.10.1950, fr.
- 59502** RGS, Farroupilha, NEH, in alte graminosis, 26.10.1956, fl., leg. O. Camargo 883 (21).

Habit: Bunchy, erect, up to 50 cm high.

Habitat: More or less dry Campos.

Distribution in RGS: Northeastern quadrant.

General distribution: South Brazil, Uruguay, B. Aires.

Poa montevidensis Arech.,

Herter, Flor. Urug. (1930) 38.

- 2724** RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus saxosis, 2.10.1935, fr., leg. C. Orth SJ, det. Chase.
- 29198** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus saxosis, 12.9.1945, fr.
- 34143** RGS, ibidem, 16.10.1945, fr.
- 34204** RGS, ibidem, 16.10.1946, fr.
- 34444** RGS, Taquarí, NEL, in incultis, 7.11.1946, fr., leg. P. Capparelli.
- 56954** RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 24.9.1955, fl.
- 57145** RGS, ibidem, 18.9.1955, fl.
- 57146** RGS, ibidem, 24.9.1955, fl.

Habit: Bunchy, up to 50 cm high.

Habitat: Dry, stony, Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: RGS, Uruguay.

Poa trivialis L., Spec. Plant. (1753) 67; Cabrera, Fl. Alred. B. Aires (1953) 65.

(21) = **Poa trivialis**, det. H. Longhi, 04.01.1974.

62332 RGS, Montenegro, NEL, in tritis, 19.9.1957, fl., leg. O. Camargo 1782.

Habit: Prostrate to ascending.

Habitat: Disturbed soil.

Distribution in RGS: Cultivated area, rare.

General distribution: RGS, Uruguay, B. Aires.

POLYPOGON Desf.

Polypogon elongatus H.B.K., Nov. Gen. et Spec. 1 (1815) 134; Döll, FB 2,3 (1878) 43, tab. 11; Herter, Flor. Urug. (1930) 33; Cabrera, Fl. Alr. B. Aires (1953) 78.

874 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte dumosis, 5.10.1933, fl., C. Orth SJ, det. Chase, Parodi, Swallen.

28387 RGS, ibidem, 3.11.1945, fr., det. Swallen.

32557 RGS, Pareci p. Montenegro, NEL, in incultis, 22.10.1945, fl., leg. E. Henz SJ.

35164 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus alte dumosis, 15.1.1947, fl.,

38481 RGS, Vila Manresa p. P. Alegre, NEL, in humidis alte dumosis, 1.12.1948, fr., det. Swallen.

49216 RGS, Pareci p. Montenegro, NEL, ad viam ferream in dumosis, 26.11.1950, fl.

59209 RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 22.1.1956, fl.

59734 RGS, Farroupilha, in dumosis, 28.5.1956, fr., leg. O. Camargo 735.

Habit: Erect, Slender, up to 1,5 m high.

Habitat: Shrubby, more or less wet places.

Distribution in RGS: Northeastern quadrant.

General distribution: From Minas Gerais to Uruguay and B. Aires.

Polypogon monspeliensis (Döll) Desf., Fl. Atlant. 1 (1798) 67; Herter, Flor. Urug. (1930) 33; Cabrera, Fl. Alred. B. Aires (1953) 78. — *Alopecurus monspeliensis* L., Spec. Plant. (1753) 61.

56470 RGS, Torres, NEC, ad ipsum mare in rupestribus, 19.1.1959, fr., det. Parodi (Swallen: *P. rioplatensis* Hert.).

Habit: Low, cespitose.

Habitat: Rocky places, in the surf.

Distribution in RGS: Northeastern sea coast.

General distribution: Native to Europa; RGS, Uruguay, B. Aires.

PSEUDECHINOLAENA Stapf

Pseudechinolaena polystachya (H.B.K.) Stapf, in Prain, Fl. Trop. Afr. 9 (1919) 495; Cabrera, Fl. Alred. B. Aires (1953) 105; Reitz, Sellowia 7 (1956) 108. — *Echinolaena polystachya* H.B.K., Nov. Gen. et Spec. 1 (1816) 119.

723 RGS, Vila Manresa p. P. Alegre, NEL, in silva, 5.1.1933, fl., det. Chase.

4344 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in araucarieto, 7.1.1941, ster., det. Swallen.

7023 SC, Lages, in silvula campestri, 1935, ster., leg. A. Bruxel SJ.

7037 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

7743 RGS, S. Leopoldo, NEL, in silva campestri, 1907, fr., leg. F. Theissen SJ.

9188 RGS, Jarí p. Tupanciretan, NWH, in silva campestri, 26.1.1942, fr.

9594 RGS, ibidem, 28.1.1942, fr.

28634 RGS, Nonoai ad fl. Uruguay, NWU, in silva campestri, 3.1945, fl.

29331 RGS, Vila Manresa p. P. Alegre, NEL, in silva campestri, 26.10.1945, fl., det. Parodi.

30738 RGS, Vila Oliva p. Caxias, NEH, in araucarieto, 28.1.1946, fr., det. Swallen.

39578 RGS, Cachoeirinha p. Gravataí, NEL, in silva campestri, 7.1.1949, fr., leg. Swallen.

41540 RGS, Vila Manresa p. P. Alegre, NEL, in silva, 14.5.1949, fr., leg. Swallen.

41793 RGS, Esteio p. S. Leopoldo, NEL, in silva campestri, 1.6.1949, fr., det. Swallen.

45627 RGS, S. Francisco de Paula, NEH, in araucarieto, 31.1.1950, fr.

49688 RGS, Lagoa dos Quadros p. Torres, NEC, in silva riparia, 18.1.1951, fr.

54946 RGS, Vila Oliva p. Caxias, NEH, in humidis umbrosis, 24.2.1949, fr., det. Swallen.

55372 SC, Serra da Boa Vista p. S. José, ad silvam, 4.2.1953, fr., leg. R. Reitz 7154, det. Swallen.

56780 RGS, Vila Oliva p. Caxias, NEH, in araucarieto, 8.2.1955, fr.

58592 RGS, Taimbesinho p. S. Francisco de Paula, NEH, in araucarieto, 12.2.1956, fr., fr.,

59025 RGS, Estação de Silvicultura p. Sta Maria, CL, in umbrosis, 12.2.1956, fr., leg. O. Camargo 166.

59832 RGS, Farroupilha, NEH, in silva, 15.2.1957, fl., leg. O. Camargo 1143.

59833 RGS, *ibidem*, 25.3.1957, fr., leg. O. Camargo 1313.

61964 RGS, Lageado, NEL, in silva, 4.12.1957, fl., leg. O. Camargo 2730.

64209 SC, S. Joaquim, in umbrosis, 10.1.1958, fl., leg. J. Mattos.

Habit: Prostrate, rooting at the lower joints, branches up to 50 cm long.

Habitat: Interior of rain forest.

Distribution in RGS: Frequent in the whole of the forested area.

General distribution: Through the whole of tropical and subtropical America.

SACCIOLEPIS Nash

Sacciolepis strumosa (Presl.) Chase,

1049 RGS, S. Leopoldo, NEL, in paludosis alte dumosis, 16.5.1933, fr., leg. C. Orth SJ, det. Chase, Parodi.

7032 SC, Lages, in paludosis alte dumosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

39536 RGS, Cachoeirinha p. Gravataí, NEL, in paludosis alte dumosis, 7.1.1949, fl.

42212 RGS, Fazenda do Arroio p. Osório, NEC, in paludosis alte dumosis, 4.1.1950, fr.

45351 RGS, Serra da Rocinha p. Bom Jesus, NEH, in paludosis alte graminosis, 18.1.1950, fr.

46177 RGS, Fazenda do Arroio p. Osório, NEC, in paludosis, 6.3.1950, fr.

54929 RGS, Vila Oliva p. Caxias, NEH, in paludosis alte dumosis, 24.2.1954, fr., det. Swallen.

56674 RGS, *ibidem*, 8.2.1955, fr., det. Swallen (Parodi: *S. campestris* (Nees)).

Habit: Culms up to 1 m high.

Habitat: Densely vegetated swamps.

Distribution in RGS: Northeastern quadrant.

General distribution:

Sacciolepis viloides (Trin.) Chase.

Panicum viloides Trin., De Gram. Pan. () 171; Döll, FB 2,2 (1877) 232, tab. 32; Ekman, Ark. f. Bot. 13,n.10 (1913) 27.

35145 RGS, Fazenda da Ronda p. Vacaria, NEH, in paludosis dense dumosis, 15.1.1947, fr., det. Parodi.

36467 RGS, Cambará p. S. Francisco de Paula, NEH, in paludosis alte graminosis, 2.1948, fr., det. Swallen.

62966 RGS, Bom Jesus, NEH, in paludosis, 6.1.1958, fl., leg. O. Camargo 3124.

62967 RGS, ibidem, 6.1.1958, fl., leg. O. Camargo 3146.

Habit: Very similar (identical?) to the precedent.

Habitat: Densely vegetated swamps.

Distribution in RGS: Northeastern Highlands.

General distribution: From Guyana to RGS.

SCLEROPOA Griseb.

Scleropoa rigida (L.) Gris., Spic. Fl. Rum. 2 (1844) 431; Herter, Flor. Urug. (1930) 38. — *Poa rigida* L., Spec. Plant. (1753) 5.

852 RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 15.10.1942, fr., leg. C. Orth SJ, det. Chase.

26979 RGS, ibidem, 10.1944, fr., det. Parodi.

29335 RGS, ibidem, 26.10.1945, fr., det. Parodi.

38079 RGS, ibidem, 17.11.1948, fr., det. Swallen.

56955 RGS, ibidem, 15.10.1955, fl.

Habit: Densely cespitose, up to 30 cm high.

Habitat: Disturbed soil.

Distribution in RGS: Northeastern Lowlands.

General distribution: Native to Europe; RGS, Uruguay, B. Aires.

SETARIA Beauv.

Setaria caespitosa Hack. et Arech., Gram. Urug. (1894) 146, fig. 15; Herter, Flor. Urug. (1930) 31; Cabrera, Fl. Alred. B. Aires (1954) 104, fig. 26 G.

60370 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fl., leg. O. Camargo 583.

Habit: Cespitose, 30-50cm high.

Habitat: Dry, grassy Campos.

Distribution in RGS: Western Campos area.

General distribution: RGS, Uruguay, B. Aires.

Setaria fiebrigii Herrm., Setaria (1909) 56; Cabrera, Fl. Alred. B. Aires (1953) 104.

721 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 5.12.1932, fr., leg. C. Orth SJ, det. Parodi (Chase: *S. setosa*(L.) Beauv.)

- 30739** RGS, Vila Oliva p. Caxias, NEH, in campestribus alte graminosis, 28.1.1946, fr., det. Swallen.
- 41799** RGS, Esteio p. S. Leopoldo, NEL, in campestribus siccis alte graminosis, 1.6.1949, fr.
- 42794** RGS, Sapucaia p. S. Leopoldo, NEL, in arenosis dumosis, 5.8.1949, fr., det. Swallen.
- 43628** RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus alte graminosis, 28.9.1949, fr., det. Swallen.
- 51085** RGS, Linha Pinhal p. Montenegro, NES, in dumosis secundariis, 20.11.1950, fr., leg. A. Sehnem.
- 57132** RGS, Esteio p. S. Leopoldo, NEL, in graminosis siccis, 14.11.1955, fr.

Habit: Bunchy, erect, 40 cm high.

Habit: Dry, grassy, Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: RGS, Uruguay, B. Aires.

Setaria geniculata (Lam.) Beauv., Ess. Agrost. 51, 178 (1812); Herter, Flor. Urug. (1930) 31; Cabrera, Fl. Alred. B. Aires (1953) 104; Reitz, Sellowia 7 (1956) 108. — *Panicum geniculatum* Lam., Encycl. Méthod. 4 (1797) 727; Döll, FB 2,2 (1878) 737.

722 RGS Vila Manresa p. P. Alegre, NEL, in incultis, 6.12.1932, fr., leg. C. Orth SJ, det. Chase.

838 RGS, ibidem, in campestribus siccis, 8.2.1933, fr., leg. C. Orth SJ, det. Parodi.

1851 RGS, Vacaria, NEH, in campestribus siccis graminosis, 6.1.1935, fr., leg. Dutra, det. Swallen.

2905 RGS, S. Borja, SW, in campestribus dense graminosis, 1943, fr., leg. fr. Baglione.

3901 RGS, Livramento, SW, in campestribus siccis, 12.1.1941, fr.

7008 SC, in campestribus siccis dumosis, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

7437 RGS, S. Leopoldo, NEL in campestribus siccis, 1907, fr., leg. F. Theissen SJ.

9867 RGS, Tupanciretan, NWH, in campestribus siccis, 29.1.1942, fr., det. Parodi, Swallen.

9873 RGS, ibidem, 29.1.1942, fr.

25566 RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus siccis, 1.1944, fr.

28473 RGS, Nonoai ad fl. Uruguay, NWU, in campestribus alte graminosis, 3.1945, fr.

28852 SC, Trindade p. Florianópolis, in arenosis siccis, 5.2.1945, fr., leg. A. Rohr SJ.

- 29166 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 12.9.1945, fr.
- 29381 RGS, ibidem, 3.11.1945, fr.
- 29687 RGS, Parecí p. Montenegro, NEL, in campestribus siccis, 10.9.1945, fr.
- 31232 RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis, 13.1.1946, fr.
- 32228 RGS, Azulega p. S. Francisco de Paula, NEH, in campestribus alte graminosis, 15.2.1946, fr.
- 33021 RGS, Parecí p. Montenegro, NEL, in campestribus alte graminosis, 10.10.1945, fr.
- 35161 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 15.1.1947, fr.
- 35607 RGS, S. Leopoldo, NEL, in campestribus alte graminosis, 20.12.1946, fr., leg. B. Wagner SJ.
- 36438 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus graminosis, 2.1948, fr., det. Swallen.
- 36443 RGS, ibidem, 2.1948, fr., det. Swallen.
- 37975 RGS, Barra do Ribeiro p. P. Alegre, NEL, in arenosis siccis, 14.11.1948, fr., det. Swallen.
- 39999 RGS, Linha Bonita p. Montenegro, NES, in incultis, 19.1.1949, fr.
- 40980 RGS, ad fl. Gravataí inferius p. P. Alegre, NEL, in campestribus siccis, 10.4.1949, fr., det. Swallen.
- 46324 RGS, Cristo Rei p. S. Leopoldo, NEL, in campestribus arenosis, 17.3.1950, fr., det. Swallen.
- 48341 SC, Figueiredo p. Bom Retiro, in campestribus subhumidis, 28.12.1948, fr., leg. R. Reitz 3472, det. Swallen.
- 52741 RGS, Uruguaiana, SW, in campestribus, 1.8.1952, fr., leg. et det. Beetle.
- 54341 SC, Laguna, prope litus in arenosis, 20.2.1952, leg. R. Reitz 5036, det. Swallen.
- 55072 RGS, Vila Oliva p. Caxias, NEH, in campestribus alte dumosis, 24.2.1954, fr.
- 55341 SC, Rio Negrinho, in campestribus dumosis, 26.1.1953, fr., leg. R. Reitz 7190, det. Swallen.
- 55358 SC, Massiambú p. Palhoça, in arenosis, 5.2.1953, fr., leg. R. Reitz 7168, det. Swallen.
- 55386 SC, Anitápolis p. Palhoça, in dumosis, 2.4.1953, fr. leg. R. Reitz 7128, det. Swallen.
- 55422 SC, Massiambú p. Palhoça, in arenosis, 2.3.1953, fr., leg. R. Reitz 7069, det. Swallen.
- 56005 RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 19.11.1954, fr.

57025 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in siccis graminosis, 26.10.1955, fr.

57511 RGS, Estação de Silvicultura p. Sta Maria, CN, in siccis graminosis, 30.11.1955, fl., fr., leg. O. Camargo 70.

58032 SC, Ibirama, in incultis, 20.10.1953, fr., leg. R. Reitz 8274, fr., det. Swallen.

58650 RGS, Estação de Silvicultura p. Sta Maria, CN, in alte graminosis, 20.1.1956, fr., leg. O. Camargo 70.

59390 RGS, Farroupilha, NEH, in siccis alte graminosis, 15.11.1956, fr. leg. O. Camargo 955.

60372 RGS, Estação de Silvicultura p. Sta Maria, CN, 8.4.1956, fr., leg. O. Camargo 615.

60373 RGS, ibidem, 8.4.1956, fr., leg. O. Camargo 580.

61982 RGS, Taquarí, NEL, in incultis, 8.12.1957, fr., leg. O. Camargo 2765.

64208 SC, S. Joaquim, in incultis, 10.1.1958, fr., leg. J. Mattos.

64243 SC, Lages, 17.2.1958, fl., leg. J. Mattos, in campestribus.

Habit: Up to 50 cm high.

Habitat: Dry Campos; often weed on cultivated soil.

Distribution in RGS: Frequent over the whole cultivated and Campos area.

General distribution: South Brazil, Uruguay, B. Aires, Chile.

Setaria lachnea (Nees) Kunth,

Panicum lachneum Nees, Agrost. Brasil. (1829) 248; – P. setosum Swartz, Fl. Ind. occid. 1 () 139; Döll, FB 2,2 (1877) 162.

1013 RGS, S. Leopoldo, NEL in subsiccis umbrosis, 14.8.1934, fr. leg. C. Orth SJ, det. Chase.

36473 RGS, Cambará p. S. Francisco de Paula, NEH, in araucarieto, 2.1948, fr., det. Swallen.

40540 RGS, Sapucaia p. S. Leopoldo, NEL, in subhumidis umbrosis, 17.3.1949, fr.

40817 RGS, Granja Neugebauer p. Itapoan, NEL, in silva ad viam, 3.4.1949, fr., det. Swallen.

42108 RGS, Schwabenschneis p. Novo Hamburgo, NES, in humidis umbrosis, 23.06.1949, fr.

42514 RGS, Sta Rita p. Farroupilha, NEH, in humidis, 13.7.1949, fr.

46373 RGS, Gramado p. Canela, NEH, in dumosis umbrosis, 20.3.1950, fr.

46460 RGS, Lagoa da Pinguela p. Osório, NEC, in dumosis umbrosis, 27.3.1950, fr.

59776 RGS, Farroupilha, NEH, in silva subhumida, 12.3.1957, fr., leg. O. Camargo 1274.

60371 RGS, Estação de Silvicultura p. Sta Maria, CL, in dumosis, 8.4.1956, fr., leg. O. Camargo 594.

61983 RGS, Taquarí, NEL, in silva aperta, 8.12.1957, fl., leg. O. Camargo 2756.

64272 SC, S. Joaquim, in umbrosis, 10.1.1958, fl., leg. J. Mattos.

Habit: Slender, culms up to 1 m high.

Habitat: More or less wet and shady places.

Distribution in RGS: Over the whole of the forested area.

General distribution: From Mexico to RGS.

Starria onurus (Willd.) Griseb.,

11962 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis dumosis, 1943, fr.

29189 RGS, ibidem, 12.9.1945, fr., det. Swallen.

33515 RGS, S. Leopoldo, NEL, in campestribus siccis dumosis, 17.1.1946, fr., leg. L. Stahl SJ.

36439 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus graminosis, 2.1948, fr., det. Swallen.

44464 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis alte dumosis, 19.1.1949, fr.

57018 RGS, ad montem Sapucaia, NEL, in alte graminosis, 26.10.1955, fl.

57208 RGS, Vila Manresa p. P. Alegre, NEL, in alte graminosis, 22.10.1955, fr.

57215 RGS, ibidem, 15.10.1955, fl.

61981 RGS, Farroupilha, NEH, in alte graminosis, 18.11.1957, fr., leg. O. Camargo 2555.

62771 RGS, P. Alegre, NEL, 1.11.1957, fr., leg. O. Camargo 2393.

Habit: Slender, up to 75 cm high.

Habitat: Dry Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: RGS, Uruguay.

Setaria poiretiana (Schult.) Hitchc., Panicum poiretianum Schult., Mant. 2 () 229. — P. speciosum A. Br., Ind. sem. hort. Berol., App. () 22; Döll, FB 2,2 (1877) 154.

863 RGS, Vila Manresa p. P. Alegre, NEL, ad rivum in alte dumosis, 4.1.1933, fr., leg. C. Orth SJ, det. Chase, Parodi, Swallen.

1073 RGS, S. Leopoldo, NEL, ad silvam campestrem in humidis, 5.5.1934, fr., leg. C. Orth SJ, det. Parodi (Chase: P. sulcatum Aubl.).

- 32838** RGS, Butterberg p. Montenegro, NES ad rivum in alte dumosis, 17.12.1945, fr.
- 40475** RGS, Vila Manresa p. P. Alegre, NEL, ad rivum in dense dumosis, 14.3.1949, fr., det. Swallen.
- 40789** RGS, Sapucaia p. S. Leopoldo, NEL, ad rivum in dumosis, 1.4.1949, fr., det. Swallen.
- 40919** RGS, ad montem das Cabras p. S. Leopoldo, NEL, in humidis alte herbosis, 8.4.1949, fr.
- 42294** RGS, ad montem Steinkopf p. S. Leopoldo, NEL, in subpaludosis dense dumosis, 28.6.1949, fr.
- 46369** RGS, Gramado p. Canela, NEH, in humidis alte dumosis, 26.3.1950, fr.
- 47206** RGS, Kappesberg p. Montenegro, NES, in subpaludosis alte graminosis, 4.7.1950, fr.
- 50695** RGS, ibidem, 25.3.1950, fr., leg. A. Sehnem SJ 4885.
- 58054** SC, Ibirama, ad silvam, 1.3.1954, fr., leg. R. Reitz 8333, det. Swallen.

Habit: Up to 1,5 high.

Habitat: Wet to swampy places, among high grasses and shrubs.

Distribution in RGS: Northeastern Lowlands and rain forest belt.

General distribution: From Amazonia to RGS.

Setaria scandens Schrad., in Schult. Mant. 2 () 279. — *Panicum scandens* Trin., Diss. 2 () 166; Döll, FB 2,2 (1877) 170.

46489 RGS, Lagoa da Pinguela p. Osório, NEC, in dumosis secundariis, 27.3.1950, fr., det. Parodi.

Habit: Up to 75 cm high, culm very rough, climbing.

Habitat: Wet, secondary growth.

Distribution in RGS: No other findplace: NEC.

General distribution: From Amazonia to RGS.

Setaria vaginata Spreng., Syst. Veg. 4,2 (1827) 33; Herter, Flor. Urug. (1930) 31; Cabrera, Fl. Alred. B. Aires (1953) 104. — *Panicum setosum* Swartz, Fl. Ind. occid. 1 () 139; Döll, FB 2,2 (1877) 162.

56332 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in alte graminosis, 2.1.1955, fr., det. Parodi (Swallen: *S. fiebrigii* Herrm.).

Habit: Culms up to 1 m high.

Habitat: More or less wet Campos.

Distribution in RGS: No other findplace.

General distribution: From the West Indies and Mexico to Uruguay and B. Aires.

Setaria verticillata (L.) Beauv., Essay (1812) 51; Herter, Flor. Urug. (1930) 31; Cabrera, Fl. Alred. B. Aires (1953) 104. — *Panicum verticillatum* L., Spec. Plant. ed. 2 (1762) 82; Döll 2,2 (1877) 172.

1012 RGS, S. Leopoldo, NEL, in incultis, 15.10.1934, fr., leg. C. Orth SJ, det. Chase.

Habit: Erect, 50 cm high.

Habitat: Dry Campos; weedy on cultivated soil.

Distribution in RGS: Only findplace.

General distribution: From Rio de Janeiro to Uruguay and B. Aires.

Setaria vulpiseta (Lam.) Hitchc. et Chase,

28079 RGS, Vila Oliva p. Caxias, NEH, in subpaludosis dumosis, 2.1945, fr. leg. P. Buck.

28492 RGS, Nonoai ad Fl. Uruguay, NWU, ad rivum in humidis, 3.1945, fr.

30716 RGS, Vila Oliva p. Caxias, NEH, ad rivum in alte dumosis, 28.1.1946, fr.

31322 RGS, ibidem, 24.2.1946, fr.

40314 RGS, Sta Rita p. Farroupilha, NEH, in paludosis alte dumosis, 29.1.1949, fr., det. Swallen.

42420 RGS, Pareci p. Montenegro, NEL, in humidis alte dumosis, 7.7.1949, fr.

45712 RGS, Sta Rita p. Farroupilha, NEH, in paludosis alte dumosis, 7.2.1950, fr.

46465 RGS, Lagoa da Pinguela p. Osório, NEC, in paludosis alte dumosis, 27.3.1950, fr.

49813 SC, Itapiranga p. fl. Uruguay, ad fl. in alte dumosis, 6.2.1951, fr.

59835 RGS, Farroupilha, NEH, ad silvam, 25.3.1957, fr., leg. O. Camargo 1316.

59836 RGS, ibidem, 18.3.1957, fr., leg. O. Camargo 1163.

59994 RGS, ibidem, 25.3.1957, fr., leg. O. Camargo 1184.

60274 RGS, ibidem, 14.1.1957, fr., leg. O. Camargo 1083.

Habit: Culms up to 2 m high.

Habitat: Wet, shrubby places among high grasses.

Distribution in RGS: Over the whole northern area of the State.

General distribution:

SORGHASTRUM Nash

Sorghastrum agrostoides (Speg.) Hitchc., Contr.U.S.Nat.Herb. 17, 3 (1913) 211. — *Andropogon agrostoides* Speg., An.Soc.Ci.Arg. 16 (1883) 136.

56740 RGS, Vila Oliva p. Caxias, NEH, in alte graminosis, 8.2.1955, fr., det. Parodi (Swallen: Setosum (Gris.) Hitchc.).

Habit: Culms up to 1 m high.

Habitat: Dry, grassy Campos among shrubs.

Distribution in RGS: No other findplaces.

General distribution:

Sorghastrum minarum (Nees) Hitchc.,

Trachypogon minarum Nees, Agrost. Bras. (1829) 349; Ekmann, Ark. f. Bot. 11, n.4 (1912) 9. — Sorghum minarum (Nees) Hack., FB 2,3 (1883) 276, tab. 63.

6954 SC, Lages, in alte graminosis, 1935, fl., fr., leg. A. Bruxel SJ, det. Swallen.

6961 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

7020 SC, ibidem, 1935, fr., leg. A. Bruxel SJ, det. Swallen.

58990 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 5.1.1956, fr., leg. O. Camargo 143.

Habit: Erect, stiff, up to 1 m high.

Habitat: Dry Campos.

Distribution in RGS: Western Campos, apparently rare.

General distribution: From Minas Gerais to RGS.

Sorghastrum nutans (L.) Nees,

Andropogon nutans L., Spec. Plant. (1753) 1045; Ekmann, Ark. f. Bot. 11, n.4 (1911) 9. — Sorghum nutans A. Gray, ssp. linnaeanum Hack., FB 2,3 (1883) 276.

1100 RGS, Sapucaia p. S. Leopoldo, NEL, in incultis, 28.9.1934, fr., leg. C. Orth SJ, det. Chase, Parodi.

Habit: Culms up to 1,5 m high.

Habitat: Dry Campos among high grasses and shrubs.

Distribution in RGS: Northeastern Lowlands.

General distribution: From southern USA to RGS.

Sorghastrum parviflorum (Desv.) Hitchc. et Chase.

1010 RGS, Portão p. S. Leopoldo, NEL, in alte paludosis, 8.9.1934, fr., leg. C. Orth SJ, det. Chase.

Habit: Culms up to 2 m high.

Habitat: Swampy places among high grasses and shrubs.

Distribution in RGS: No other findplaces.

General distribution:

Sorghastrum setosum (Gris.) Hitchc.,

55068 RGS, Vila Oliva p. Caxias, NEH, in paludosis alte graminosis, 24.2.1954, fr., det. Swallen.

63456 SC, S. Joaquim, in campestribus, 18.2.1958, fl., leg. J. Mattos.

Habit: Culms up to 150 cm high.

Habitat: Swampy Campos.

Distribution in RGS: Northeastern Highlands.

General distribution:

Sorghastrum stipoides(H.B.R.) Nash,

Andropogon stipoides H.B.K., Nov. Gen. et Spec. 1 (1816) 189. — *Sorghum nutans* A. Gray. ssp. *avenaceum* (Michx.) Hack., var. *stipoides* (HBK) Hack., FB 2,3 (1883) 274.

36865 RGS, Morro da Polícia p. P. Alegre, NEL, in campestribus siccis alte graminosis, 6.11.1948, fr., det. Swallen.

43604 RGS, ibidem, 28.9.1949, fr., det. Swallen.

44077 RGS, Barreto Vianna p. S. Leopoldo, in campestribus alte dumosis, 24.10.1949, fr., det. Swallen.

44480 RGS, Granja Neugebauer p. Itapoan, NEL, in alte graminosis 19.11.1949, fr., det. Beetle.

54463 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 12.11.1953, fr., det. Swallen.

57138 RGS, ibidem, 4.11.1955, fl.

57510 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 18.11.1955, fr., leg. O. Camargo 19.

58925 RGS, ibidem, 1.3.1956, leg. O. Camargo 78.

61985 RGS, Farroupilha, NEH, in alte graminosis, 25.11.1957, fl., leg. O. Camargo 2629.

62750 RGS, ibidem, 10.11.1957, fl., leg. O. Camargo 2506.

Habit: Culms up to 15 m high.

Habitat: More or less dry, shrubby, Campos.

Distribution in RGS: Apparently over the whole of the northern half of the State.

General distribution: From Mexico to RGS.

SPARTINA Schreb.

Spartina ciliata Kunth, Rev. 1 () 94; Döll, FB 2,3 (1878) 85; Ekman, Ark. f. Bot. 13, n.10 (1913) 44; Herter, Flor. Urug. (1930) 39; Reitz, Sellowia 7 (1956) 109.

52295 Paraná, Paranaguá, ad litus in arenosis, 8.3.1951, leg. G. Hatschbach 2150, det. Parodi.

56450 RGS, Cidreira, NEC, in arenosis prope litus, 17.2.1950, fl., det. Parodi, Swallen (*S. ciliata* Brongn.).

Habit: Culms stiff, up to 75 cm high.

Habitat: More or less fixed sand dunes.

Distribution in RGS: Sea coast.

General distribution: From Paraná to RGS.

SPOROBOLUS R. Br.

Sporobolus aeneus (Trin.) Kunth, Enum. 1 () 213; Ekman, Ark. f. Bot. 13, n.10 (1913) 41. — *Vilfa aenea* Trin., Spec. Gram. 2 () tab. 23; Döll, FB 2,3 (1878) 33.

44462 RGS, Granja Neugebauer p. P. Alegre, NEL, in campestribus siccis alte graminosis, 19.11.1949, fr., det. Parodi.

Habit: Bunchy, erect, up to 50 cm high.

Habitat: Dry, shrubby Campos.

Distribution in RGS: No other findplace.

General distribution: From Ceará to RGS.

Sporobolus eximius (Nees) Ekman, Ark. f. Bot. 13, n.10 (1913) 41, fab. 2, f. 3. — *Vilfa eximia* Nees ap. Trin., Act. Petrop. 1845, 77; Döll, FB 2,3 (1878) 35.

44022 RGS, Morro do Osso p. P. Alegre, NEL, in campestribus siccis, 21.10.1949, fr., det. Swallen.

Habit: Bunchy, up to 75 cm high.

Habitat: Dry, grassy, Campos.

Distribution in RGS: No other findplace.

General distribution: From S. Paulo to RGS.

Sporobolus poiretii (R. et S.) Hitchc., Bartonia 14 (1932) 32; Cabrera, Fl. Alred. B. Aires (1953) 79; Reitz, Sellowia 7 (1956) 109. — *Axonopus poiretii* R. et S., Syst. Veg. 2 (1877) 318.

713 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, leg. C. Orth SJ, det. Swallen (Chase: *S. tenacissimus* Veauv.).

877 RGS, ibidem, in incultis, 5.12.1933, fr., leg. C. Orth SJ (Chase: *S. bereteroanus* (Trin.) Hitchc.).

1958 RGS, Carauno p. Bom Jesus, NEH, in campestribus siccis, 2.2.1934, fr., leg. J. Dutra, det. Chase.

25587 RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus siccis, 1.1944, fr., det. Swallen.

29181 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 12.9.1945, fr., leg. Swallen.

- 29377 RGS, ibidem, 3.11.1945, fr., det. Swallen.
- 31417 SC, Sombrio p. Araranguá, in arenosis alte graminosis, 31.1.1946, fr., det. Swallen.
- 32648 RGS, Parecí p. Montenegro, NEL, in campestribus siccis, 5.12.1945, fr., leg. E. Henz SJ.
- 33194 RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis, 16.1.1946, fr.
- 33490 RGS, S. Leopoldo, NEL, in campestribus siccis, 17.7.1946, fl.
- 33910 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis dense graminosis, 18.9.1946, fr.
- 34217 Chile, Las Animas p. Valdivia, 14.2.1937, fr., in siccis arenosis, leg. et det. H. Gunckel (S. bertoroanus (Trin.) Hitchc.).
- 38076 RGS, Vila Manresa p. P. Alegre, NEL, in agro, 17.11.1948, fr., det. Swallen.
- 39848 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in arenosis dumosis, 12.1.1949, fr.
- 42449 RGS, Parecí p. Montenegro, NEL, in campestribus siccis, 7.7.1949, fr., det. Swallen.
- 44712 RGS, Lagoa dos Barros p. Osório, NEC, in arenosis alte graminosis, 24.11.1949, fr.
- 45945 RGS, Lagoa dos Quadros p. Torres, NEC, in arenosis dumosis, 21.2.1950, fr.
- 54953 RGS, Vila Oliva p. Caxias, NEH, in graminosis humidis, 24.2.1954, fr., det. Swallen.
- 55348 SC, Rio Tavares in Insula Florianópolis, ad viam in incultis tritis, 13.5.1953, fr., leg. R. Reitz 7182, det. Swallen.
- 55390 SC, Anitápolis p. Palhoça, in dumosis, 2.4.1953, leg. R. Reitz 7124, det. Swallen.
- 55428 SC, Rio Tavares in Insula Florianópolis, in arenosis dense graminosis, 23.12.1952, fr., leg. R. Reitz 7060, det. Swallen.
- 55438 SC, Mafra, in campestribus alte graminosis, 26.1.1935, fr., leg. R. Reitz 7045, det. Swallen.
- 57017 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fl.
- 57224 RGS, Vila Manresa p. P. Alegre, NEL, in humidis, 15.10.1955, fr.
- 58926 RGS, Estação de Silvicultura p. Sta Maria, CL, in humidis alte graminosis, 1.3.1956, fr., leg. O. Camargo 75.
- 59386 RGS, Farroupilha, NEH, in alte graminosis, 15.11.1956, fl., leg. O. Camargo 993.
- 59993 RGS, ibidem, 25.3.1956, fr., leg. O. Camargo 1185.
- 60368 RGS, Estação de Silvicultura p. Sta Maria, CL, in alte graminosis, 8.4.1956, fr., leg. O. Camargo 612.

62968 RGS, Taquarí, NEL, in incultis, 10.12.1957, fl., leg. O. Camargo 3039.

63093 RGS, Pelotas, SEC, in incultis, 10.3.1956, fl., leg. Schlichting.

64224 SC, S. Joaquim, in incultis, 10.1.1958, fl., leg. J. Mattos.

Habit: Bunchy, erect, up to 60 cm high.

Habitat: Dry Campos, often weedy on disturbed soil.

Distribution in RGS: Over the whole of the cultivated and Campos area.

General distribution: Common through most of South America; cosmopolitan.

Sporobolus pseudairoides Parodi,

50862 Paraná, Florestal p. Piraquara, in dumosis secundariis, 6.2.1949, fr., leg. G. Hatschbach 1172, det. Parodi.

59733 RGS, Farroupilha, NEH, in alte graminosis, 28.5.1956, fr., leg. O. Camargo 730.

Habit: Vigorous bunches, up to 1 m high.

Habitat: Somewhat wet, secondary growth.

Distribution in RGS: Northeastern Highlands.

General distribution:

Sporobolus pyramidalis Beauv., Flor. Owar. 2 () 36; Reitz, Selowia 7 (1956) 109.

2500 SC, Maratá p. P. União, in incultis dumosis, 22.12.1936, fr., leg. C. Orth SJ, det. Swallen.

40688 RGS, Vila Manresa p. P. Alegre, NEL, in incultis dumosis, 26.3.1949, fr., det. Swallen.

49854 SC, Itapiranga ad fl. Uruguay superius, ad viam in incultis, 6.2.1951, fr.

58060 SC, Saiguassú p. S. Francisco, in arenosis, 3.1.1954, fr., det. leg. R. Reitz 8364, det. Swallen.

Habit: Bunchy, up to 1 m high.

Habitat: Secondary growth on former cultivated land.

Distribution in RGS: One Only findplace.

General distribution: Native to tropical Africa.

Sporobolus virginicus (L.) Kunth, Rev. Gram. 1 () 67; Ekman, Ark. f. Bot. 13,n.10 (1913) 41; Retiz, Selowia 7 (1956) 109. — *Vilfa virginica* L., Spec. Plant. (1753) 63; Döll, FB 2,3 (1878) 30, tab. 8, fig. 1.

52288 Paraná, Paranaguá, ad litus in arenosis, 14.1.1950, fl., leg. G. Hatschbach 1677, det. Parodi.

54334 SC, Praia Braba p. Itajaí, in arenosis late repens, 10.2.1950, fr.

55404 SC, Rio Tavares in Insula Florianópolis, prope litus in arenosis, 12.3.1953, leg. R. Reitz 7105, det. Swallen.

56851 SC, Arroio do Silva p. Araranguá, ad litus in arenosis, 19.2.1955, fr.

Habit: Prostrate, rooting at almost all joints.

Habitat: Dry sandy places.

Distribution in RGS: To be expected at the northern sea coast.

General distribution: From Guyana to RGS.

STENOTAPHRUM Trin.

Stenotaphrum secundatum (Walt.) O. Kuntze, Rev. Gen. Plant. 2 (1891) 794; Cabrera, Fl. Alred. B. Aires (1953) 104; Reitz, Sellowia 7 (1956) 109. — *Ischaemum secundatum* Walt., Flor. Carolin. (1788) 249. — *Stenotaphrum glabrum* Trin., Gram. Panic. (1826) 60; Döll, FB 2,2 (1877) 300, tab. 39;

720 RGS, Vila Manresa p. P. Alegre, NEL, in pascuo, 5.12.1932, fl., leg. C. Orth SJ, det. Swallen, Chase, Parodi: *S. secundatum* (Walt.) O.K.

7448 RGS, S. Leopoldo, NEL, in graminosis, 1907, ster., leg. F. Theissen SJ.

35140 RGS, Fazenda da Ronda p. Vacaria, NEH, in campestribus siccis, 15.1.1947, fr.

38585 RGS, ad montem das Cabras p. S. Leopoldo, NEL, in arenosis graminosis, 7.12.1948, fl.

47586 SC, Azambuja p. Brusque, in pascuo, 28.2.1948, fl., leg. R. Reitz 2871, det. Parodi (*S. glabrum* Trin.).

47613 SC, Azambuja p. Brusque, in pascuo, 28.2.1948, fl., leg. R. Reitz 2871, det. Parodi.

55368 SC, Barra do Sul p. Araquari, in arenosis, 9.4.1953, leg. R. Reitz 7158, det. Swallen.

55388 SC, Anitápolis p. Palhoça, in dumosis, 2.4.1953, fr., leg. R. Reitz 7126, det. Swallen.

55406 SC, Palhoça p. Florianópolis, in arenosis, 22.12.1952, fr., det. Swallen.

55432 SC, Barra do Sul p. Araquari, in arenosis, 9.1.1953, fl., leg. R. Reitz 7053, det. Swallen.

55433 SC, Rio Tavares in Insula Sta Catarina, in arenosis, 12.3.1953, fr., leg. R. Reitz 7051, det. Swallen.

56508 RGS, Torres, NEC, in arenosis, 19.1.1955, fl.

60670 RGS, Farroupilha, culta, 7.5.1957, fr., leg. O. Camargo 1368.

Habit: Prostrate, densely matted.

Habitat: Native to RGS? Often cultivated in meadows.

Distribution in RGS: Over the whole of the cultivated area.
General distribution: From Guyana to Uruguay and B. Aires.

STIPA L.

Stipa amethystina Steud., Syn. Pl. Gram. (1855) 124.

44474 RGS, Granja Neugebauer p. Itapoan, NEL, in campestribus siccis alte dumosis, 19.11.1949, fr., det. Parodi.

57126 RGS, Esteio p. S. Leopoldo, NEL, in siccis alte graminosis, 14.11.1955, fr.

Habit: Culms up to 1 m high.

Habitat: Dry, shrubby Campos.

Distribution in RGS: Chile, RGS.

General distribution: Chile, RGS.

Stipa filifolia Nees, Agrost. Bras. (1829) 10; Döll, FB 2,3 (1878) 10; Herter, Flor. Urug. (1930) 34.

38125 RGS, Morro da Policia p. P. Alegre, NEL, in campestribus siccis alte graminosis, 20.11.1948, fr., det. Swallen.

43622 RGS, ibidem, 28.9.1949, fl.

43705 Morro Sta Teresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 3.10.1949, fl.

44039 RGS, Morro do Osso p. P. Alegre, NEL, in campestribus alte graminosis, 21.10.1949, fl., det. Swallen.

Habit: Bunches up to 50 cm high, culms filiform.

Habitat: Dry Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: RGS and Uruguay.

Stipa juergensii Hack.,

44021 RGS, Morro do Osso p. P. Alegre, NEL, in campestribus siccis, 21.10.1949, fr., det. Swallen.

Habit: Culms up to 60 cm high.

Habitat: Dry Campos.

Distribution in RGS: Central and northeastern Lowlands.

General distribution: Uruguay and RGS.

Stipa megapotamica Spreng., Acta Petrop. 1829, 77; Döll, FB 2,3 (1878) 9; Herter, Flor. Urug. (1930) 34; Cabrera, Fl. Alred. B. aires (1953) 82.

1097 RGS, ad montem Sapucaia p. S. Leopoldo, NEL, in campestri-

bus siccis alte graminosis, 17.10.1934, fr., det. Parodi (Chase: *S. hackelii* Arech.).

2736 RGS, S. Leopoldo, NEL, in campestribus siccis alte dumosis, 30.10.1935, fr., leg. C. Orth SJ, det. Chase.

8819 RGS, Bom Jesus, NEH, in campestribus subsiccis alte graminosis, 15.1.1942, fr., det. Swallen.

32493 RGS, Parecí p. Montenegro, NEL, in campestribus alte graminosis, 17.10.1945, fr., leg. E. Henz SJ.

33084 Uruguay, Juan Jackson p. Soriano, in campestribus siccis alte graminosis, 11.1937, fr., leg. B. Rosengurtt B-2317, det. Parodi.

43644 RGS, Morro das Abertas p. P. Alegre, NEL, in campestribus siccis alte graminosis, 30.9.1949, fr.

62751 RGS, Porto Alegre, NEL, in campestribus, 2.11.1957, fl., leg. O. Camargo 2385.

63391 RGS, Pelotas, SEC, in dumosis, 5.10.1954, fl., leg. J. Sacco 189, det. Parodi, Swallen.

Habit: Dense bunches, culms up to 50 cm high.

Habitat: Dry Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: RGS, Uruguay, B. Aires.

Stipa melanosperma Presl, ap. Spegazzini. An. Mus. Nac. Mont. 4 (1901) 67; Ekman, Ark. f. Bot. 13, n.10 (1913) 40; Herter, Flor. Urug. (1930) 34.

1953 RGS, Quilombo p. S. Leopoldo, NEL, in campestribus siccis alte dumosis, 9.10.1934, fl., fr., leg. C. Orth SJ, det. Chase.

2733 RGS, ibidem, 23.10.1933, fl., leg. C. Orth SJ, det. Swallen.

8351 RGS, Montserrat p. P. Alegre, NEL, in campestribus siccis alte graminosis, 1.12.1941, fr.

11767 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 1943, fl.

28929 RGS, P. Alegre, NEL, in campestribus siccis alte graminosis, 1944, fl., leg. Irm. Augusto.

33044 Uruguay, Juan Jackson p. Soriano, in rupestribus siccis graminosis, leg. B. Rosengurtt PE-4483, det. Chase, Parodi.

44030 RGS, Morro do Osso p. P. Alegre, NEL, in campestribus siccis alte graminosis, 21.10.1949, fl., det. Swallen.

63375 RGS, Pelotas, SEC, in campestribus siccis, 10.11.1954, fr., leg. J. Sacco 217.

Habit: Culms up to 1 m high.

Habitat: Dry, more or less shrubby, Campos.

Distribution in RGS: Northeastern Lowlands.

General distribution: From Paraná to Uruguay.

***Stipa nutans* Hack.,**

2911 RGS, S. Borja, SW, in siccis alte dumosis, 1942, fr., leg. Fr. Baglione.

43863 RGS, ad fl. Caí inferius p. P. Alegre, NEL, in siccis alte graminosis, 12.10.1949, fr., det. Swallen.

Habit: Culms up to 75 cm high.

Habitat: Dry, more or less shrubby Campos.

Distribution in RGS: Central and northeastern Lowlands.

General distribution: RGS, Uruguay.

***Stipa sellowia* Nees, ap. Trin. in Act. Petrop. 1848, 38; Döll, FB 2,3 (1878) 8; Ekman, Ark. f. Bot. 1, n.10 (1913) 40.**

56318 RGS, Fazenda Englert p. S. Francisco de Paula, NEH, in campestribus alte graminosis, 2.1.1955, fr., det. Parodi.

56354 RGS, ibidem, 2.1.1955, fr., det. Parodi.

Habit: Culms up to 1,5 m high.

Habitat: More or less dry Campos.

Distribution in RGS: Northeastern Highlands.

General distribution: From Paraná to RGS.

TRACHYPOGON Nees

***Trachypogon canescens* Nees, Agrost. Bras. (1829) 111; Cabrera, FL Alred. B. Aires (1953) 111. — *T. polymorphus* Hack., var. *canescens* (Nees) Hack., FB 2,3 (1883) 263.**

7024 SC, Lages, in campestribus siccis alte graminosis, 1953, fr., leg. A. Bruxel SJ.

29149 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis alte graminosis, 12.9.1945, fr.

29149 RGS, ibidem, 12.9.1949, fr.

30745 RGS, Vila Oliva p. Caxias, NEH, in campestribus siccis alte graminosis, 28.1.1946, fr.

32757 RGS, Pareci p. Montenegro, NEL, in campestribus siccis dense graminosis, 17.10.1945.

38096 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 17.11.1948, fr.

44031 RGS, Morro do Osso p. P. Alegre, NEL, in siccis alte graminosis, 21.10.1949, fr., Swallen: *T. filifolius* (Hack.) Hitchc.).

44471 RGS, Granja Neugebauer p. Itapoan, NEL, in siccis alte graminosis, 19.11.1949, fr.

55995 RGS, Vila Manresa p. P. Alegre, NEL, in campestribus alte graminosis, 19.11.1954, fr.

- 60708** SC, S. Joaquim, in campestribus, 28.12.1955, fr., leg. J. Mattos.
58511 RGS, *ibidem*, 19.11.1955, fr.

Habit: Dense bunches, up to 75 cm high.

Habitat: Dry, somewhat shrubby Campos.

Distribution in RGS: Northeastern Lowlands and Highlands.

General distribution: Southern Brazil, Uruguay, B. Aires.

TRICHACHNE Nees

Trichachne insularis (L.) Nees, Agrost. Bras. (1829) 66; Herter, Flor. Urug. Addit.; Reitz, *Sellowia* 7 (1956) 109. — *Andropogon insularis* L., Spec. Plant. ed. 2 (1762) 1480. — *Panicum leucophaeum* H.B.K., Nov. Gen. et Spec. 1 (1816) 97; Döll, FB 2,2 (1877) 136.

1067 RGS, S. Leopoldo, NEL, in incultis, 8.11.1933, fr., leg. C. Orth SJ, det. Chase.

10835 RGS, Marcelino Ramos ad. fl. Uruguay, NWU, in cultis, 1.1943, fr., leg. E. Friderichs SJ.

10837 RGS, Lagoa Vermelha, NEH, ad viam, 1.1943, fr., leg. E. Friderichs SJ.

39936 RGS, ad montem Ferrabraz p. Novo Hamburgo, NEL, in arenosis alte graminosis, 12.1.1949, fr.

41217 RGS, ad fl. Caí inferius p. P. Alegre, NEL, in incultis, 22.4.1949, fr., det. Swallen.

41481 RGS, Estação Azevedo p. Montenegro, NEL, in incultis, 6.5.1949, fr., det. Swallen.

53278 RGS, Caaró p. S. Luiz, in incultis alte graminosis, 24.11.1952, fr.

55353 SC, Rio Tavares in Insula Sta Catarina, in arenosis alte dumosis, 13.5.1953, leg. R. Reitz 7174, det. Swallen.

58930 RGS, Estação de Silvicultura p. Sta Maria, CN, in alte graminosis, 28.2.1956, fl., leg. O. Camargo 33.

59524 RGS, Farroupilha, NEH, in dumosis alte graminosis, 22.10.1956, fl., leg. O. Camargo 855.

59834 RGS, *ibidem*, 25.3.1957, fr., leg. O. Camargo 1314.

60338 RGS, Estação de Silvicultura p. Sta Maria, CL, in graminosis, 8.4.1956, fr., leg. O. Camargo 554.

60339 RGS, *ibidem*, 8.4.1956, fr., leg. O. Camargo 600.

62913 RGS, Taquarí, NEL, in campestribus, 10.12.1957, fl., leg. O. Camargo 3026.

Habit: Up to 1 m high.

Habitat: Disturbed soil; secondary brushwood.

Distribution in RGS: Over the whole cultivated area.

General distribution: From southern USA through South America.

TRICHOLAENA Schrad.

- Tricholaena rosea** Nees, Flor. Afr. Austr. 1 () 17; Reitz, Sellowia 7 (1956) 108 sub *Rhychelytrum roseum* (Nees) Tapf et Hubb.
- 1065** RGS, S. Leopoldo, NEL, in arenosis incultis, 5.5.1934, fr., leg. C. Orth SJ, det. Parodi.
- 10831** RGS, Marcelino Ramos, NWU, ad viam, 1.1943, fr., leg. E. Friderichs SJ.
- 10838** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 1.1943, fr., det. Parodi.
- 26825** RGS, Belém Velho p. P. Alegre, NEL, in incultis, 27.9.1944, fr., leg. P. Capparelli, det. Parodi.
- 29188** RGS, Vila Manresa p. P. Alegre, NEL, in campestribus siccis, 12.9.1945, fr., det. Swallen.
- 29251** RGS, ibidem, 10.10.1945, fr.
- 32493** RGS, Pareci p. Montenegro, NEL, in incultis, 17.10.1945, fr.
- 33489** RGS, S. Leopoldo, NEL, in incultis, 17.7.1946, fr., leg. E. Henz SJ.
- 37386** RGS, Vila Manresa p. P. Alegre, NEL, in incultis, 5.7.1946, fr.
- 42118** RGS, Schwabenschneis p. S. Leopoldo, NES, in incultis, 23.6.1949, fr.
- 42313** RGS, Langschneis p. Montenegro, NES, ad viam ferream, 2.7.1949, fr.
- 57013** RGS, in summo monte Sapucaia p. S. Leopoldo, NEL, in alte graminosis, 26.10.1955, fl.
- 58882** RGS, Estação de Silvicultura p. Sta Maria, CL, in incultis, 28.2.1956, fl., leg. O. Camargo.
- 58991** RGS, ibidem, 5.1.1956, fl., leg. O. Camargo 147.
- 59871** RGS, Farroupilha, NEH, in incultis, 25.3.1957, fr., leg. O. Camargo 1197.
- 62335** RGS, Montenegro, NEL, in incultis, 19.9.1957, fl., leg. O. Camargo 1781.
- 62684** RGS, P. Alegre, NEL, in incultis, 2.11.1957, fl., leg. O. Camargo 2383.

Habit: Culms up to 75 cm high.

Habitat: Disturbed soil.

Distribution in RGS: Weedy over the whole of the cultivated area.

General distribution: Native to South Africa, very common in South Brazil.

TRIODIA Brown

Triodia brasiliensis (Stedud.) Lindman, Ekman, Ark. f. Bot. 11,n.4

(1912) 40; Herter, Flor. Urug. (1930) 38. — *Uralepis brasiliensis* Stedud., Syn. () 218.

25591 RGS, Fazenda Sta Cecília p. S. Gabriel, SW, in campestribus siccis alte graminosis, 1.1944, fr., det. Swallen.

38344 RGS, Esteio p. S. Leopoldo, NEL, in campestribus dense graminosis, 24.11.1948, fr., det. Swallen.

49153 RGS, ibidem, 20.11.1950, fr., det. Beetle.

57237 RGS, ibidem, 11.11.1955, fl.

58973 RGS, Estação de Silvicultura p. Sta Maria, CN, in alte graminosis, 25.1.1956, fr., O. Camargo.

60336 RGS, ibidem, 8.4.1956, fr., leg. O. camargo 582.

Habit: Dense bunches, up to 75 cm high.

Habitat: Dry Campos.

Distribution in RGS: Widely scattered: western and northeastern Lowlands.

General distribution: Southern Brazil.

TRIPOGON Roth ap. R. et S.

Tripogon spicatus (Nees) Ekman, Ark. f. Bot. 11, n.4 (1912) 36. — *Diplachne spicata* (Nees) Döll, FB 2,2 (1878) 157.

36435 RGS, Cambará p. S. Francisco de Paula, NEH, in campestribus subsiccis, 2.1948, fr., det. swallen.

Habit: Erect, culms 30 cm high.

Habitat: More or less dry Campos.

Distribution in RGS: Northeastern Highlands.

General distribution: From Piauí to RGS.

SPECIES CULTIVATED OR ESCAPED (22).

Arundo donax L.: 988, 60987.

Avena fatua L.: 57230.

sativa L.: 1090,

sterilis L.: 2720.

strigosa Schreber: 29053.

Bambusa arundinacea Retz: 742, 37020, 58931, 59003, 59679.

Briza maxima L.: 2520, 6633.

- Coix lacrima-jobi** L.: 872, 42087, 47221, 63676.
Hordeum vulgare L.: 1092.
Euchlaena mexicana Schrajer: 859, 2272, 52946, 55629.
Lolium multiflorum Lam.: 2730, 29293, 32685, 42306, 57221, 57354,
 59735, 63174.
 perenne L.: 845, 2906, 6990, 35535.
Oryza sativa L.: 3075, 3076, 3077, 59026, 60922, 62998.
Panicum miliaceum L.: 1937.
Pennisetum clandestinum Hochst.: 9599.
Phyllostachys aurea A. et C. Riviere: 41557, 47101.
Saccharum officinarum L.: 60369.
Sorghum dochna (Forsk.) Snowden: 1414, 41558, 53108.
 halepense (L.) Pers.: 60813.
 saccharatum (L.) Moench: 60812.
 vulgare Pers.: 876, 32575.
Zea mays L.: 7006, 33133.

SPECIES CITED FOR RGS, BUT NOT IN MY HERBARIUM (122).

- Agrostis pectinana** Hack. et Arech.: Araujo, det. Parodi.
Andropogon altus Hitchc.: Araujo, det. Hitchc.
 exaratus Hack.: Malme, det. Ekman.
 plumiger Ekman: Malme, det. Ekman.
 salzmanni (Trin.) Hack: Araujo, det. Parodi.
 semiberbis (Nees) Kunth: Dutra, det. Dutra.
Aristida complanata Trin.: Döll, Lindman (Parodi, Swallen: A. laevis
 Kunth)
 macrophylla Hack.: Tupanciretan.
 megapotamica Spreng.: Parodi, Asplund.
 murina Cav.: Araujo, det. Parodi.
 recurvata H.B.K.: Lindman.

Axonopus arcatus Mez: S. Leopoldo, P. Alegre, Soledade, Rio Pardo.

- argentinus Parodi: Tupanciretan, Parodi.
 brasiliensis (Spreng.) Kulm.: Rambo, det. Swallen (lost).
 fissifolius (Raddi) Kulm.: Rambo, det. Parodi (lost).
 furcatus (Flügge) Hitchc.: Araujo, det. Parodi.
 iridaceus Mez: Araujo, det. Parodi.
 otusifolius (Raddi) Chase: Rambo, det. Swallen (lost).
 scoparius (Flügge) Hitchc.: Rambo, det. Parodi (lost).
Bouteloua megapotamica (Spreng.) O.K.: Araujo, det. Parodi.

Briza erecta Lam.: Malme.

lindmani Ekman: P. Alegre, Lindman.

sellowii Nees: Rambo, det. Parodi (lost.).

Bromidium anomalum Döll: RGS, Lindman

Chloris retusa Lagasca: Rambo, det. Parodi (lost.).

Chusquea juergensii Hack.: Rio Pardo.

phacelolophora Pilger: Cruz Alta.

Danthonia breviseta Hack.: Rambo, det. Parodi (lost.).

montana Döll: Rambo, det. Parodi (lost.).

Deschampsia juergensii Hackel: Soledade.

Digitaria connivens (Trin.) Mez: Malme, Rio Grande.

cuyabensis (Trin.) Parodi: Dutra.

lanuginosa (Nees) Henr.: S. Leopoldo.

lugens Nees: Araujo, det. Parodi.

pallens Parodi: Araujo, det. Parodi.

pedicellaris (Trin.) Prain: Araujo, det. Parodi.

phaeothrix (Trin.) Parodi: Rambo, det. Parodi (lost.).

Echinochloa implexa (Poir.) Chase: Rambo, det. Parodi (lost.).

Elionurus bisetosus Hack. et Lindman.: Rio Grande.

viridulus Hack.: Araujo, det. Parodi.

Eragrostis capillaris Steud.: Lindman, RGS.

monandra Hack: Soledade.

perennis Döll: Cruz Alta.

purpurascens Nees: RGS, Lindman.

rojasii Hack.: Araujo, det. Chase.

Eriochloa montevideensis (Nees) Gris.: Araujo, det. Parodi.

Eriochrysis warminigniana (Hack.) Pilger: Rambo, det. Swallen (lost.).

Guadua lindmanii Camus: Silveira Martins.

tomentosa Hack. et Lindman: Ijuí.

Gymnopogon haumanii Parodi: Araujo, det. Parodi.

radiatus (L.) Parodi: Araujo, det. Parodi.

spicatus (Spreng.) Kze: Araujo, det. Parodi.

Hordeum stenostachys Godr.: Araujo, det. Parodi.

Hymnachne donacifolia (Raddi) Chase: Rambo, det. Parodi, Swallen (lost.).

Ichnanthus candidans (Nees) Döll: S. Leopoldo.

leiocarpus Kunth: Rio Pardo, S. Leopoldo.

nemoralis (Schrad.) Hitchc. et Chase: S. Leopoldo, Dutra.

panicaoides Beauv.: Rambo, det. Parodi cum?

ruprechti Döll: Rambo, det. Swallen cum?

tenuis (Presl) Hitchc. et Chase: Rambo, det. Swallen cum?

Imperata tenuis Hack.: S. Leopoldo.

Melica papilionacea L.: Livramento.

Merostachys multiramea Hack.: Rio Pardo, Valley of the Taquary.

Panicum appressum Lam.: Dutra.

campestre Nees: Rambo, det. Parodi (lost).

apillare L.: RGS, Lindman.

dichotomiflorum Michx.: Araujo, det. Parodi.

glabripes Döll: Rio Pardo, S. Leopoldo; Dutra.

gracilipes Hack.: Fazenda Leitão, Rio Pardo.

junceum Nees: P. Alegre; Dutra.

parvifolium Lam.: Rambo, det. Parodi (lost).

plantagineum Linck: Dutra.

prionitis Nees: Rambo, det. Parodi (lost).

proliferum Lam.: Dutra.

rude Nees: Rambo, det. Swallen (lost).

stenophyllum Hack.: Rambo, det. Swallen (lost).

uncinatum Raddi: RGS, Lindman.

Paspalum alnum Chase: Araujo, det. Parodi.

brunneum Mez: S. Leopoldo.

dvincenzzii Parodi: Araujo, det. Parodi.

durifolium Mez: S. Leopoldo.

equitans Mez: Araujo, det. Parodi.

erianthoides Lindman: Araujo, det. Parodi.

falcatum Nees: Venâncio Aires.

glaucescens Hack.: Dutra.

hydrophilum Henr.: Montenegro.

indecorum Mez, Araujo, det. Parodi.

intermedium Munro: Rambo, det. Parodi (lost).

modestum Mez: Araujo, det. Parodi.

platycaulon Poir.: Malme, Lindman.

rufum Nees: Rambo, det. Swallen (lost).

stellatum H.B.K.: Araujo, det. Chase.

uruguayense Arech.: Bornmüller.

virgatum L.: RGS, Lindman.

warmingii: Rambo, det. Parodi (lost).

Phalaris arundinacea L.: Araujo, det. Parodi.

mimor Retz: Araujo, det. Parodi.

platensis Parodi: Araujo, det. Parodi.

Piptochaetium ovatum (Trin. et Ruper.) Desv.: Araujo, det. Parodi.

setosum Arech.: Araujo, Parodi.

Polypogon maritimus Willd.: Rio Grande.

Rottboellia balansae Hack.: Sto Angelo.

Sacciolepis myuros (Lam.) Chase: Araujo, det. Parodi.

Setaria flava Kunth: Dutra.

globulifera (Steud.) Gris.: RGS, Rambo (lost).

imberbis R. et S.: RGS, Rambo, det. Swallen (lost).

- paucifolia* (Morong) Londm.: Sta Angelo.
- Sorghastrum albescens** (Hack.): Bom Jesus.
- scaberrimum* (Nees) Herter: Rambo, det. Swallen (lost).
- Spartona montevidensis** Arech.: RGS.
- stricta* (Ait.) Roth, Malme, Rio Grande.
- Sporobolus brasiliensis** (Raddi) Hack.: Bom Jesus, Rio Pardo.
- indicus* (L.) R. Br.: Lindman, Malme.
- multicaulis* Hack.: det. Parodi.
- multinodis* Hack.: S. Leopoldo, Rio Pardo.
- ramosissimus* Kunth: Lindman, RGS.
- Stipa bavioensis** Speg.: Rambo, det. Parodi (lost).
- charruana* Arech.: Araujo, det. Parodi.
- filicaumis* Delil.: Rambo, det. Parodi (lost).
- neesiana* Trin. et Rupr.: Araujo, det. Parodi.
- papposa* Nees: Araujo, det. Parodi.
- Triodia hackelii** Arech.: Araujo, det. Parodi.

LITERATURE (local only).

Araujo, A. A. de

- 1942 Gramíneas úteis da Flora Riograndense. — Secretaria de Agricultura, Bol. n.100, P. Alegre.
- 1948 O Gramado. Disclimax da Vegetação Campestre. — Secretaria da Agricultura, Série C, P. Alegre.
- Arechavaleta, J.**
- 1894 Las Gramíneas uruguayas. — Montevideo 1874.
- 1897 Idem. — An. Mus. Hist. Nat. Montevideo 1 (1897) 29-271; 213-556.
- 1903 Idem. — Ibidem, 4 (1903) 87-121.

Barreto I.L.

- 1954 Las Especies afines de *Paspalum virgatum* en la América del Sur. — Rev. Arg. Agron. 21, n.3 (1954) 125-142.
- 1956 Las especies afines de *Paspalum plicatulum* en el Rio do Sul (Brasil). — Rev. Arg. Agron. 23, n.2 (1956) 53-70.

Bertoni, M. S.

- 1918 Graminaceas de las regiones florestales del Alto Paraná. — An. Ci. Parag. 2. ser., 6 de Bot. (1918) 142-166.
- Bornmüller, A.**
- 1934 Gramineae. — Florula Riograndensis, Rev. Sudam. Bot. 1 (1934) 132-133.

Cabrera, A. L.

- 1943 Gramineae – Manual de la flora de los alrededores de Buenos Aires (1953)

Dutra, J.

- ? A taquara-assú, *Bambusa riograndensis* n. sp. – Rev. Agric. do RGS ano 6 () 102-104, 1 pl.
1938 Les Bambuses de Rio Grande do Sul. – Rev. Sudam. Bot. 5 (1938) 145-152, 3 fig.

Ekman, E. L.

- 1911 Neue brasiliische Gräser. – Ark. f. Bot. 10, n.17 (1911).
1912 Beiträge zur Gramineenflora von Misiones. – Ark. f. Bot. 11, n.4 (1912).
1913 Die Gräser des brasiliischen Staates Paraná. – Ark. f. Bot. 13, n.10 (1913).

Hackel, E.

- 1883 Gramineae. – Flora Brasiliensis (FB) 2,3 (1883) 245-326.
1885 Andropogoneae novae. – Flora 7 (1885) 115-128; Flora 8 (1885) 131-145.
1889 Andropogonae. – DC. Monogr. Phan. 6 (1889) 1-716.
1904 Gramineae. Chod. et Hassl. Plant. Hasslerianae. – Bull. Herb. Boiss. 2 série, 4 (1904) 265-282.

Henrard, J.T.

- 1929- 1933 A monograph of the genus *Aristida* l. – Med. Herb. Leiden, n. 58, (1929-1933).

Herter, W.

- 1925 Treinta y tres gramíneas nuevas para el país. – Com Herb. Osten Montevideo, n.1 (1924) 1-13; 29-31.
1929 Las Gramíneas de la República O. del Uruguay. – An. Mus. Hist. Nat. Montevideo 2. ser. 3 (1929) 43-56.
1930 Gramineae. Florula Uruguayensis (1930) 26-39.

Lindman, C.A.M.

- 1900a Beiträge zur Gramineenflora Südamerikas. – Kungl. Sv. Vet. Akad. Handl. 34, n.6 (1900) 1-52.
1900b Vegetationen in Rio Grande do Sul. – Stockholm 1900.

Malme, G.O.A.

- 1936 Zur Kenntnis der Phanerogamenflora des Sandgebietes im Süden von Rio Grande do Sul. – Sv. Bot. Tidskr. 30 (1936) 9-11

Mez, C.

- 1917a Novae Species Panicearum. – Notizbl. Dahlem 7 (1917) 45-78.
1917b Generis Paspali Species novae. – Fedde Rept. 15 (1917) 27-32; 60-76.

Osten, C. and Herter. W.

- 1927-29 Las Gramíneas de la República O. del Uruguay. — Plantae Uruguayenses 5. — An.Mus.Hist.Nat.Mont. 2. serie (1927) 301-319; ibidem 3 (1929) 43-56.

Parodi, L. R.

- 1919 Las Chlorídeas de la República Argentina. — Rev. Fac. Agr. Vet. B. Aires 2 (1919) 233-335.
- 1920 Notas sobre las especies de Briza de la Flora Argentina. — Rev. Fac. Agron. Vet. B. Aires 3 (1920) 113-138.
- 1922 Las Gramíneas de la región de Concordia (Prov. de Entre Ríos). — Rev. Fac. Agron. Veter. B. Aires 4 (1922) 24-102.
- 1925a Gramineas bonaerenses. — Rev. Cent. Est. Agr. B. Aires (1925) 120, 17...; 121, 111...
- 1925b Las Gramíneas de género *Pennisetum* de la Flora argentina. — An.Mus.Ci.Nat.B.Aires 32 (1925) 501-526.
- 1928a Revisión de las Gramíneas argentinas del género *Sporobolus*. — Rev.Fac.Agron.Vet. B. Aires 6 (1928) 115-168.
- 1928b Sinopsis de las Gramíneas argentinas del género *Muehlenbergia*. — Physis 9 (1928) 205-222.
- 1936a Contribución al conocimiento de las especies del género *Poa* en la Flora Uruguaya. — Rev. Arg. Agron. 3 (1936) 133-152.
- 1936b Las Bambuseas indígenas en la Mesopotamia argentina. — Rev.Arg.Agr. 3 (1936) 229-244.
- 1937 Contribución al estudio de las Gramíneas de género *Paspalum* de la Flora uruguaya. — Rev.Mus.La Plata (1937) 211-250.
- 1941 Estudio sobre el Género *Chusquea* en la Argentina. — Rev. Arg. Agron. 8, n.4 (1941) 331-345.
- 1944 Revisión de las Gramíneas americanas del género *Piptochaetium*. — Rev. del Mus. de la Plata, nueva ser. secc. bot. 6 (1944) 213-310.
- 1946a Las especies de *Sorghum* cultivadas en la Argentina. — Rev.Arg.Agron. 13,n.1 (1946) 1-32.
- 1946b *Agrostis ramboi* Par., nueva especies. — Bol.Soc.Arg.Botánica 1, n.2 (1946) 119-122.
- 1946c Gramíneas bonaerenses, 4. ed. — Centro Estudiantes Fac. Agron. B. Aires, 1946.
- 1947 Las Gramíneas del género *Bromus* adventicias en la Argentina. — Rev. Arg. Agron. 14, n.1 (1947) 1-19.
- 1949a Las Gramíneas sudamericanas del género *Deschampsia*. — Darwiniana 8 (1949) 415-475.
- 1949b Las especies afines a *Camamagrostis viridi-flavescens*. — Rev. Arg. Agron. 16,n.2 (1949) 61-77.

Reitz, R.

1956 Gramineae. Manipulus Monocotyledonearum Catarinen-sium. — *Sellowia* 7 (1956) 96-109.

INDEX.

Agrostis hygrometrica Nees	9
<i>juergensii</i> Hack	9
<i>koelerioides</i> E. Desv.	9
<i>longiberbis</i> Hack.	10
<i>montevidensis</i> Spreng.	10
<i>ramboi</i> Parodi	11
Amphibromus rubrivalvis (Trin.) Swallen	11
Andropogon arenarius Hack.	12
<i>bicornis</i> L.	12
<i>condensatus</i> H.B.K.	12
<i>consanguineus</i> Kunth	13
<i>glaucuscens</i> H.B.K.	13
<i>hassleri</i> Hack.	14
<i>hypogynus</i> Hack.	14
<i>imberbis</i> Hack.	14
<i>lateralis</i> Nees	14
<i>leucostachyus</i> H.B.K.	17
<i>lindmanii</i> Hack.	17
<i>macrothrix</i> Trin.	18
<i>paniculatus</i> Kunth	18
<i>saccharoides</i> Swartz	20
<i>selloanus</i> Hack.	20
<i>tener</i> (Nees) Kunth	21
<i>ternatus</i> (Spreng.) Nees	22
<i>virgatus</i> Desv.	22
Anthoxanthum odoratum L.	23
Aristida altissima Arech.	24
<i>circinnalis</i> Lidm.	25
<i>implexa</i> Trin.	25
<i>laevis</i> Kunth	26
<i>pallens</i> Cav.	27
<i>teretifolia</i> Arech.	28
<i>uruguayensis</i> Henr.	28
<i>venustula</i> Arech.	28
Arthrostylidium trinii (Rupr.) Munro	29
Arundinaria ulei Hack.	29

Arundinella hispida (Willd.) O.K.	30
xonopus affinis Chase	30
brasiliensis (Spreng.) Kuhl	31
compressus (Swartz) Beauv.	32
suffultus (Mikan) Parodi	33
ulei Hack.	34
Brachiaria plantaginea (Linck) Hitchc.	35
Briza calotheca (Trin.) Hack.	35
hackelii (Lindm.) Ekm.	36
juergensii Hack.	36
lamarckiana Nees	37
minor L.	37
neesii Döll	38
poaemorpha (Presl) Henr.	38
rotundata Hack.	39
rufa (Hook et Arn.) Steud.	39
stricta (Presl) Steud.	40
subaristata Lam.	41
triloba Nees	42
uniolae Nees	43
Bromus auleticus (Trin.) ex Nees	44
brachyanthera Döll	45
catharticus Vahl	46
Calamagrostis armata (Döll) Parodi	47
beyrichiana Nes	47
longearistata (Wedd.) Hack.	47
montevidensis Nees	48
viridi-flavescens (Poir.) Steud.	49
Cenchrus echinatus L.	50
myosuroides H.B.K.	51
pauciflorus Benth.	51
Chaetotropis chilensis Kunth.	52
Chloris argentina (Hack.) Lillo et Parodi	52
berroi Arech.	53
canterai Arech.	53
caribaea Spreng.	53
ciliata Swartz	54
distichophylla Lag.	54
dusenii Ekman	55
gayana Kunth	56
polydactyla Swartz	56
pycnothrix Trin.	56
uliginosa Hack.	57
Chusquea meyeriana Rupr.	57

<i>ramosissima</i> Lindman	58
<i>tenella</i> Nees	58
Cortaderia rudiuscula Stapf	59
<i>seloana</i> (Schult.) Aschers. et Graebner	59
Cynodon dactylon (L.) Pers.	60
Danthonia cirrhata Hack.	61
<i>dusenii</i> Ekman	62
<i>secundiflora</i> Presl	62
<i>tenuifolia</i> Döll	62
Deschampsia caespitosa (L.) Beauv.	63
Digitaria aequiglumis (Hack. et Arech.) Parodi	63
<i>horizontalis</i> Willd.	63
<i>sacchariflora</i> (Raddi) Henr.	64
<i>sanguinalis</i> (L.) Scop.	64
<i>violascens</i> Link	65
Echinochloa colonum (L.) Link	66
<i>crus-galli</i> (L.) Beauv.	66
<i>crus-pavonis</i> (H.B.K.) Schultes	66
<i>spectabilis</i> (Nees) Link	68
Eleusine indica (L.) Gaertn.	68
<i>tristachya</i> (Lam.) Kunth	69
Elionurus adustus (Trin.) Ekman	70
<i>candidus</i> (Trin.) Hack.	70
<i>rostratus</i> Nees	71
Eragrostis airoides Nees	71
<i>articulata</i> Nees	72
<i>bahiensis</i> (Schrad.) Schult.	72
<i>brasiliensis</i> Hack.	74
<i>cilianensis</i> (All.) Link	74
<i>ciliaris</i> (L.) Link	74
<i>expansa</i> Link	74
<i>flaccida</i> Lindman	75
<i>hypnoides</i> (Lam.) B.S.P.	76
<i>lugens</i> Nees	76
<i>megapotamica</i> (Spreng.) R. et Schl.	77
<i>neesii</i> Trin.	77
<i>pilosa</i> (L.) Beauv.	79
<i>polytricha</i> Nees	79
<i>trichocolea</i> Hack. et Ar.	80
<i>virescens</i> Presl	80
Erianthemum bulbosum Parodi	80
Erianthus angustifolius Nees	80
<i>asper</i> Nees	81
<i>trinii</i> Hack.	82

Eriochloa polystachya H.B.K.	82
<i>punctata</i> (L.) Hamilt.	83
Eriochrysis cayennensis Beauv.	84
Festuca ampliflora Döll	85
<i>australis</i> Nees	85
<i>dertonensis</i> (All.) Aschers. et Graebn.)	85
<i>fimbriata</i> Nees	86
<i>myurs</i> L.	86
<i>ulochaeta</i> (Nees) Döll	87
Guadua trinii Ness	88
Gymnopogon biflorus Pilger	88
Heteropogon villosus Nees	89
Holcus lanatus L.	89
Hordeum compressum Griseb.	90
<i>pusillum</i> Nutt.	90
Hymenachne amplexicaulis (Rudge) Nees.	91
Hyparrhenia hirta (L.) Stapf	91
Ichnanthus pallens (Swartz) Munro	91
Imperata brasiliensis Trin.	93
<i>contracta</i> (H.B.K.) Hitchc.	93
<i>tenuis</i> Hack.	93
Ischaemum urvilleanum Kunth	93
Koeleria phleoides (Vill.) Pers.	94
Lasiacis divaricata (L.) Hitzc.	95
Leersia hexandra Swartz	95
Leptocoryphium lanatum (H.B.K.) Nees	96
Luziola peruviana Gmel.	97
Manisuris loricata (Trin.) Kuntze	98
Melica aurantiaca Lam.	98
<i>sarmentosa</i> Nees	99
Melinis minutiflora Beauv.	100
Merostachys burchellii Munro.	101
Microchloa indica (L.) O.K.	102
Muehlenbergia schreberi Gmel.	102
Olyra humidis Nees	103
<i>latifolia</i> L.	103
Oplismenus hirtellus (L.) Beauv.	104
<i>setarius</i> (Lam.) R. et Schult.	104
Oryza subulata Nees	105
Panicum anomalum Trin.	105
<i>aquaticum</i> Poir.	106
<i>aristellum</i> Döll	106
<i>bergii</i> Arech.	106

<i>boliviense</i> Hack.	107
<i>chloroticum</i> Nees	107
<i>cyanescens</i> Nees	108
<i>decipiens</i> Nees	108
<i>demissum</i> Trin.	109
<i>dusenii</i> Hack.	110
<i>elephantipes</i> Nees	110
<i>fultum</i> Hack.	110
<i>glutinosum</i> Swartz	111
<i>gouinii</i> Fourn.	111
<i>grumosum</i> Nees	112
<i>gynerioides</i> (Gris.) Pilger	112
<i>hebotes</i> Trin.	112
<i>helobium</i> Mez	113
<i>latiglume</i> Döll	114
<i>laxum</i> Swartz	114
<i>maximum</i> Jacq.	115
<i>milioides</i> Nees	115
<i>millegrana</i> Poir.	116
<i>missionum</i> Mez	116
<i>numidianum</i> Lam.	117
<i>olyroides</i> H.B.K.	117
<i>ovuliferum</i> Trin.	117
<i>paludivagum</i> Hitchc. et Chase	118
<i>pantrichum</i> Hack.	118
<i>pilosum</i> Swartz	118
<i>polycladum</i> Ekman	119
<i>polygonatum</i> Schrad.	119
<i>procurrens</i> Nees	119
<i>racemosum</i> Spreng.	120
<i>repens</i> L.	120
<i>rhizogonum</i> Hack.	120
<i>rivulare</i> Trin.	121
<i>sabulorum</i> Lam.	121
<i>schenckii</i> Hack.	123
<i>secundum</i> Trin.	123
<i>sellowii</i> Nees	124
<i>stigmosum</i> Trin.	124
<i>stoloniferum</i> Poir.	125
<i>subjuncuum</i> Ekman	125
<i>superatum</i> Hack.	126
<i>trichantium</i> Nees	126
<i>tricholaenoides</i> Steud.	126
<i>Paspalum acuminatum</i> Raddi	127

<i>arechavaletae</i> Hack.	127
<i>arenarium</i> Schrad.	127
<i>conjugatum</i> Berg	128
<i>conspersum</i> Schrad.	129
<i>corcovadense</i> Raddi	129
<i>cromyorrhizon</i> Trin.	130
<i>dilatatum</i> Poir.	130
<i>distichum</i> L.	131
<i>exaltatum</i> Presl	131
<i>guaraniticum</i> Parodi	132
<i>guenoarum</i> Arech.	132
<i>haumanii</i> Parodi	132
<i>hyalinum</i> Nees	133
<i>inaequivalve</i> Raddi	133
<i>juergensii</i> Hack.	134
<i>maculosum</i> Trin.	134
<i>mandiocanum</i> Trin.	134
<i>multiflorum</i> Döll	135
<i>nicorae</i> Parodi	136
<i>notatum</i> Flügge	136
<i>orbiculatum</i> Poir.	138
<i>paniculatum</i> L.	138
<i>plicatulum</i> Michx.	139
<i>plumosum</i>	141
<i>polyphyllum</i> Nees	141
<i>pumilum</i> Nees	142
<i>quadrifarrium</i> Lam.	143
<i>regnellii</i> Mez	143
<i>repens</i> Berg	144
<i>rojasii</i> Hack.	144
<i>umbrosum</i> Trin.	144
<i>urvillei</i> Steud.	144
<i>vaginatum</i> Swartz	146
<i>yaguaronense</i> Henr.	146
Pennisetum latifolium Spreng.	147
<i>purpureum</i> Schumah	147
Phalaris angusta Nees	148
Pharus glaber H.B.K.	149
Piptochaetium confusum Parodi	150
<i>lasianthum</i> Gris.	150
<i>montevidense</i> (Spreng.) Parodi	151
<i>panicoides</i> (Lam.) Desv.	152
<i>ruprechtianum</i> Desv.	152
<i>stipoides</i> (Trin. et Rupr.) Hack.	153

Poa annua L.	153
bonariensis (Lam.) Kunth	154
bradei Pilger	154
lanigera Nees	154
montevidensis Arech.	155
trivialis L.	
Polypogon elongatus H.B.K.	156
monspeliensis (Döll) Desf.	156
Pseudechinolaena polystachya (H.B.K.) Stapf	157
Sacciolepis strumosa (Presl) Chase	158
viloides (Trin.) Chase	158
Scleropoa rigida (L.) Gris.	159
Setaria caespitosa Hack.	159
fiebrigii Herrmann	159
geniculata (Lam.) Beauv.	160
lachnea (Nees) Kunth	162
onurus (Willd.) Gris.	163
poiretiana (Schult.) Hitchc.	163
scandens Schrad.	164
vaginata Spreng.	164
verticillata (L.) Beauv.	165
vulpiseta (Lam.) Hitchc. et Chase	165
Sorghastrum agrostoides (Speg.) Hitchc.	165
minarum (Nees) Hitchc.	166
nutans (L.) Nees	166
parviflorum (Desv.) Hitchc. et Chase	166
setosum (Gris.) Hitchc.	167
stipoides (H.B.K.) Nash	167
Spartina ciliata Kunth	167
Sporobolus aeneus (Trin.) Kunth	168
eximus (Nees) Ekman	168
poretii (R. et Sch.) Hitchc.	168
pseudairoides Parodi	170
pyramidalis Beauv.	170
virginicus (L.) Kunth	170
Stenotaphrum secundatum (Walt.) O.K.	171
Stipa amethystina Steud.	172
filifolia Nees	172
juergensii Hack.	172
megapotamica Spreng.	172
melanosperma Presl.	173
nutans Hack.	174
sellowiana Nees	174
Trachypogon canescens Nees	174

Trichachne insularis (L.) Nees	175
Tricholaena rosea Nees	176
Triodia brasiliensis (Steud.) Lindman	176
Tripogon spicatus (Nees) Ekman	177
Species cultivated or escaped (22)	177
Species cited for Rio Grande do Sul, not in my Herbarium	178

11.9.1961.

PESQUISAS

PUBLICAÇÕES DE BOTÂNICA

1. **Die Auslese im Naturversuch** – B. Rambo, SJ. – Pesquisas 1, 1957, 131-219.
2. **Die Alte Südfloren in Brasilien** – B. Rambo, SJ. – Pesquisas 2, 1958, 177-198.
3. **An Historical Approach to Plant Evolution** – B. Rambo, SJ. – Pesquisas 2, 1958, 199-222.
4. **Uma coleção de pteridófitos do Rio Grande do Sul** – Aloysio Sehnem, SJ. – Pesquisas, 2, 1958, 223-229 e 6 est. fora do texto.
5. **Cyperaceae Riograndenses** – B. Rambo, SJ. – Pesquisas 3, 1959, 353-453.
6. **Towards the concept of the species in plant evolution** – B. Rambo, SJ. – Pesquisas 3, 1959, 455-493.
7. **Uma coleção de pteridófitos do Rio Grande do Sul, cont.** – Aloysio Sehnem, SJ. – Pesquisas 3, 1959, 495-576 e 5 est. fora do texto.
8. **Die Südgrenze des brasiliandischen Regenwaldes** – B. Rambo, SJ. – Pesquisas 1960, Bot. nr. 8; 41 pp.
9. **Euphorbiceae riograndenses** – B. Rambo, SJ. – Pesquisas 1960, Bot. nr. 9; 78 pp.
10. **Uma coleção de pteridófitos do Rio Grande do Sul IV** – Aloysio Sehnem, SJ. – Pesquisas 1960, Bot. nr. 10; 44 pp. e 5 est. fora do texto.
11. **Solaceae riograndenses** – B. Rambo, SJ. – Pesquisas 1961, Bot. nr. 11; 69 pp.
12. **Migration routes of the south Brazilian forest** – B. Rambo, SJ. – Pesquisas 1961, Bot. nr. 12; 54 pp.
13. **Uma coleção de pteridófitos do Rio Grande do Sul V** – Aloysio Sehnem, SJ. – Pesquisas 1961, Bot. nr. 13; 42 pp. e 10 est. fora do texto.
14. **Der Küstenwald in Rio Grande do Sul (Südbrasiliens)** – Roberto M. Klein – Pesquisas 1961, Bot. nr. 14; 39 pp. e 6 tab., 1 mapa fora do texto.
15. **Labiatae riograndenses** – B. Rambo, SJ. – Pesquisas 1962, Bot. nr. 15; 46 pp.
16. **Convolvulaceae riograndenses** – B. Rambo, SJ. – Pesquisas 1962, Bot. nr. 16; 31 pp.
17. **Umbelliferae riograndenses** – B. Rambo, SJ. – Pesquisas 1962, Bot. nr. 17; 39 pp.
18. **Rubiaceae riograndenses** – B. Rambo, SJ. – Pesquisas 1962, Bot. nr. 18; 76 pp.
19. **Observações sobre o prótalo de *Trichomanes pilosum* Raddi** – Aloysio Sehnem, SJ. – Pesquisas 1965, Bot. nº 19; 12 pp. 4 fig.
20. **Myrtaceae riograndenses** – B. Rambo, SJ. – Pesquisas 1965, Bot. nr. 20; 64 pp.
21. **Verbenaceae Riograndenses** – B. Rambo, SJ. – Pesquisas 1965, Bot. nr. 21; 62 pp.
22. **Melastomataceae Riograndenses** – B. Rambo, SJ. – Pesquisas 1966, Bot. nr. 22; 48 pp.
23. **Leguminosae Riograndenses** – B. Rambo, SJ. – Pesquisas 1966, Bot. nr. 23; 170 pp.
24. **Malvaceae Riograndenses** – B. Rambo, SJ. – Pesquisas 1967, Bot. nr. 24, 52 pp.
25. **Bromeliaceae Riograndenses** – B. Rambo, SJ. – Pesquisas 1967, Bot. nr. 25, 27 pp.
26. **Amarantaceae Riograndenses** – B. Rambo, SJ. – Pesquisas 1968, Bot. nr. 26, 30 pp.
27. **Musgos Sul-brasileiros** – Aloysio Sehnem, Pesquisas 1969, Bot. nr. 27; 33 pp. 5 Est.
28. **Musgos Sul-brasileiros II** – Aloysio Sehnem, Pesquisas 1970, Bot. nr. 28, 96 pp. 21 Est.
29. **Musgos Sul-brasileiros III** – Aloysio Sehnem, Pesquisas 1972, Bot. nr. 29, 70 pp.
30. **Musgos Sul-brasileiros IV** – Aloysio Sehnem, Pesquisas 1976, Bot. nr. 30, 79 pp.
31. **As Filicíneas do sul do Brasil, sua Distribuição geográfica, sua Ecologia e suas Rotas de Migração** – Aloysio Sehnem, Pesquisas 1977, Bot. nr. 31, 108 pp.
32. **Musgos Sul-brasileiros V** – Aloysio Sehnem, Pesquisas 1978, Bot. nr. 32, 170 pp.
33. **Musgos Sul-brasileiros VI** – Aloysio Sehnem, Pesquisas 1979, Bot. nr. 33, 149 pp.
34. **Musgos Sul-Brasileiros VII** – A. Sehnem, S.J., Pesquisas 1980, Botânica nº 34, 121 pp.
35. **Contribuição ao estudo dos fungos agaricais da Mata Nativa de Araucaria angustifolia (Bertol.) O. Kze. da floresta nacional de São Francisco de Paula, Rio Grande do Sul** – Antonio Batista Pereira, Pesquisas 1984, Botânica nº 35, 73 pp.

Composto e Impresso na Gráfica UNISINOS